

[国内版](#)[国际版](#)

Comparison of cytokine and phosphoprotein profiles in idiopathic and Crohn

[All](#)[Images](#)[Videos](#)[翻译成中文](#)[关闭取词](#)

9,770 Results

Any time ▾

Management of perianal fistulas in Crohn's disease: An up ...

www.ncbi.nlm.nih.gov › ... › v.21(5); 2015 Feb 7

Feb 07, 2015 · INTRODUCTION. **Perianal** fistulas are a very disabling manifestation and source of morbidity for **Crohn's** disease (CD) patients. Population-based studies showed that the cumulative incidence of **perianal** fistulas in CD range from 23% to 38%[1,2].Disease location, age at diagnosis, **fistula** type, presence or absence of abscesses and intestinal strictures may influence the natural ...

Cited by: 29

Author: Manuela Marzo, Carla Felice, Daniela Pu...

Publish Year: 2015

Review article: medical, surgical and radiological ...

onlinelibrary.wiley.com/doi/10.1111/j.1365-2036.2010.04486.x/full

Tacrolimus. **Tacrolimus** was assessed in a **placebo-controlled trial** in which it was given orally for 10 weeks in 42 **Crohn's anal fistula** patients. There was significant clinical improvement in the tacrolimus group (45%) compared with **placebo** (9%) but no significant improvement in **fistula** ...

Published in: **Alimentary Pharmacology & Therapeutics** - 2011

Authors: P Tozer · David Burling · A Gupta · Robin K S Phillips · Ailsa Hart

Affiliation: **Imperial College London**

About: **Radiology** · **Fistula** · **Surgery**

Predictors of response to infliximab in patients with ...

<https://www.sciencedirect.com/science/article/pii/S0016508502001610>

Background & Aims: Identifying predictors of response to infliximab in **Crohn's** disease may lead to better selection of patients for this therapy.Methods: One hundred patients with either inflammatory or fistulous **Crohn's** disease and at least 3 months of follow-up after infliximab infusion were evaluated. Clinical response and duration of response were the primary outcome measures.

Cited by: 322

Author: Mansour A. Parsi, Jean-Paul Achkar, Sc...

Publish Year: 2002

Current treatment of rectovaginal fistula in Crohn's disease

https://www.researchgate.net/publication/50907865_Current...

Objective To **compare** the healing rate after several types of surgical repair of rectovaginal **fistula** (RVF) in **Crohn's** disease, and to identify factors predicting a successful outcome.Patients ...

검색결과 약 3개 (0.48초)

Comparison of cytokine and phosphoprotein profiles in idiopathic and Crohn's disease-related perianal fistula에 대한 학술자료

Hyperactivated B cells in human inflammatory bowel ... - Noronha - 64회 인용

Serum Cytokine Profiles in Children with Crohn's Disease

<https://www.ncbi.nlm.nih.gov/articles/PMC5187493> ▼ 이 페이지 번역하기

E Vasilyeva 저술 - 2016 - 5회 인용 - 관련 학술자료

2016. 12. 13. - **Crohn's disease** (CD) is a form of inflammatory bowel **disease** (IBD) affecting the ... pain, distension, bloating, vomiting, **perianal fistulae**, and abscesses. ... However, little is known about **differences** in serum **cytokine profiles** ... Potentially this information could suggest biomarkers **related** to the age of onset.

누락된 검색어: phosphoprotein idiopathic

관련 질문

Does Crohn's disease cause fistulas?



Does fistula mean Crohn's?



What is fistulizing Crohn's disease?



2 Name of Journal: *World Journal of Gastrointestinal Pathophysiology*
Manuscript NO: 48183
Manuscript Type: ORIGINAL ARTICLE

Observational Study

Comparison of cytokine and phosphoprotein profiles in idiopathic and Crohn's disease-related perianal fistula

Haddow JB *et al.* Perianal fistula immunopathology

James B Haddow, Omar Musbahi, Thomas T MacDonald, Charles H Knowles

Abstract

BACKGROUND

Perianal fistulae are either primary (idiopathic) or secondary [commonly associated with Crohn's disease, (CD)]. It is assumed, although not proven, that the pathophysiology differs.

AIM

To systematically compare the clinical phenotypes, cytokine and phosphoprotein profiles of idiopathic and CD-related perianal fistulae.

METHODS

Sixty-one patients undergoing surgery for perianal fistula were prospectively recruited (48 idiopathic, 13 CD) into a cohort study. Clinical data, including the Perineal Disease Activity Index (PDAI) and EQ-5D-5L were collected. Biopsies of the fistula tract, granulation tissue, internal opening mucosa and rectal mucosa were obtained at surgery. Concentrations of 30 cytokines and 39 phosphoproteins were

Match Overview

1	Crossref 33 words Rothlin, Carla V., Jonathan A. Leighton, and Sourav Ghosh. " Tyro3, Axl, and Mertk Receptor Signaling in Inflammatory B ...	1%
2	Internet 32 words crawled on 28-Aug-2019 f6publishing.blob.core.windows.net	1%
3	Internet 14 words crawled on 04-Dec-2008 ajp.amjpathol.org	<1%
4	Internet 14 words crawled on 06-Oct-2019 www.tandfonline.com	<1%



All

Images

Videos

关闭取词

15,100 Results

Any time ▼

Management of perianal fistulas in Crohn's disease: An up ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4316082>

Feb 07, 2015 · INTRODUCTION. **Perianal** fistulas are a very disabling manifestation and source of morbidity for **Crohn's** disease (CD) patients. Population-based studies showed that the cumulative incidence of **perianal** fistulas in CD range from 23% to 38%[1,2].Disease location, age at diagnosis, **fistula** type, presence or absence of abscesses and intestinal strictures may influence the natural ...

Cited by: 34

Author: Manuela Marzo, Carla Felice, Daniela Pu...

Publish Year: 2015

British Library EThOS: Studies in the aetiology ...

<https://ethos.bl.uk/OrderDetails.do?uin=uk.bl.ethos.766107> ▼

Over 12,000 clinical and 23,500 laboratory measurements were made. Results: We found no clear differences in the clinical characteristics or the overall **cytokine** or **phosphoprotein profiles** between **idiopathic** and **Crohn's perianal** fistulae. We also found no differences in the DCE-MRI measurements between the groups.

Author: James Boyd Haddow

Publish Year: 2018

Perianal Crohn's disease and hidradenitis suppurativa: a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4511252>

Jul 22, 2015 · **Crohn's disease** (CD) and Hidradenitis suppurativa (HS) are both **chronic inflammatory diseases**. The **pathogenesis** of these **diseases** is multifactorial, due to the interaction of genetic and environmental factors leading to a deregulated local immune response where T ...

Cited by: 12

Author: Francesco Giudici, Laura Maggi, Raffaell...

Publish Year: 2015

UpToDate

www.uptodate.com/contents/search?search=crohns

Similar **cytokine profiles** in children with juvenile **idiopathic** arthritis ... Primary immunodeficiencies: Classification ... dysregulation with colitis – Patients develop severe, early-onset inflammatory bowel disease (**Crohn** disease or ulcerative colitis), even as early as infancy.