



国内版

国际版

Multi-institutional retrospective analysis of FOLFIRI in patients with advanced biliar



All

Images

Videos

翻译成中文

关闭取词

26,800 Results

Any time ▼

## Second-line chemotherapy in advanced biliary cancer ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4689003>

Dec 23, 2015 · Types of second-line chemotherapy used were the following: FOLFOX or XELOX or 5-fluorouracil plus cisplatin in 128 patients, FOLFIRI in 75 patients (in 13 of whom in combination with bevacizumab), 5-fluorouracil or capecitabine monotherapy in 39 patients, gemcitabine plus cisplatin in 17 patients and other regimens in 66 patients.

Cited by: 15

Author: Lorenzo Fornaro, Caterina Vivaldi, Stefan...

Publish Year: 2015

## Study of Nivolumab in Patients With Advanced Refractory ...

<https://clinicaltrials.gov/ct2/show/NCT02829918> ▼

Jul 12, 2016 · Study of Nivolumab in Patients With Advanced Refractory Biliary Tract Cancers. Prior therapy with anti-PD-1, anti-PD-L1, anti-PD-L2, anti-CD137, or anti-CTLA-4 antibody (including ipilimumab or any other antibody or drug specifically targeting T-cell costimulation ...

## A multicenter randomized phase II study of nivolumab in ...

[https://ascopubs.org/doi/abs/10.1200/JCO.2018.36.4\\_suppl.TPS541](https://ascopubs.org/doi/abs/10.1200/JCO.2018.36.4_suppl.TPS541)

A multicenter randomized phase II study of nivolumab in combination with gemcitabine/cisplatin or ipilimumab as first-line therapy for patients with advanced unresectable biliary tract cancer.

Cited by: 1

Author: Vaibhav Sahai, Kent A. Griffith, Mark Zalu...

Publish Year: 2018

## Advanced biliary tract carcinomas: a retrospective ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4236575>

Background. The aim of this retrospective study was to evaluate the OS, PFS, and safety in patients with aBTCs treated in first line chemotherapy by modified one day GEMOX or Gem/Carb regimen in our institution comprising three different hospitals.

Cited by: 9

Author: Frédéric Fiteni, Marine Jary, Franck Monn...

Publish Year: 2014

## Patients with recurrent biliary tract cancer have a better ...

[https://www.researchgate.net/publication/242018469\\_Patients\\_with\\_recurrent\\_biliary...](https://www.researchgate.net/publication/242018469_Patients_with_recurrent_biliary...)

Patients with recurrent biliary tract cancer have a better prognosis than those with unresectable disease: retrospective analysis of a multi-institutional experience with patients of advanced ...

**Name of Journal:** *World Journal of Gastrointestinal Oncology*

**Manuscript NO:** 48207

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Study*

**Multi-institutional retrospective analysis of FOLFIRI in patients with advanced biliary tract cancers**

Jonathan D Mizrahi, Valerie Gunchick, Kabir Mody, Lianchun Xiao, Phanikeerthi Surapaneni, Rachna T Shroff, Vaibhav Sahai

**Abstract**

**BACKGROUND**

Gemcitabine plus platinum (GP) is the standard of care first-line treatment for advanced biliary tract cancers (BTC). There is no established second-line therapy and retrospective

### Match Overview

1	<b>Internet</b> 49 words crawled on 10-Jul-2019 <a href="http://cslide.ctimeetingtech.com">cslide.ctimeetingtech.com</a>	2%
2	<b>Crossref</b> 21 words &NA; . "15th World Conference on Lung Cancer .", <i>Journal of Thoracic Oncology</i> , 2013.	1%
3	<b>Internet</b> 20 words crawled on 28-Feb-2019 <a href="http://mafiadoc.com">mafiadoc.com</a>	1%
4	<b>Internet</b> 17 words crawled on 24-Dec-2010 <a href="http://ash.confex.com">ash.confex.com</a>	1%
5	<b>Internet</b> 14 words crawled on 10-Oct-2017 <a href="http://www.impactjournals.com">www.impactjournals.com</a>	1%
6	<b>Crossref</b> 13 words "Abstracts", <i>Journal of Thoracic Oncology</i> , 2009	1%
7	<b>Crossref</b> 12 words A. Boere, Ingrid, and Maria E.L. van der Burg. "Review of Do se-intense Platinum and/or Paclitaxel Containing Chemo ...	1%
8	<b>Internet</b> 12 words crawled on 24-Jul-2018 <a href="http://www.lib.kobe-u.ac.jp">www.lib.kobe-u.ac.jp</a>	1%
9	<b>Internet</b> 11 words crawled on 08-May-2019 <a href="http://ir.nucana.com">ir.nucana.com</a>	<1%
10	<b>Internet</b> 10 words crawled on 14-Aug-2018 <a href="http://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a>	<1%



[国内版](#)[国际版](#)

A Multi-Institutional Retrospective Analysis of FOLFIRI in Patients with Advan

[All](#)[Images](#)[Videos](#)[翻译成中文](#)[关闭取词](#)

27,800 Results

Any time ▾

## Second-line chemotherapy in advanced biliary cancer ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4689003>

Dec 23, 2015 · Background. **Biliary tract cancers** (BTC) represent an uncommon group of **malignancies** that includes **intra-** and extra-hepatic cholangiocarcinoma, tumors of the gallbladder and tumors of the ampulla of Vater [ 1 ]. The only potentially curative approach to BTC is represented by radical resection in early stage; however,...

Cited by: 13

Author: Lorenzo Fomaro, Caterina Vivaldi, Stefan...

Publish Year: 2015

## Systemic Therapy for Biliary Tract Cancers - The Oncologist

[theoncologist.alphamedpress.org/content/13/4/415.full](http://theoncologist.alphamedpress.org/content/13/4/415.full)

Apr 01, 2008 · Introduction. **Biliary tract cancers** (BTCs) are invasive **carcinomas** that arise from the epithelial lining of the **gallbladder** and **bile** ducts. The term BTC includes both cholangiocarcinoma, which has been used to refer to **cancers** arising in the intrahepatic, perihilar, or distal **biliary tree**,...

Author: Aram F. Hezel

Published in: Oncologist · 2008

Authors: Aram F Hezel · Andrew X Zhu

Affiliation: Partners Healthcare

About: Fluorouracil · **Biliary tract** neoplasm · Deoxycytidine

## Patients with recurrent biliary tract cancer have a better ...

[https://www.researchgate.net/publication/242018469\\_Patients\\_with\\_recurrent\\_biliary...](https://www.researchgate.net/publication/242018469_Patients_with_recurrent_biliary...)

**Patients with** recurrent **biliary tract** cancer have a better prognosis than those with unresectable disease: **retrospective analysis** of a **multi-institutional** experience with **patients of advanced** ...

## [PDF] Efficacy of a sequential treatment strategy with GEMOX ...

<https://www.tandfonline.com/doi/pdf/10.1080/0284186X.2016.1191670>

reported that FOLFIRI was the most frequently used second-line treatment (32.7%) after failure of GEM-platinum first-line chemotherapy [14]. In the present study, we conducted a **retrospective analysis** of the effectiveness of sequential GEMOX-FOLFIRI-based chemotherapy in a cohort of 52 **patients** with recurrent or **advanced** BTC. **Patients** and methods

## Initial Report of Second-Line FOLFIRI in Combination with ...

[theoncologist.alphamedpress.org/lookup/resid/24/4/475?view=full&urtype=cgi](http://theoncologist.alphamedpress.org/lookup/resid/24/4/475?view=full&urtype=cgi)

Apr 01, 2019 · Initial Report of Second-Line FOLFIRI in Combination with Ramucirumab in **Advanced** Gastroesophageal Adenocarcinomas: **A Multi-Institutional Retrospective Analysis** Predictors of