

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 1,350,000 条结果 (用时 0.46 秒)

## Google 学术: Abdominal Metastases of Soft Tissue Sarcoma – A Systematic Review.

... as sites of metastases from sarcomas of soft tissue - Mazon - 被引用次数: 217

Soft tissue sarcomas - Cormier - 被引用次数: 447

Metastases to soft tissue: A review of 118 cases over a ... - Plaza - 被引用次数: 159

## Treatment of Adult Soft Tissue Sarcomas: An Overview - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4837937/> ▾ 翻译此页

作者: S Arifi - 2015 - 被引用次数: 8 - 相关文章

2015年9月7日 - The Union for International Cancer Control (UICC) recommends to .... Meta- analysis of adjuvant chemotherapy in adult soft tissue sarcomas ... with high-risk STS including abdominal or retroperitoneal location [5, 42, 43].

Primary Locally Advanced ... · Treatment of Metastatic Soft ... · Targeted Therapies

## Adult Soft Tissue Sarcoma Treatment (PDQ®) - PDQ Cancer ...

<https://www.ncbi.nlm.nih.gov/books/NBK66046/> ▾ 翻译此页

作者: PDQATE Board - 2018 - 被引用次数: 1 - 相关文章

2019年1月29日 - In a British study of 260 patients with soft tissue sarcoma diagnosed .... Alvegård TA, Berg NO: Histopathology peer review of high-grade soft tissue sarcoma: ... Soft tissue sarcoma of the abdomen and thoracic visceral organs.

## Treatment of advanced, metastatic soft tissue sarcoma: latest evidence ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5524246/> - 翻译此页

作者: GK In - 2017 - 被引用次数: 20 - 相关文章

2017年6月15日 - Historically, there have been few systemic treatment options for this ..... Metastatic soft tissue sarcoma, an analysis of systemic therapy and ...

## Soft-tissue Sarcoma Metastases Identified on Abdomen and Pelvis CT ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2758993/> ▾ 翻译此页



国内版 国际版

Chat with Bing

Abdominal metastases of primary extremity soft tissue sarcom



Sign in



Add Bing Firefox extension

ALL

IMAGES

VIDEOS

开启取词

50,600 Results

Any time

### [PDF] Functional Outcome Measurement in Patients with Lower ...

<https://link.springer.com/content/pdf/10.1245/s10434-019-07698-w.pdf>

in the treatment of patients with **extremity soft tissue sarcoma** (STS) has been increasingly recognized in the last three decades. This **systematic review** aimed to investigate how FO is measured in surgically treated lower-**extremity** STS patients. Methods. A **systematic** search of PubMed, Web of Sci-ence, and Scopus was performed based on the PRISMA

Author: Gilber Kask, Ian Barner-Rasmussen, ... Publish Year: 2019

### Primary retroperitoneal soft tissue sarcoma: Imaging ...

<https://www.sciencedirect.com/science/article/pii/S0748798316310101>

Because **soft tissue sarcoma** accounts for only a third of retroperitoneal **tumours** arising extrinsic to the solid **abdominal viscera**, other diagnoses must be considered. 26 Percutaneous core needle biopsy usually confirms the diagnosis and is the gold standard for diagnosis, but rarely lesions are not amenable or high risk for biopsy and the differential diagnosis based on imaging becomes crucial. **MRI** is ...

Cited by: 22 Author: C. Messiou, E Moskovic, D. Vanel, C. Mo...

Publish Year: 2017

### Metastases to soft tissue - Plaza - 2008 - Cancer - Wiley ...

<https://onlinelibrary.wiley.com/doi/full/10.1002/cncr.23151>

Nov 26, 2007 - **Metastases** to **soft tissue** are usually noted in patients with advanced disease, but occasionally they can present in patients with no known **primary** tumor and masquerade as a **soft tissue sarcoma**. 4 Multiple case reports and small series of patients have been reported in the literature addressing this rare phenomenon. 2, 3, 8, 9, 12 Autopsy ...

Cited by: 183 Author: Jose Antonio Plaza, Delia Perez-Montiel, ...

Publish Year: 2008

### Abdominal soft tissue sarcoma: A multicenter retrospective ...

[https://www.researchgate.net/publication/43081014\\_Abdominal\\_soft\\_tissue\\_sarcoma\\_A...](https://www.researchgate.net/publication/43081014_Abdominal_soft_tissue_sarcoma_A...)

Primary mesenchymal tumors of the colon are extremely rare tumors among **soft tissue sarcomas**. These **tumors** are more aggressive and have poorer prognosis than **adenocarcinoma** of the colon.

### Patterns of metastatic spread following resection of ...

<https://onlinelibrary.wiley.com/doi/abs/10.1002/ssu.2980040106>

Eighty-five patients developed **metastatic disease**: 13 had isolated local recurrence, 43 had isolated

resection, especially regarding the surveillance of potential AM/RM.

## *AIM*

To systematically analyse incidence, diagnosis, treatment and outcome of AM/RM in eSTS patients.

## *METHODS*

In this systematic review, 899 studies available in PubMed and published between 2000 and 2018 were screened, identifying 17 original articles focused on AM or RM in eSTS. Article selection was based on the PRISMA guidelines, using the search terms [abdominal metastasis AND soft tissue sarcoma] and [soft tissue sarcoma metastasis abdomen]. All studies published between January, 2000 and December, 2018 were screened. Further articles were identified by cross-searching article references, with final search date being February 18, 2019. Due to limited data and different reporting techniques used, the present review focused on descriptive analysis of the studies included

### Match Overview

1 Internet 13 words  
crawled on 28-Aug-2019  
[f6publishing.blob.core.windows.net](http://f6publishing.blob.core.windows.net)

&lt;1%

## Abdominal metastases of primary extremity soft tissue sarcom



ALL

IMAGES

VIDEOS

关闭取词

85,400 Results

Any time ▾

## Synovial sarcoma of the abdominal wall: Imaging findings ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4391800>

Feb 28, 2015 · The **anterior abdominal wall** is an uncommon site of **synovial sarcoma**, which is the fourth most common type of **soft-tissue sarcoma**. MR imaging remains the optimal modality for diagnosis and staging, best local control is achieved by radical surgical excision, and because of its tendency for late local **recurrence** and **metastases**, a follow-up period of at least 10 years seems ...

Cited by: 3

Author: Robbert J. de Haas, Johannes J. Bonenk...

Publish Year: 2015

## Soft-tissue Sarcoma Metastases Identified on Abdomen and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2758993>

Jul 28, 2009 · We included 124 patients with 15 types of **extremity soft tissue** sarcomas evaluated with CT of the C/A/P. **Primary** outcomes were (1) location of **metastatic** disease in relation to (2) sarcoma type. Twenty patients (16%) presented with or developed abdomen/pelvis **metastases** and 10 of the 15 types of **soft tissue** sarcomas had **abdominal** or pelvic ...

Cited by: 10

Author: David M. King, Donald A. Hackbarth, Chri...

Publish Year: 2009

## Functional Outcome Measurement in Patients with Lower ...

<https://link.springer.com/article/10.1245/s10434-019-07698-w> ▾

Aug 12, 2019 · Flow diagram showing flow of studies retrieved for **systematic review** of functional outcome measurement in patients with lower-**extremity soft tissue** sarcoma Study Characteristics The included studies were published between 1984 and 2018.

Author: Gilber Kask, Ian Barner-Rasmussen, J... Publish Year: 2019

## Primary retroperitoneal soft tissue sarcoma: Imaging ...

<https://www.sciencedirect.com/science/article/pii/S0748798316310101>

Because **soft tissue sarcoma** accounts for only a third of retroperitoneal **tumours** arising extrinsic to the solid **abdominal viscera**, other diagnoses must be considered. 26 Percutaneous core needle biopsy usually confirms the diagnosis and is the gold standard for diagnosis, but rarely lesions are not amenable