

검색결과 약 28,700개 (0.56초)

Two-stage revision arthroplasty for coagulase-negative staphylococcal periprosthetic joint infection of the hip and knee에 대한 학술자료

Prosthetic joint infections: bane of orthopedists, ... - Lentino - 470회 인용

... **infection** by frozen section during **revision arthroplasty** - Athanasou - 213회 인용

Prosthetic-joint infections - Zimmerli - 2370회 인용

🔗 <https://www.ncbi.nlm.nih.gov/pmc/PMC5226970> ▼ 이 페이지 번역하기

Two-stage Revision for Periprosthetic Hip and Knee Joint Infections

SG Kini 저술 - 2016 - 15회 인용 - 관련 학술자료

2016. 11. 30. - Keywords: **Infection, Hip, Knee, Prosthetic joint, Two-stage revision** of infection , particularly with **coagulase negative staphylococcal infections**. ... The **second stage** of revision **hip and knee arthroplasty** involves spacer ...

[Abstract](#) · [INTRODUCTION](#) · [FIRST STAGE](#) · [SECOND STAGE](#)

🔗 <https://www.ncbi.nlm.nih.gov/pmc/PMC6123506> ▼ 이 페이지 번역하기

Management of Periprosthetic Joint Infection - NCBI

C Li 저술 - 2018 - 9회 인용 - 관련 학술자료

2018. 9. 4. - The **infection** rate after **hip** or shoulder replacement is usually less than 1%, after **knee** replacement less than 2% and after elbow replacement is between 1.9% and 10.3%. ... **Coagulase-negative staphylococci** and **Staphylococcus aureus** are the most common microorganisms in **hip** and **knee PJI**⁸⁾.



国内版

国际版

Two-Stage Revision Arthroplasty for Coagulase-Negative Staphylococcal Periprosth



All

Images

Videos

翻译成中文

关闭取词

30,800 Results

Any time ▾

Two-stage Revision for Periprosthetic Hip and Knee Joint ...

<https://openorthopaedicsjournal.com/VOLUME/10/PAGE/579/FULLTEXT> ▾

Two-stage Revision for Periprosthetic Hip and Knee Joint Infections Sunil Gurpur Kini *, 1 , Ayman Gabr 1 , Rishi Das 1 , Mohamed Sukeik 2 , Fares Sami Haddad 1 1 Department of Trauma and Orthopaedics, University College London Hospital, 235 Euston ...

Cited by: 8

Author: Sunil Gurpur Kini, Ayman Gabr, Rishi Da...

Publish Year: 2016

Two-stage Revision for Periprosthetic Hip and Knee Joint ...

https://www.researchgate.net/publication/311509907_Two-stage_Revision_for...

Two-stage revision arthroplasty of the hip for infection using an interim articulated Prostalac hip spacer: a 10 to 15-year follow-up study. J Bone Joint Surg Br 2009; 91(11): 1431-7.

Two Stage Hip Revision in Periprosthetic Infection ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2923340>

Jun 11, 2010 - DISCUSSION. Nowadays, two-stage revision arthroplasty is considered the gold standard treatment method for deep periprosthetic infection [22], however it involves a high cost for the medical healthcare system, and much effort for patients, paramedical staffs and surgeons. It needs at least two operations to cure.

Cited by: 19

Author: Giovanni Pignatti, Shingo Nitta, Nicola Ra...

Publish Year: 2010

Prosthetic Joint Infection - Recon - Orthobullets

<https://www.orthobullets.com/recon/5004/prosthetic-joint-infection> ▾

Coagulase-negative Staphylococcus (chronic infections) most common fungal pathogen Pain and Blood Loss Management after Total Hip and Knee Arthroplasty - The Secret Sauce for Rapid Recovery Total Joints - Charles DeCook, MD, Arthritis & Total Joint Specialist, Atlanta (COA 2018, 1.1) ... - Prosthetic Joint Infection 2/4/2017 226 . 6 . 30 ...

Methicillin-resistant Staphylococcal periprosthetic joint ...

<https://bmcinfectdis.biomedcentral.com/articles/10.1186/s12879-016-1366-9> ▾

Feb 01, 2016 · Periprosthetic joint infections (PJI) of the hip and knee are disastrous complications that occur in approximately 1 to 2 % of patients after total joint arthroplasties [].The management of PJI may require long-term antibiotic suppression, surgical debridement, one-stage or 2-stage revision, resection arthroplasty, arthrodesis, or amputation.

Cited by: 6

Author: Feng-Chih Kuo, Feng-Chih Kuo, Shih-Hsi...

Publish Year: 2015

Two-stage revision arthroplasty for coagulase-negative staphylococcal periprosthe



All

Images

Videos

翻译成中文

关闭取词

35,800 Results

Any time ▾

Two-stage Revision for Periprosthetic Hip and Knee Joint ...

https://www.researchgate.net/publication/311509907_Two-stage_Revision_for...

Keywords: Infection, Hip, Knee, Prosthetic joint, Two-stage revision. INTRODUCTION Prosthetic joint infection (PJI) remains one of the most dreadful complications following total joint arthroplasty.

Two-stage Revision for Periprosthetic Hip and Knee Joint ...

<https://openorthopaedicsjournal.com/VOLUME/10/PAGE/579/FULLTEXT> ▾

Two-stage revision arthroplasty of the **hip** for infection using an interim **articulated Prostalac hip spacer**: a 10 to 15-year follow-up study. **J Bone Joint Surg Br** 2009; 91(11): 1431-7. CrossRef PubMed

Cited by: 11

Author: Sunil Gurbur Kini, Ayman Gabr, Rishi Das...

Publish Year: 2016

Two Stage Hip Revision in Periprosthetic Infection ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2923340>

Jun 11, 2010 · Nowadays, **two-stage revision arthroplasty** is considered the **gold standard treatment method** for **deep periprosthetic infection**, however it involves a high cost for the **medical healthcare system**, and much effort for patients, paramedical staffs and surgeons. It needs at least **two operations** to cure.

Cited by: 19

Author: Giovanni Pignatti, Shingo Nitta, Nicola Ra...

Publish Year: 2010

Two-Stage Exchange Arthroplasty for Periprosthetic Joint ...

<https://www.sciencedirect.com/science/article/pii/S0883540319305911>

Jun 15, 2019 : Two-Stage Exchange **Arthroplasty** for **Periprosthetic Joint Infection**: The Rate and Reason for the Attrition After the First Stage Author links open overlay panel Qiaojie Wang MD a b Karan Goswami MD, MRCS a Feng-Chih Kuo MD a c Chi Xu MD a ...

Author: Qiaojie Wang, Qiaojie Wang, Karan G...

Publish Year: 2019

Revision arthroplasty for periprosthetic joint infections ...

<https://www.sciencedirect.com/science/article/pii/S1045452719300040>

Name of Journal: *World Journal of Orthopedics*

Manuscript NO: 49236

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Two-stage revision arthroplasty for coagulase-negative staphylococcal
periprosthetic joint infection of the hip and knee

Veltman ES *et al.* Mid-term follow-up of two-stage revision arthroplasties

Ewout S Veltman, ⁶Dirk Jan F Moojen, Marc L van Ogtrop, Rudolf W Poolman

Abstract

BACKGROUND

Match Overview

| | | |
|---|--|-----|
| 1 | Internet 22 words crawled on 25-Aug-2019 mrnjournal.biomedcentral.com | 1% |
| 2 | Crossref 19 words Joris B.W. Elbers, Borg Leijten, Hannah E. van Werven, Patrick D. Sturm, Bart Jan Kullberg, Berend W. Schreurs. "Antib | 1% |
| 3 | Crossref 14 words X. Yuan, C. Chen, J. Zhou, M. Han, X. Wang, C. Wang, X. H e. "Organ Donation and Transplantation From Donors With ... | <1% |
| 4 | Crossref 11 words Christophe Pattyn, Thomas De Geest, Pieter Ackerman, Emmanuel Audenaert. "Preformed gentamicin spacers in two ... | <1% |
| 5 | Crossref 11 words C.L. Romanò, G. Manzi, S. Scarponi, N. Logoluso, D. Geor ... e. "Practice and guidelines for treating periprosthetic joint in | <1% |
| 6 | Internet 11 words crawled on 04-Jun-2018 home.biolsci.org | <1% |
| 7 | Crossref 10 words Koo, K.H.. "Impregnation of vancomycin, gentamicin, and ceftaxime in a cement spacer for two-stage cementless re ... | <1% |
| 8 | Crossref 10 words "Long-Term Health-Related Quality of Life After Surgery for Adolescent Idiopathic Scoliosis and Spondylolisthesis", The | <1% |
| 9 | Crossref 10 words Leif Claassen, Christian Plaass, Kiriakos Daniilidis, Tilman C allies, Gabriela von Lewinski. "Two-Stage Revision Total K | <1% |