

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 49308

Manuscript Type: CASE REPORT

Severe mental disorders following anti-retroviral treatment in a patient on peritoneal dialysis: A case report and literature review

He QE *et al.* Mental disorders after anti-retroviral

Qi-En He, Min Xia, Guang-Hui Ying, Xue-Lin He, Jiang-Hua Chen, Yi Yang

Abstract

BACKGROUND

Antiviral drugs are widely used in populations with viral infection caused by

Match Overview

There are no matching sources for this report.

[国内版](#)[国际版](#)

Severe mental disorders following anti-retroviral treatment in a patient on pe

[All](#)[Images](#)[Videos](#)[翻译成中文](#)[关闭取词](#)

9,880 Results

Any time ▾

No Influence of Dialysis on Mirtazapine – A Case Report ...

https://www.researchgate.net/publication/23642609_No_Influence_of_Dialysis_on...

Given that **pain** represents the number one reason that **patients** visit general practitioners, with 45% of visits related to **pain**, and that **depression** is the most common **mental health disorder**, with ...

Quality of life, mental health and ... - PubMed Central (PMC)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2611965>

The aim of the present study was to investigate a) self-reported physical, psychological, social and environmental well-being, b) mental health factors such as **anxiety and depression**, as well as c) illness beliefs of **health locus of control** in haemodialysis and **peritoneal dialysis patients**.

Cited by: 195

Author: M Ginieri-Coccossis, P Theofilou, C Syno...

Publish Year: 2008

Patient Ratings of Dialysis Care With Peritoneal Dialysis ...

<https://jamanetwork.com/journals/jama/fullarticle/198181> ▾

Feb 11, 2004 · Conclusions. Nearly 100 000 **patients with incident end-stage renal disease (ESRD)** in the United States must choose between hemodialysis and peritoneal dialysis each year in light of a limited number of donor kidneys for transplantation. Since the 1970s, **patients** have had the choice of these 2 modalities.

Cited by: 291

Author: Haya R. Rubin, Nancy E. Fink, Laura C. P...

Publish Year: 2004

Author: Haya R. Rubin, MD, PhD

Calciophylaxis in end-stage renal disease prior to dialytic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4334332>

Calciophylaxis in end-stage renal disease prior to dialytic **treatment**: a **case report and literature review**
Ramón Alberto Tamayo-Isla and Mauro Cuba de la Cruz Internal Medicine Department, Pietersburg
Provincial Hospital, Polokwane Kidney and **Dialysis** Centre, Limpopo Province, South Africa

Cited by: 5

Author: Ramón Alberto Tamayo-Isla, Mauro Cuba ...

Publish Year: 2015

Ertapenem-Induced Delirium: A Case Report and Literature ...

https://www.researchgate.net/publication/273400898_Ertapenem-Induced_Delirium_A_Case...

A **literature** search of similar **case** reports has shown that seizures during ertapenem **treatment** may develop in **patients** with prior known CNS **disorders** and/or severe renal dysfunction that predicts ...

Hyperammonemic Encephalopathy Associated With ...

theoncologist.alphamedpress.org/content/21/4/514.full

Apr 01, 2016 · Hyperammonemic Encephalopathy Associated With Fibrolamellar Hepatocellular Carcinoma: **Case Report**, **Literature Review**, and Proposed **Treatment Algorithm** ... On day 3 after



All

Images

Videos

翻译成中文

关闭取词

11,500 Results

Any time ▼

Quality of life, mental health and health beliefs in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2611965>

Nov 14, 2008 · The aim of the present study was to investigate a) self-reported physical, psychological, social and environmental well-being, b) **mental health** factors such as **anxiety and depression**, as well as c) **illness** beliefs of **health** locus of control in haemodialysis and **peritoneal dialysis** patients.

Cited by: 200

Author: M Ginieri-Coccossis, P Theofilou, C Syno...

Publish Year: 2008

Skin disorders in peritoneal dialysis patients: An ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4936344>

Jul 06, 2016 · But, data about **peritoneal dialysis** patients regarding skin changes is lacking in the **literature** except for a few studies subjecting only skin color changes and xerosis. The authors aimed to examine all skin changes in **peritoneal dialysis** (PD) **patients** followed up in the authors' PD unit.

Cited by: 1

Author: Meltem Gursu, Sami Uzun, Derya Topcuo...

Publish Year: 2016

Posterior reversible encephalopathy syndrome in an adult ...

<https://bmcnephrol.biomedcentral.com/articles/10.1186/1471-2369-15-10> ▼

Jan 13, 2014 · Clinically, PRES is characterized by visual disturbances, **headache**, nausea, change in **mental status**, and seizure. **Seizures** are cited as the commonest manifestation of PRES, occurring in up to 90% of reported cases [3, 4]. **Seizure type** is mainly generalized, but **focal onset** ...

Cited by: 20

Author: Brett R Graham, George B Pylypchuk

Publish Year: 2014

Author: Brett R Graham, George B Pylypchuk

[PDF] Calciphylaxis in end-stage renal disease prior to dialytic ...

https://www.researchgate.net/publication/272843068_Calciphylaxis_in_end-stage_renal...

Introduction: Calciphylaxis is a rare medical condition that is usually **diagnosed** in **patients** suffering from **end-stage renal disease** who are already receiving **renal replacement therapy** and in ...

[PDF] The influence of different dialysis modalities on the ...

<https://www.tandfonline.com/doi/pdf/10.1080/08870446.2016.1226307>

patients treated with different **dialysis** modalities. Methods: A **literature** search was conducted for English language articles in the CINAHL, Medline and PubMed databases published from January 1990 through May 2016. Specifically, we sought articles that would compare the HR-QOL of hemodialysis (HD)



国内版

国际版

Severe mental disorders following anti-retroviral treatment in



All

Images

Videos

翻译成中文

关闭取词

42,200 Results

Any time ▾

Dialysis (Hemodialysis): Purpose, Procedure, and Complications

<https://www.webmd.com/a-to-z-guides/kidney-dialysis> ▾

It's a **treatment** that takes over your kidney functions if those organs stop doing their job. There are two types of **dialysis**: Hemodialysis: Your blood is put through a filter outside your body ...

Author: Kelli Miller

Quality of life, mental health and health beliefs in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2611965>

Nov 14, 2008 · The aim of the present study was to investigate a) self-reported physical, psychological, social and environmental well-being, b) **mental** health factors such as anxiety and depression, as well as c) **illness** beliefs of health locus of control in haemodialysis and **peritoneal dialysis patients**.

Cited by: 200

Author: M Ginieri-Coccossis, P Theofilou, C Synodi...

Publish Year: 2008

Quality of life, mental health and health beliefs in ...

<https://bmcnephrol.biomedcentral.com/articles/10.1186/1471-2369-9-14> ▾

Nov 14, 2008 · The study examines differences regarding quality of life (QoL), **mental** health and **illness** beliefs between in-centre haemodialysis (HD) and continuous ambulatory **peritoneal dialysis** (CAPD/PD) **patients**. Differences are examined between **patients** who recently commenced **treatment** compared to **patients** on long term **treatment**. 144 End-Stage Renal Disease (ESRD) **patients** were recruited from ...

Cited by: 200

Author: M Ginieri-Coccossis, P Theofilou, C Synodi...

Publish Year: 2008

Author: M Ginieri-Coccossis

Skin disorders in peritoneal dialysis patients: An ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4936344>

Jul 06, 2016 · But, data about **peritoneal dialysis patients** regarding skin changes is lacking in the **literature** except for a few studies subjecting only skin color changes and xerosis. The authors aimed to examine all skin changes in **peritoneal dialysis** (PD) **patients** followed up in the authors' PD unit.

Cited by: 1

Author: Meltem Gursu, Sami Uzun, Derya Topcuoğlu...

Publish Year: 2016

Severe mental illness and chronic kidney disease: a cross ...

<https://www.dovepress.com/severe-mental-illness-and-chronic-kidney-disease-a-cross...> ▾

The higher prevalence of CKD in **patients** with SMI may, in part, be explained by more frequent blood testing as compared to the general population. Conclusion: CKD is identified more commonly among **patients** with SMI than in the general population. Keywords: **severe mental illness**, schizophrenia, bipolar