

World Journal of *Clinical Cases*

World J Clin Cases 2019 December 6; 7(23): 3915-4171



**REVIEW**

- 3915** Overview of organic anion transporters and organic anion transporter polypeptides and their roles in the liver
Li TT, An JX, Xu JY, Tuo BG

ORIGINAL ARTICLE**Observational Study**

- 3934** Value of early diagnosis of sepsis complicated with acute kidney injury by renal contrast-enhanced ultrasound
Wang XY, Pang YP, Jiang T, Wang S, Li JT, Shi BM, Yu C
- 3945** Value of elastography point quantification in improving the diagnostic accuracy of early diabetic kidney disease
Liu QY, Duan Q, Fu XH, Fu LQ, Xia HW, Wan YL
- 3957** Resection of recurrent third branchial cleft fistulas assisted by flexible pharyngotomy
Ding XQ, Zhu X, Li L, Feng X, Huang ZC
- 3964** Therapeutic efficacy of acupuncture combined with neuromuscular joint facilitation in treatment of hemiplegic shoulder pain
Wei YH, Du DC, Jiang K
- 3971** Comparison of intra-articular injection of parecoxib *vs* oral administration of celecoxib for the clinical efficacy in the treatment of early knee osteoarthritis
Lu L, Xie Y, Gan K, Huang XW

Retrospective Study

- 3980** Celiomesenteric trunk: New classification based on multidetector computed tomography angiographic findings and probable embryological mechanisms
Tang W, Shi J, Kuang LQ, Tang SY, Wang Y

Prospective Study

- 3990** Interaction of arylsulfatases A and B with maspin: A possible explanation for dysregulation of tumor cell metabolism and invasive potential of colorectal cancer
Kovacs Z, Jung I, Szalman K, Baniás L, Bara TJ, Gurzu S

CASE REPORT

- 4004** Recuperation of severe tumoral calcinosis in a dialysis patient: A case report
Westermann L, Isbell LK, Breitenfeldt MK, Arnold F, Röthele E, Schneider J, Widmeier E

- 4011** Robotic wedge resection of a rare gastric perivascular epithelioid cell tumor: A case report
Marano A, Maione F, Woo Y, Pellegrino L, Geretto P, Sasia D, Fortunato M, Orcioni GF, Priotto R, Fasoli R, Borghi F
- 4020** Primary parahiatal hernias: A case report and review of the literature
Preda SD, Pătrașcu Ș, Ungureanu BS, Cristian D, Bințișan V, Nica CM, Calu V, Strâmbu V, Sapalidis K, Șurlin VM
- 4029** Diagnosis of Laron syndrome using monoplex-polymerase chain reaction technology with a whole-genome amplification template: A case report
Neumann A, Alcántara-Ortigoza MÁ, González-del Ángel A, Camargo-Diaz F, López-Bayghen E
- 4036** *In-vitro* proliferation assay with recycled ascitic cancer cells in malignant pleural mesothelioma: A case report
Anayama T, Taguchi M, Tatenuma T, Okada H, Miyazaki R, Hirohashi K, Kume M, Matsusaki K, Orihashi K
- 4044** Distant metastasis in choroidal melanoma with spontaneous corneal perforation and intratumoral calcification: A case report
Wang TW, Liu HW, Bee YS
- 4052** Secondary Parkinson disease caused by breast cancer during pregnancy: A case report
Li L
- 4057** Pulmonary embolism and deep vein thrombosis caused by nitrous oxide abuse: A case report
Sun W, Liao JP, Hu Y, Zhang W, Ma J, Wang GF
- 4063** Micronodular thymic tumor with lymphoid stroma: A case report and review of the literature
Wang B, Li K, Song QK, Wang XH, Yang L, Zhang HL, Zhong DR
- 4075** Diffuse large B cell lymphoma with bilateral adrenal and hypothalamic involvement: A case report and literature review
An P, Chen K, Yang GQ, Dou JT, Chen YL, Jin XY, Wang XL, Mu YM, Wang QS
- 4084** Urethral pressure profilometry in artificial urinary sphincter implantation: A case report
Meng LF, Liu XD, Wang M, Zhang W, Zhang YG
- 4091** Hydroxyurea-induced cutaneous squamous cell carcinoma: A case report
Xu Y, Liu J
- 4098** Recurrent hypotension induced by sacubitril/valsartan in cardiomyopathy secondary to Duchenne muscular dystrophy: A case report
Li JM, Chen H
- 4106** Complete duodenal obstruction induced by groove pancreatitis: A case report
Wang YL, Tong CH, Yu JH, Chen ZL, Fu H, Yang JH, Zhu X, Lu BC

- 4111** Radiological aspects of giant hepatocellular adenoma of the left liver: A case report
Zheng LP, Hu CD, Wang J, Chen XJ, Shen YY
- 4119** Mixed serous-neuroendocrine neoplasm of the pancreas: A case report and review of the literature
Xu YM, Li ZW, Wu HY, Fan XS, Sun Q
- 4130** Rigid esophagoscopy combined with angle endoscopy for treatment of superior mediastinal foreign bodies penetrating into the esophagus caused by neck trauma: A case report
Wang D, Gao CB
- 4137** Left armpit subcutaneous metastasis of gastric cancer: A case report
He FJ, Zhang P, Wang MJ, Chen Y, Zhuang W
- 4144** Bouveret syndrome: A case report
Wang F, Du ZQ, Chen YL, Chen TM, Wang Y, Zhou XR
- 4150** Fatal complications in a patient with severe multi-space infections in the oral and maxillofacial head and neck regions: A case report
Dai TG, Ran HB, Qiu YX, Xu B, Cheng JQ, Liu YK
- 4157** Management of massive fistula bleeding after endoscopic ultrasound-guided pancreatic pseudocyst drainage using hemostatic forceps: A case report
Ge N, Sun SY
- 4163** Pure squamous cell carcinoma of the gallbladder locally invading the liver and abdominal cavity: A case report and review of the literature
Jin S, Zhang L, Wei YF, Zhang HJ, Wang CY, Zou H, Hu JM, Jiang JF, Pang LJ

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Consolato M Sergi, FRCP (C), MD, PhD, Professor, Department of Lab Medicine and Pathology, University of Alberta, Edmonton T6G 2B7, Canada

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in PubMed, PubMed Central, Science Citation Index Expanded (also known as SciSearch®), and Journal Citation Reports/Science Edition. The 2019 Edition of Journal Citation Reports cites the 2018 impact factor for WJCC as 1.153 (5-year impact factor: N/A), ranking WJCC as 99 among 160 journals in Medicine, General and Internal (quartile in category Q3).

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Yan-Xia Xing*

Proofing Production Department Director: *Xiang Li*

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Semimonthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Bao-Gan Peng, Sandro Vento

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2307-8960/editorialboard.htm>

EDITORIAL OFFICE

Jin-Lei Wang, Director

PUBLICATION DATE

December 6, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Pure squamous cell carcinoma of the gallbladder locally invading the liver and abdominal cavity: A case report and review of the literature

Shan Jin, Lu Zhang, Yuan-Feng Wei, Hai-Jun Zhang, Cheng-Yan Wang, Hong Zou, Jian-Ming Hu, Jin-Fang Jiang, Li-Juan Pang

ORCID number: Shan Jin (0000-0001-9906-8256); Lu Zhang (0000-0002-6852-4236); Yuan-Feng Wei (0000-0001-6338-7546); Hai-Jun Zhang (0000-0002-3577-3324); Cheng-Yan Wang (0000-0002-7905-9157); Hong Zou (0000-0003-3051-8006); Jian-Ming Hu (0000-0001-7790-7979); Jin-Fang Jiang (0000-0001-8631-5617); Li-Juan Pang (0000-0003-0017-7634).

Author contributions: Jin S and Zhang L were the patient's pathologists, reviewed the literature, and contributed to manuscript drafting; Wei YF reviewed the literature and contributed to manuscript drafting; Zhang HJ and Wang CY performed the etiological analyses and interpretation and contributed to manuscript drafting; Zou H analyzed and interpreted the clinicopathological features; Hu JM and Jiang JF performed the molecular mechanism consultation, reviewed the literature, and drafted the manuscript; Pang LJ was responsible for the revision of the manuscript for important intellectual content; all authors issued final approval for the version to be submitted.

Supported by the National Natural Science Foundation of China, No. 81560053; the Youth Science and Technology Innovation Leading Talents Project of Corps, No. 2017CB004; International Science and Technology Cooperation Promotion Plan of Shihezi University, No. GJHZ201805;

Shan Jin, Lu Zhang, Yuan-Feng Wei, Hai-Jun Zhang, Cheng-Yan Wang, Hong Zou, Jian-Ming Hu, Jin-Fang Jiang, Li-Juan Pang, Department of Pathology, the First Affiliated Hospital to Shihezi University School of Medicine and Shihezi University School of Medicine, Shihezi 832002, Xinjiang Uygur Autonomous Region, China

Shan Jin, Lu Zhang, Yuan-Feng Wei, Hai-Jun Zhang, Cheng-Yan Wang, Hong Zou, Jian-Ming Hu, Jin-Fang Jiang, Li-Juan Pang, Key Laboratory of Xinjiang Endemic and Ethnic Diseases (Ministry of Education), Shihezi University School of Medicine, Shihezi 832002, Xinjiang Uygur Autonomous Region, China

Corresponding author: Li-Juan Pang, MD, PhD, Chief Physician, Professor, Department of Pathology, Shihezi University School of Medicine and the First Affiliated Hospital to Shihezi University School of Medicine, North 2nd Road, Shihezi 832002, Xinjiang Uygur Autonomous Region, China. oceen123456@163.com
Telephone: +86-993-2850955
Fax: +86-993-2015620

Abstract

BACKGROUND

Gallbladder squamous cell carcinoma (GBSCC) is a rare subtype of malignancy and accounts for only 2%-3% of gallbladder malignancies. Due to its rapid development, most patients with GBSCC initially present with an advanced stage of the disease and hence a poor prognosis. The clinicopathological and biological features of SCC remain to be fully elucidated, owing to its uncommon occurrence. The majority of currently available data only described individual case reports or series analyses of trivial cases.

CASE SUMMARY

A 64-year-old man was admitted for progressively poor abdominal distension and pain. Liver computed tomography (CT) showed infiltration of gallbladder carcinoma into the adjacent liver, and enlarged retroperitoneal lymph nodes. The patient underwent radical cholecystectomy. Part of the mass was grey and soft, and the neoplastic section showed a purulent-necrotic lesion. Hematoxylin and eosin staining revealed a moderately differentiated SCC. Immunohistochemical studies showed strong staining of the tumor for AE1/3 and CK5/6. Staining for CK19, CK7, and CAM5.2 was positive in the cytoplasm. Systemic chemotherapy was not administered because of the patient's poor physical condition. After five

Xinjiang Production and Construction Corps Key Areas Innovation Team Project, No. 2018CB002.

Informed consent statement:

Written informed consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

CARE Checklist (2016) statement:

The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: June 19, 2019

Peer-review started: June 26, 2019

First decision: September 9, 2019

Revised: September 30, 2019

Accepted: October 15, 2019

Article in press: October 15, 2019

Published online: December 6, 2019

P-Reviewer: González-González R

S-Editor: Dou Y

L-Editor: Wang TQ

E-Editor: Liu MY



months, CT and magnetic resonance cholangiopancreatography showed multiple metastases in the liver and abdominal cavity.

CONCLUSION

Squamous components of GBSCC may explain the complex biological behavior, and CD109 may be involved in the pathogenesis.

Key words: Gallbladder; Squamous cell carcinoma; Squamous metaplasia; Cholecystectomy; Case report

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Squamous cell carcinoma of gallbladder (GBSCC) is a rare subtype, and it has a poor prognosis. The squamous components that may explain the complex biological behavior and its poor prognosis show greater capacities of proliferation and local invasive in gallbladder carcinoma. Surgical resection has been considered as the basis of treatment for years, and complete resection is associated with increased survival. CD109 may be involved in the pathogenesis of GBSCC and serves as a novel target for intervention.

Citation: Jin S, Zhang L, Wei YF, Zhang HJ, Wang CY, Zou H, Hu JM, Jiang JF, Pang LJ. Pure squamous cell carcinoma of the gallbladder locally invading the liver and abdominal cavity: A case report and review of the literature. *World J Clin Cases* 2019; 7(23): 4163-4171
URL: <https://www.wjnet.com/2307-8960/full/v7/i23/4163.htm>
DOI: <https://dx.doi.org/10.12998/wjcc.v7.i23.4163>

INTRODUCTION

Gallbladder carcinoma (GBC) is a common type of cancer in the gastrointestinal tract, and adenocarcinoma (AC) is the most common type of gallbladder cancer^[1]. Squamous cell carcinoma (SCC) is a rare subtype and is responsible for only 2%-3% of gallbladder malignancies^[2,3]. Due to its rapid development, the majority of patients with gallbladder SCC (GBSCC) initially present with an advanced stage of the disease and hence a poor prognosis^[1]. Although patients underwent surgical interventions, the 5-year survival rate remained at approximately 1%. The large majority of SCCs are invasive carcinoma entity, often invading the entire gallbladder wall. It had been speculated that adenosquamous carcinomas (ASCs)/SCCs are more aggressive than ordinary gallbladder ACs. It has been testified that the squamous component of GBCs proliferates at a higher rate than the glandular component. However, despite their high proliferation rate, these tumors appeared to less frequently present with lymph node metastasis than gallbladder ACs. Their aggressive biological behavior has been attributed to their potential for direct extension and early invasion into the liver and neighboring organs, such as the stomach, duodenum, and transverse colon. Generally, SCC is aggressive, with infiltration into the liver, and rarely metastasizes to lymph nodes^[4]. The clinicopathological and biological features of SCC remain to be fully elucidated^[5,6], owing to its uncommon occurrence. The majority of currently available data only described individual case reports or series analyses of trivial cases. Therefore, it is imperative to describe the clinicopathological and biological features of SCC^[5,6]. To improve our understanding of SCC of the gallbladder, we conducted a retrospective case study and review of the literature^[6].

CASE PRESENTATION

Chief complaints

A 64-year-old man presented with symptoms of constantly worsening right upper quadrant distension and pain for 1 d.

History of present illness

The patient's symptoms started a month ago with symptoms of right upper quadrant distension and pain, which had worsened the last 48 h.

History of past illness

The patient did not have a previous medical history.

Personal and family history

The patient denied any personal or family history.

Imaging examinations

Ultrasonography (USG) of the abdomen showed a 5.8 cm × 4.5 cm heterogeneous mass localized in both the gallbladder and liver and a gallbladder with remarkably thickened wall, indicating a malignancy. Computed tomography (CT) and contrast-enhanced CT of the liver revealed infiltration of GBC into the adjacent liver, with enlarged retroperitoneal lymph nodes (Figure 1).

The patient underwent radical cholecystectomy for GBC. The gallbladder was intraoperatively observed to adhere tightly to the greater omentum. Moreover, miliary lesions were observed in a portion of the omentum. Neoplastic entities were present at the bottom of the gallbladder, which had invaded the V hepatic segments of the liver.

The gallbladder and a part of the liver tissue were surgically resected (dimension of 13 mm × 5 mm × 8 mm). The gallbladder contained dark green bile. The thickness of the gallbladder wall was approximately 0.2-0.3 cm, and the size of the tumor near the neck of the gallbladder was approximately 5 cm × 4 cm × 6 cm. Part of the mass was grey and soft, and the neoplastic section of the tumor revealed purulent necrosis. The boundaries of the mass and surrounding liver tissue were vague (Figure 2).

Pathological findings

Hematoxylin and eosin staining: For paraffin-embedded tissue specimens, consecutive 3-5 μm thick sections were cut and used for histological analysis^[7]. Hematoxylin and eosin staining revealed moderately differentiated SCC. The tumor cells invaded the full-thickness of the serosa and approached the surrounding adipose tissue. Part of the tumor also infiltrated the liver tissue. Lymph nodes in the hepatic duodenal ligament were examined, and resected lymph nodes were negative for carcinoma metastasis (Figure 3).

Immunohistochemical staining: Immunohistochemical analysis showed strong staining for AE1/3 and CK5/6 in the tumor. Staining for CK19, CK7, and CAM5.2 was positive in the cytoplasm, and P40 and Ki-67 were positive in the nucleus. In contrast, the expression of hepatocyte marker and AFP was not detected (Figure 4).

FINAL DIAGNOSIS

The final pathological diagnosis was moderately differentiated SCC of the gallbladder.

TREATMENT

Due to the patient's poor physical condition, systemic chemotherapy was not administered.

OUTCOME AND FOLLOW-UP

After five months, the patient was re-admitted to the hospital owing to fever (Table 1). A space-occupying lesion in the right posterior aspect of the first hepatic portal region was observed by abdominal USG, and considered metastatic cancer (Figure 5). Re-examination of CT and magnetic resonance cholangiopancreatography (MRCP) revealed postoperative changes in the GBC and multiple metastases in the liver and abdominal cavity (Figure 6).

DISCUSSION

GBC is one of the most aggressive malignancies of the gastrointestinal tract^[1]. GBC includes several subtypes, including AC, SCC, and ASC. Particularly, SCC has a very low incidence rate and represents only 2%-3% of all GBC cases. The progression and overall prognosis of SCC appear to be worse than those of AC and ASC^[4,8,9]. ASC of




Figure 1 Gallbladder carcinoma infiltrating the adjacent liver, with enlarged retroperitoneal lymph nodes. A: Computed tomography; B: Contrast enhanced computed tomography.

the gallbladder is also rare, and its incidence rate is only 7%-10.6%. ASC includes squamous cells and glandular components. The squamous components that may explain the complex biological behavior and poor prognosis show greater capacities of proliferation and local invasion than the glandular components. Lymphatic metastasis and hematogenous metastasis are the principal types of tumor metastasis.

In clinical practice, pain occurs in 66% of patients, and is the most significant symptom. In patients with GBC, the detection of a palpable mass in the right upper quadrant is not specific, but could be a crucial hint that indicates the presence of a malignancy. Some patients may present with a tumor in the right hypochondrial region. However, it could be clinically misdiagnosed as abscess of the gallbladder. USG is the preferred method for diagnosis, and suspected cases can be diagnosed by CT, percutaneous transhepatic cholangiography, MRCP, and fine needle aspiration^[4]. SCC of the gallbladder has a very poor prognosis, and the involvement of the serosa and lymphatic metastasis are the most important prognostic factors. The size of tumors at diagnosis is the most significant determinant of survival. Therefore, it is especially important to improve the survival index, thus providing useful strategies for early diagnosis and prevention of SCC of the gallbladder^[1].

Studies have shown that surgical resection has been considered as the cornerstone of treatment for many years, and complete resection is associated with increased survival^[10,11]. However, most patients died approximately 6 mo after diagnosis without radical surgery^[4]. Currently, cholecystectomy, hepatectomy, and lymphadenectomy contribute to patient's recovery. Some studies demonstrated that the 5-year survival rate may still be low owing to systemic failure. Weatherall *et al.*^[11] have confirmed that although patients with SCC of the gallbladder have a larger tumor size and higher possibility of involvement of adjacent organs, complete resection is possible^[12]. However, there is limited information on the optimal treatment for SCC of the gallbladder. Therefore, physicians have been striving to optimize assisted therapy and improve survival. After surgical resection, the majority of physicians offer adjuvant chemotherapy using 5-fluorouracil. The benefits of the best adjuvant therapy or accompanying chemoradiotherapy remain controversial.

Various hypotheses have been proposed for the etiology of gallbladder SCC, including: (1) Ectopic squamous epithelium with malignant transformation^[1]; (2) Metaplastic squamous epithelium with malignant transformation which describes the evolution of metaplasia-dysplasia-carcinoma in progressive development^[1,13]. The gallbladder may trigger differentiation of glandular cells into squamous cells due to chronic irritation from gallstones. Thereafter, the squamous metaplastic cells may undergo malignant transformation into tumor cells^[4,8,13-16]; and (3) Adenocarcinoma with squamous metaplasia. Here, squamous cell elements of mixed ASC of the gallbladder undergo excessive growth and eventually replace all the adenocarcinoma components, resulting in SCC development^[4,8,13-16].

The molecular mechanisms for the occurrence and progression of SCC remain unclear. Further studies are needed to determine the etiopathogenetic and molecular differences between this histologic type and conventional adenocarcinomas. Therefore, it is essential to investigate the malignant biological behavior and clinicopathological characteristics of GBSCC. According to studies, CD109 may be associated with the development and clinicopathological characteristics of SCC. CD109 is a glycosylphosphatidylinositol-anchored cell-surface glycoprotein that negatively regulates transforming growth factor- β (TGF- β) signaling as a novel TGF- β co-receptor^[17-22]. CD109 is a member of the α 2-macroglobulin/C3, C4, C5 family of



Figure 2 Gross morphology of a 5 cm × 4 cm × 6 cm tumor seen near the gallbladder neck and liver tissue. Part of the mass was grey and soft, part of the mass section showing purulent necrosis was visible, and the boundary between the mass and surrounding liver tissue was not clear.

thioester-containing proteins^[21,22]. CD109 could be a potential biological marker for SCC of the gallbladder^[23,24]. Furthermore, a higher level expression of the CD109 protein was also reported in SCCs of other organs^[24-27]. TGF- β is a multipotent cytokine implicated in many physiological processes and regulates a wide variety of cellular processes including cell proliferation, differentiation, adhesion, extracellular matrix deposition, wound healing, and inflammation^[18,28-30]. TGF- β signal transduction maintains the homeostasis of epithelial cells^[24,31], and defective TGF- β signaling may lead to cellular hyper-proliferation, reduced apoptosis, and increased genomic instability^[24,32]. The transducing factor of the TGF- β signaling pathway is commonly downregulated in human SCCs^[23,24]. CD109 is a co-receptor of TGF- β 1 and increases the binding of TGF- β 1 to its receptor^[24,33]. CD109 is associated with caveolin-1, which can promote the localization of TGF- β receptors in degraded cavities^[24,34]. Therefore, CD109 regulates TGF- β signaling and enhances TGF- β receptor endocytosis actively. CD109 upregulation in SCC of the gallbladder potentially inhibits TGF- β signal transduction and subsequently promotes SCC progression. In summary, CD109 may be involved in the pathogenesis of gallbladder SCC and serves as a novel target for intervention^[24].

Recent findings demonstrate that the clinicopathological features of SC remain to be well recognized. Consequently, the therapeutic interventions remain to be fully elucidated. Although biomarkers for predicting the prognosis of GBSCC have been recognized, they still lack clinical applicability currently. Therefore, it is vital to record the biological and clinicopathological characteristics of GBSCC.

CONCLUSION

Only scarce reports of SCC of the gallbladder have been reported; GBSCC is a rare subtype, and it has a poor prognosis. The squamous components that may explain the complex biological behavior and poor prognosis show greater capacities of proliferation and local invasion in GBC. Surgical resection has been considered as the basis of treatment for years, and complete resection is associated with increased survival. CD109 may be involved in the pathogenesis of gallbladder SCC and serves as a novel target for intervention.

Table 1 Timeline

Time	Measure	Results
November 8, 2017	Abdominal ultrasonography	A 5.8 cm × 4.5 cm heterogeneous mass localized in both the gallbladder and liver and a gallbladder with remarkably thickened wall, indicating a malignancy
November 15, 2017	CT and contrast-enhanced CT	Gallbladder carcinoma infiltrating the adjacent liver, with enlarged retroperitoneal lymph nodes
November 22, 2017	Radical cholecystectomy	The gallbladder adhered tightly to the greater omentum. Neoplastic entities were present at the bottom of the gallbladder, which had invaded the V hepatic segments of the liver
November 23, 2017	Pathological findings	A moderately differentiated SCC
November 24, 2017	Immunohistochemistry	A moderately differentiated SCC
April 15, 2018	Abdominal ultrasonography	A space-occupying lesion in the right posterior aspect of the first hepatic portal region, which was considered as cancer metastasis
April 16, 2018	CT and magnetic resonance cholangiopancreatography	Postoperative changes in gallbladder carcinoma and multiple metastases in the liver and abdominal cavity

CT: Computed tomography; SCC: Squamous cell carcinoma.




Figure 3 Hematoxylin and eosin staining of tumor cells. A: Image showing diffuse growth replacing the mucosa and invading the muscularis propria; B: Image showing densely packed tumors formed by squamous cells; C: Image showing squamous cell carcinoma with keratin pearl formation; D: Image showing that tumor cell proliferation is markedly atypical, and karyokinesis is increased.




Figure 4 Immunohistochemical staining shows tumor cells in gallbladder squamous cell carcinoma ($\times 100$). A: AE1/AE3 (+); B: CK5/6 (+); C: CAM5.2 (+); D: CK7 (+); E: CK19 (+); F: P40 (+); G: Hepatocyte (-); H: AFP (-); I: Ki-67 (60%+).




Figure 5 Abdominal ultrasonography revealed a space-occupying lesion in the right posterior aspect of the first hepatic portal region, which was considered as cancer metastasis.




Figure 6 Postoperative changes in gallbladder carcinoma and multiple metastases in the liver and abdominal cavity. A and B: Computed tomography; C and D: Magnetic resonance cholangiopancreatography.

REFERENCES

- 1 **Khan N**, Afroz N, Haider N, Khan MA. A case of pure endophytic squamous cell carcinoma of the gallbladder: a rare entity with aggressive behaviour. *Turk Patoloji Derg* 2012; **28**: 181-183 [PMID: 22627641 DOI: 10.5146/tjpath.2012.01122]
- 2 **Yildiz İ**, Koca YS, Barut İ. Overlap of Acute Cholecystitis with Gallstones and Squamous Cell Carcinoma of the Gallbladder in an Elderly Patient. *Case Rep Surg* 2015; **2015**: 767196 [PMID: 26356191 DOI: 10.1155/2015/767196]
- 3 **Yoo Y**, Mun S. Synchronous double primary squamous cell carcinoma and adenocarcinoma of the extrahepatic bile duct: a case report. *J Med Case Rep* 2015; **9**: 116 [PMID: 25986701 DOI: 10.1186/s13256-015-0600-1]
- 4 **Chakrabarti I**, Giri A, Ghosh N. Cytohistopathological correlation of a case of squamous cell carcinoma of gallbladder with lymph node metastasis. *Turk Patoloji Derg* 2014; **30**: 81-84 [PMID: 24101354 DOI: 10.5146/tjpath.2013.01180]
- 5 **Li J**, Yang ZL, Ren X, Zou Q, Yuan Y, Liang L, Chen M, Chen S. ACE2 and FZD1 are prognosis markers in squamous cell/adenosquamous carcinoma and adenocarcinoma of gallbladder. *J Mol Histol* 2014; **45**: 47-57 [PMID: 23921915 DOI: 10.1007/s10735-013-9528-1]
- 6 **Liu Z**, Yang Z, Jiang S, Zou Q, Yuan Y, Li J, Li D, Liang L, Chen M, Chen S. MCM2 and TIP30 are prognostic markers in squamous cell/adenosquamous carcinoma and adenocarcinoma of the gallbladder.

- Mol Med Rep* 2016; **14**: 4581-4592 [PMID: [27748889](#) DOI: [10.3892/mmr.2016.5851](#)]
- 7 **Zhou H**, Tao L, Chen D, Gao R, Zhao L, Li G, Chen H, Ren Y, Li X, Wang Y, Pang L. Immunohistochemical findings in small cell neuroendocrine carcinoma of the cervix: a report of two cases. *Transl Cancer Res* 2017; **6**: 860-868 [DOI: [10.21037/tcr.2017.07.15](#)]
 - 8 **Roppongi T**, Takeyoshi I, Ohwada S, Sato Y, Fujii T, Honma M, Morishita Y. Minute squamous cell carcinoma of the gallbladder: a case report. *Jpn J Clin Oncol* 2000; **30**: 43-45 [PMID: [10770570](#) DOI: [10.1093/jjco/hyd010](#)]
 - 9 **Kendre P**, Kataria P, Patel A, Mittal LC, Mule T. Metastasis as initial presentation of squamous cell carcinoma of gallbladder: A rare clinical entity. *Indian J Pathol Microbiol* 2017; **60**: 440-441 [PMID: [28937396](#) DOI: [10.4103/IJPM.IJPM_113_17](#)]
 - 10 **Oohashi Y**, Shirai Y, Wakai T, Nagakura S, Watanabe H, Hatakeyama K. Adenosquamous carcinoma of the gallbladder warrants resection only if curative resection is feasible. *Cancer* 2002; **94**: 3000-3005 [PMID: [12115390](#) DOI: [10.1002/cncr.10578](#)]
 - 11 **Weatherall TJ**, Fenton M, Munene G, Dickson PV, Deneve JL. Locally Advanced, Unresectable Squamous Cell Carcinoma of the Gallbladder. *Case Rep Surg* 2015; **2015**: 424650 [PMID: [26236536](#) DOI: [10.1155/2015/424650](#)]
 - 12 **Kalayarasan R**, Javed A, Sakhuja P, Agarwal AK. Squamous variant of gallbladder cancer: is it different from adenocarcinoma? *Am J Surg* 2013; **206**: 380-385 [PMID: [23827515](#) DOI: [10.1016/j.amjsurg.2012.12.012](#)]
 - 13 **Alpuerto AC**, Mora ME, Robitsek RJ, Schubl SD. Primary Pure Squamous Cell Carcinoma of the Gallbladder Locally Invading the Liver, Duodenum, and Stomach: A Case Report and Literature Review. *Case Rep Surg* 2017; **2017**: 2534029 [PMID: [28251014](#) DOI: [10.1155/2017/2534029](#)]
 - 14 **Hosseinzadeh M**, Shokripur M, Salahi H. Primary pure squamous cell carcinoma of gallbladder presenting as acute cholecystitis. *Iran J Med Sci* 2012; **37**: 271-273 [PMID: [23390334](#)]
 - 15 **Roa JC**, Tapia O, Cakir A, Basturk O, Dursun N, Akdemir D, Saka B, Losada H, Bagci P, Adsay NV. Squamous cell and adenosquamous carcinomas of the gallbladder: clinicopathological analysis of 34 cases identified in 606 carcinomas. *Mod Pathol* 2011; **24**: 1069-1078 [PMID: [21532545](#) DOI: [10.1038/mod-pathol.2011.68](#)]
 - 16 **Rao S**, Arya A, Aggarwal S, Gupta K, Arora R, Dhawan I. Pure squamous cell carcinoma of the gall bladder. *Indian J Pathol Microbiol* 2007; **50**: 599-600 [PMID: [17883151](#)]
 - 17 **Hagiwara S**, Murakumo Y, Mii S, Shigetomi T, Yamamoto N, Furue H, Ueda M, Takahashi M. Processing of CD109 by furin and its role in the regulation of TGF-beta signaling. *Oncogene* 2010; **29**: 2181-2191 [PMID: [20101215](#) DOI: [10.1038/onc.2009.506](#)]
 - 18 **Li C**, Hancock MA, Sehgal P, Zhou S, Reinhardt DP, Philip A. Soluble CD109 binds TGF-beta and antagonizes TGF-beta signalling and responses. *Biochem J* 2016; **473**: 537-547 [PMID: [26621871](#) DOI: [10.1042/BJ20141488](#)]
 - 19 **Yokoyama M**, Ichinoe M, Okina S, Sakurai Y, Nakada N, Yanagisawa N, Jiang SX, Numata Y, Umezawa A, Miyazaki K, Higashihara M, Murakumo Y. CD109, a negative regulator of TGF-beta signaling, is a putative risk marker in diffuse large B-cell lymphoma. *Int J Hematol* 2017; **105**: 614-622 [PMID: [28032275](#) DOI: [10.1007/s12185-016-2173-1](#)]
 - 20 **Shiraki Y**, Mii S, Enomoto A, Momota H, Han YP, Kato T, Ushida K, Kato A, Asai N, Murakumo Y, Aoki K, Suzuki H, Ohka F, Wakabayashi T, Todo T, Ogawa S, Natsume A, Takahashi M. Significance of perivascular tumour cells defined by CD109 expression in progression of glioma. *J Pathol* 2017; **243**: 468-480 [DOI: [10.1002/path.4981](#)]
 - 21 **Mii S**, Hoshino A, Enomoto A, Murakumo Y, Ito M, Yamaguchi A, Takahashi M. CD109 deficiency induces osteopenia with an osteoporosis-like phenotype in vivo. *Genes Cells* 2018; **23**: 590-598 [PMID: [29767469](#) DOI: [10.1111/gtc.12593](#)]
 - 22 **Qi R**, Dong F, Liu Q, Murakumo Y, Liu J. CD109 and squamous cell carcinoma. *J Transl Med* 2018; **16**: 88 [PMID: [29625613](#) DOI: [10.1186/s12967-018-1461-3](#)]
 - 23 **Reis-Filho JS**, Torio B, Albergaria A, Schmitt FC. p63 expression in normal skin and usual cutaneous carcinomas. *J Cutan Pathol* 2002; **29**: 517-523 [PMID: [12358808](#)]
 - 24 **Dong F**, Lu C, Chen X, Guo Y, Liu J. CD109 is a novel marker for squamous cell/adenosquamous carcinomas of the gallbladder. *Diagn Pathol* 2015; **10**: 137 [PMID: [26249215](#) DOI: [10.1186/s13000-015-0375-0](#)]
 - 25 **Dong F**, Wang Y, Li L, Wang Y, Liu X, Liu J. CD109 expression is increased in cutaneous squamous cell carcinoma. *J Dermatol* 2014; **41**: 947-949 [PMID: [25271095](#) DOI: [10.1111/1346-8138.12620](#)]
 - 26 **Zhang JM**, Hashimoto M, Kawai K, Murakumo Y, Sato T, Ichihara M, Nakamura S, Takahashi M. CD109 expression in squamous cell carcinoma of the uterine cervix. *Pathol Int* 2005; **55**: 165-169 [PMID: [15826242](#) DOI: [10.1111/j.1440-1827.2005.01807.x](#)]
 - 27 **Dong F**, Liu F, Yan S, Liu X, Jiang Z, Liu J. Elevated Expression of CD109 in Esophageal Squamous Cell Carcinoma. *Pathol Oncol Res* 2015; **21**: 1273-1275 [PMID: [26122747](#) DOI: [10.1007/s12253-014-9894-3](#)]
 - 28 **Vorstenbosch J**, Nguyen CM, Zhou S, Seo YJ, Siblini A, Finnson KW, Bizet AA, Tran SD, Philip A. Overexpression of CD109 in the Epidermis Differentially Regulates ALK1 Versus ALK5 Signaling and Modulates Extracellular Matrix Synthesis in the Skin. *J Invest Dermatol* 2017; **137**: 641-649 [PMID: [27866969](#) DOI: [10.1016/j.jid.2016.09.039](#)]
 - 29 **Bizet AA**, Tran-Khanh N, Saksena A, Liu K, Buschmann MD, Philip A. CD109-mediated degradation of TGF-beta receptors and inhibition of TGF-beta responses involve regulation of SMAD7 and Smurf2 localization and function. *J Cell Biochem* 2012; **113**: 238-246 [PMID: [21898545](#) DOI: [10.1002/jcb.23349](#)]
 - 30 **Man XY**, Finnson KW, Baron M, Philip A. CD109, a TGF-beta co-receptor, attenuates extracellular matrix production in scleroderma skin fibroblasts. *Arthritis Res Ther* 2012; **14**: R144 [PMID: [22694813](#) DOI: [10.1186/ar3877](#)]
 - 31 **Taylor MA**, Lee YH, Schiemann WP. Role of TGF-beta and the tumor microenvironment during mammary tumorigenesis. *Gene Expr* 2011; **15**: 117-132 [PMID: [22268294](#)]
 - 32 **White RA**, Malkoski SP, Wang XJ. TGF-beta signaling in head and neck squamous cell carcinoma. *Oncogene* 2010; **29**: 5437-5446 [PMID: [20676130](#) DOI: [10.1038/onc.2010.306](#)]
 - 33 **Finnson KW**, Tam BY, Liu K, Marcoux A, Lepage P, Roy S, Bizet AA, Philip A. Identification of CD109 as part of the TGF-beta receptor system in human keratinocytes. *FASEB J* 2006; **20**: 1525-1527 [PMID: [16754747](#) DOI: [10.1096/fj.05-5229fje](#)]
 - 34 **Bizet AA**, Liu K, Tran-Khanh N, Saksena A, Vorstenbosch J, Finnson KW, Buschmann MD, Philip A. The TGF-beta co-receptor, CD109, promotes internalization and degradation of TGF-beta receptors. *Biochim Biophys Acta* 2011; **1813**: 742-753 [PMID: [21295082](#) DOI: [10.1016/j.bbamer.2011.01.028](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

