

Name of the Journal: *World journal of Gastroenterology*

Manuscript NO: 49807

Manuscript Type: RANDOMIZED CLINICAL TRIAL

Individualized³ home-monitoring of disease activity in adult patients with inflammatory bowel disease can be recommended in clinical practice: A randomized-clinical trial

Ankersen DV *et al.* Home-monitoring in IBD

Abstract

BACKGROUND

The optimal way to home-monitor patients with inflammatory bowel disease (IBD) for disease progression or relapse remains to be found.

Match Overview

1	Internet 21 words crawled on 04-Oct-2018 bmcgastroenterol.biomedcentral.com	<1%
2	Internet 12 words crawled on 22-Aug-2019 www.scribd.com	<1%
3	Crossref 12 words "Biomarkers in Inflammatory Bowel Diseases", Springer Science and Business Media LLC, 2019	<1%

검색결과 약 763,000개 (0.69초)

청소년에게 유해한 결과는 제외되었습니다. 만 19세 이상의 사용자는 성인인증을 통해 모든 결과를 볼 수 있습니다.

사용자 성인인증

DOP031 Individualised home-monitoring of disease activity ... - Ecco-ibd

<https://www.ecco-ibd.eu/.../dop031-individualised-home-monitori...> ▼ 이 페이지 번역하기

DOP031 Individualised home-monitoring of disease activity in adult patients with inflammatory bowel disease can be recommended implemented in clinical practice ... IBD patients were randomised in a 1-year open-label telemedicine study to ...

DOP031 Individualised home-monitoring of disease activity in adult ...

<https://academic.oup.com/ecco-jcc/article-abstract/.../4807941> - 이 페이지 번역하기

A Heida 저술 - 2018

2018. 1. 16. - Due to the chronic relapsing nature of Inflammatory bowel disease (IBD) it is of ... disease activity in adult patients with inflammatory bowel disease can be recommended implemented in clinical practice ... A total of 120 adult IBD patients were randomised in a 1-year open-label telemedicine study to screen ...

DOP031 Individualised home-monitoring of disease activity in adult ...

https://www.researchgate.net/.../322890037_DOP031_Individuali... - 이 페이지 번역하기

DOP031 Individualised home-monitoring of disease activity in adult patients with inflammatory bowel disease can be recommended implemented in clinical practice ... Methods A total of 120 adult IBD patients were randomised in a 1-year open-label telemedicine study to screen for disease activity OD or every 3M using a ...


Evidence-based clinical practice guidelines for inflammatory bowel



国内版

国际版

Individualized home-monitoring of disease activity in adult pa



Sign in



All

Images

Videos

翻译成中文

关闭取词

28,400 Results

Any time ▾

TELEmedicine for Patients with Inflammatory Bowel Disease ...

<https://www.sciencedirect.com/science/article/pii/S1551714415000610>

The TELEmedicine for Patients with Inflammatory **Bowel Disease** (TELE-IBD) study is a multicenter, randomized, clinical trial conducted over 12 months. Participants in the intervention arms complete self-testing with **TELE-IBD** on either a weekly or every other week basis based on ...

Cited by: 33

Author: Raymond K. Cross, Guruprasad Jambaul...

Publish Year: 2015

Elderly patients and inflammatory bowel disease

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4734955>

Feb 06, 2016 · Core tip: **Inflammatory bowel disease** can be mis-diagnosed as its **clinical** features are similar in younger and elderly **patients**. Therapeutic regimes may be different with elderly **patients** being less likely to have immunosuppressant drugs and Anti-TNF's either driven by clinican or patient preference and **disease** related factors.

Cited by: 19

Author: Danielle Nimmons, Jimmy K Limdi

Publish Year: 2016

Is home monitoring of inflammatory bowel disease feasible ...

https://www.researchgate.net/publication/334184336_Is_home_monitoring_of_inflammatory...

Background: **Fecal calprotectin** is a reliable surrogate marker for inflammatory activity in inflammatory bowel disease (IBD).

American Gastroenterological Association Institute ...

[https://www.gastrojournal.org/article/S0016-5085\(17\)35963-2/fulltext](https://www.gastrojournal.org/article/S0016-5085(17)35963-2/fulltext)

This document presents the official recommendations of the American Gastroenterological Association (AGA) on therapeutic drug monitoring (TDM) in **inflammatory bowel disease** (IBD). The guideline was developed by the AGA's **Clinical** Guidelines Committee and approved by the AGA Governing Board. It is accompanied by a Technical Review, which is a compilation of **clinical** evidence from which these ...

Cited by: 134

Author: Joseph D. Feuerstein, Geoffrey C. Nguye...

Publish Year: 2017



国内版

国际版

Individualized home-monitoring of disease activity in adult patients with inflamm



All

Images

Videos

翻译成中文

关闭取词

28,000 Results

Any time ▾

TELEmedicine for Patients with Inflammatory Bowel Disease ...

<https://www.sciencedirect.com/science/article/pii/S1551714415000610>

The TELEmedicine for **Patients with Inflammatory Bowel Disease** (TELE-IBD) study is a multicenter, randomized, **clinical trial** conducted over 12 months. Participants in the intervention arms complete self-testing with TELE-IBD on either a weekly or every other week ...

Cited by: 33

Author: Raymond K. Cross, Guruprasad Jambaulik...

Publish Year: 2015

Is home monitoring of inflammatory bowel disease feasible ...

https://www.researchgate.net/publication/334184336_Is_home_monitoring_of_inflammatory...

activity between the **home monitoring** and control **patients**, ... implementation of randomized **clinical trial**. Contemp Clin Trials. ... **Patients with inflammatory bowel disease** need close monitoring ...

Elderly patients and inflammatory bowel disease

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4734955>

Feb 06, 2016 - Core tip: **Inflammatory bowel disease** can be mis-diagnosed as its **clinical** features are similar in younger and elderly **patients**. Therapeutic regimes may be different with elderly **patients** being less likely to have immunosuppressant drugs and Anti-TNF's either driven by clinican or patient preference and **disease** related factors.

Cited by: 19

Author: Danielle Nimmons, Jimmy K Limdi

Publish Year: 2016

TNF Blocking Therapies and Immunomonitoring in Patients ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3976924>

Since their appearance in the armamentarium for **inflammatory bowel disease** (IBD) more than a decade ago, antitumor necrosis factor (TNF) inhibitors have demonstrated beneficial **activity** in induction and maintenance of **clinical** remission, mucosal healing, improvement in quality of life, and reduction in surgeries and hospitalizations.

Cited by: 20

Author: Romain Altwegg, Thierry Vincent

Publish Year: 2014

Academic Editor: Sophie Desplat-J&#x000e9;go

ECCO IBD Curriculum: 3. Assessment

<https://e-learning.ecco-ibd.eu/mod/data/view.php?d=149&advanced=0&paging&page=5> ▾

The assessment of **disease activity** and extent including complications ... **Individualized home-monitoring of disease activity in adult patients with Inflammatory Bowel Disease** can be recommended implemented in **clinical practice**. D. Ankersen. ECCO'18 Vienna. PPT . Topic 3.1; 3.4 ; ...