



All

Images

Videos

翻译成中文

关闭取词

40,200 Results

Any time ▾

Imaging Findings of Intrahepatic Bile Duct Adenoma ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2930166>

Aug 27, 2010 - Intrahepatic BDA is a rare benign epithelial tumor of the liver that is usually incidentally found at surgery or autopsy (1-4). Its incidence was reported to make up 1.3% of primary liver tumors (2). Craig et al. (3) reported only five cases of intrahepatic BDA in 50,000 autopsies, and until now, Allaire et al.

Intrahepatic bile duct adenoma (peribiliary gland ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4503188>

May 01, 2015 - Introduction. Intrahepatic bile duct adenoma (BDA) is a rare benign liver tumor arising from the epithelium of the intrahepatic bile ducts. Bhathal et al [1] suggests that BDA should be called a peribiliary gland hamartoma because of a similarity between bile duct adenoma and peribiliary glands in their secretory gland cell phenotype.

Author: Jianguo Wei, Dongmei Zhang, Jianfen... Publish Year: 2015

The first case of pediatric bile duct adenoma - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/S2213576614001687>

Intrahepatic bile duct adenoma (BDA) is a rare benign epithelial liver tumor derived from bile duct cells. BDA represents about 1.3% of all primary liver tumors and it is mainly found in adults incidentally on the surface of the liver at laparotomy or autopsy . To date, this case is the first pediatric bile duct adenoma all over the world . 2.1.

Author: Zhi Li, Xiaoyi Sun, Jiexiong Feng Publish Year: 2015

(PDF) The first case of pediatric bile duct adenoma

https://www.researchgate.net/.../272381692_The_first_case_of_pediatric_bile_duct_adenoma

PDF | Intrahepatic bile duct adenoma (BDA) is a rare benign epithelial liver tumor derived from bile duct cells. We report the first case of pediatric bile duct adenoma in the world.

Common bile duct villous adenoma: a case report and review ...

<https://jmedicalcasereports.biomedcentral.com/articles/10.1186/s13256-015-0789-z> ▾

Jan 21, 2016 - According to the literature, benign bile duct tumors are exceedingly uncommon. To the best of our knowledge, we report the largest extrahepatic bile duct villous adenoma described in the literature. We present a case of a 77-year-old Caucasian woman with obstructive jaundice. Laboratory tests revealed that she had elevated bilirubin and liver enzyme levels.



All

Images

Videos

关闭取词

42,100 Results

Any time ▾

Intrahepatic bile duct adenoma is a **rare** benign epithelial hepatic **tumor** derived from **bile duct** cells. We **report** the imaging findings of a patient with **bile duct adenoma**, which appeared as a small heterogeneously enhancing mass with focal small cystic change on CT and MRI.

Imaging Findings of Intrahepatic Bile Duct Adenoma ...

www.ncbi.nlm.nih.gov/pmc/articles/PMC2930166/

Is this answer helpful?  

The first case of pediatric bile duct adenoma - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/S2213576614001687>

2. Discussion. **Intrahepatic bile duct adenoma** (BDA) is a **rare** benign epithelial **liver tumor** derived from **bile duct** cells. BDA represents about 1.3% of all primary **liver tumors** and it is mainly found in adults incidentally on the surface of the **liver** at laparotomy or autopsy .To date, this **case** is the first **pediatric bile duct adenoma** all over the world .

Author: Zhi Li, Xiaoyi Sun, Jiexiong Feng **Publish Year:** 2015

(PDF) The first case of pediatric bile duct adenoma

https://www.researchgate.net/.../272381692_The_first_case_of_pediatric_bile_duct_adenoma

PDF | **Intrahepatic bile duct adenoma** (BDA) is a **rare** benign epithelial **liver tumor** derived from **bile duct** cells. We **report** the first **case** of **pediatric bile duct adenoma** in the world. Furthermore ...

3
 Name of Journal: *World Journal of Gastrointestinal Surgery*
 Manuscript NO: 50147
 Manuscript Type: CASE REPORT

Pediatric intrahepatic bile duct adenoma - rare liver tumor: A case report

Roy AK *et al.* Pediatric intrahepatic bile duct adenoma

Anup Kumar Roy, Nriyten Nath Das

Abstract

BACKGROUND

Intrahepatic bile duct adenoma (BDA) is one of the rarest of the rare benign tumors of liver in the pediatric age group. It arises from the epithelial lining of intrahepatic bile ducts. The overall incidence of BDA is 1.3% of all primary benign liver tumors. Few case reports of this rare tumor occurring in adult population are present in the literature but till date only one pediatric case has been reported in the world.

CASE SUMMARY

An 18 months old male child presented with chief complaints of mass per abdomen for 8 months. Computerized tomography abdomen revealed a well-defined exophytic solid tumor arising from the posteroinferior margin of right lobe of liver with heterogenous enhancement and cystic changes suggesting a suspicion of hepatoblastoma. Non anatomical liver resection was done for the same and a tumor of size 10 cm × 9.5 cm was excised. Histopathology of the specimen was conclusive with the diagnosis of intrahepatic bile duct adenoma which was further supported by immunohistochemistry panel testing. Post-operative period was uneventful. On follow up the child was doing

Match Overview

1	Crossref 79 words William W. Wu, Mai Gu, Di Lu. "Cytopathologic, Histopathologic, and Immunohistochemical Features of Intrahepatic Cle ...	5%
2	Crossref 54 words Arthur Zimmermann. "Chapter 41-1 Tumors and Tumor-like ... esions of Peribiliary Glands", Springer Science and Business	3%
3	Internet 44 words crawled on 14-Sep-2019 f6publishing.blob.core.windows.net	3%
4	Crossref 14 words Zhi Li, Xiaoyi Sun, Jiexiong Feng. "The first case of pediatri ... bile duct adenoma", Journal of Pediatric Surgery Case Repor	1%
5	Crossref 14 words Kyoung Doo Song, Woo Kyoung Jeong. "Benign nodules n ... micking hepatocellular carcinoma on gadoxetic acid-enhanc	1%



All

Images

Videos

关闭取词

42,100 Results

Any time ▾

The first case of pediatric bile duct adenoma - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/S2213576614001687>

Intrahepatic bile duct adenoma (BDA) is a **rare benign epithelial liver tumor** derived from **bile duct cells**. **BDA** represents about 1.3% of all primary **liver tumors** and it is mainly found in adults incidentally on the surface of the **liver** at laparotomy or autopsy . To date, this **case** is the first **pediatric bile duct adenoma** all over the world . 2.1.

Author: Zhi Li, Xiaoyi Sun, Jiexiong Feng **Publish Year:** 2015

The first case of pediatric bile duct adenoma - ScienceDirect

<https://www.sciencedirect.com/science/article/abs/pii/S2213576614001687>

Intrahepatic bile duct adenoma (BDA) is a **rare benign epithelial liver tumor** derived from **bile duct cells**. We report the first **case** of **pediatric bile duct adenoma** in the world. Furthermore, we review the diagnosis, pathology, **treatment** and prognosis of **bile duct adenoma**.

Author: Zhi Li, Xiaoyi Sun, Jiexiong Feng **Publish Year:** 2015

(PDF) The first case of pediatric bile duct adenoma

https://www.researchgate.net/.../272381692_The_first_case_of_pediatric_bile_duct_adenoma

PDF | **Intrahepatic bile duct adenoma (BDA)** is a **rare benign epithelial liver tumor** derived from **bile duct cells**. We report the first **case** of **pediatric bile duct adenoma** in the world.

Bile duct adenoma: a case report and literature review ...

<https://wjso.biomedcentral.com/articles/10.1186/1477-7819-12-125> ▾

Apr 26, 2014 · The **patient underwent a liver tumor resection**, and the histological examination revealed **bile duct adenoma (BDA)**. **BDA** is an extremely **rare benign tumor**, which is difficult to distinguish **BDA** from **hepatocellular carcinoma** definitely preoperatively, **surgical resection** is needed as a way of treatment.

Author: Lei Chen, Mai Yu Xu, Feng Chen

Published in: [World Journal of Surgical Oncology](#) · 2014

Authors: [Lei Chen](#) · [Mai Yu Xu](#) · [Feng Chen](#)