



All

Images

Videos

开启取词

82,700 Results

Any time ▾

## The Futility of Systematic Lymphadenectomy in Early-Stage ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5984865>

Lymphadenectomy is considered an integral part of comprehensive surgical staging of endometrial cancer but debate on the value of lymphadenectomy continues in early-stage endometrial cancer. The aim of our study was to determine the number of node positive patients in clinically early-stage low-grade (G1–G2) endometrioid endometrial cancer.

Cited by: 1

Author: Nidhi Nayyar, Prerna Lakhwani, Ashish G...

Publish Year: 2018

## Prognostic factors for and prognostic value of mesenteric ...

<https://www.sciencedirect.com/science/article/pii/S0748798311006494>

Unfortunately, in most cases, invasive epithelial ovarian cancer continues to be discovered at an advanced-stage.

Cited by: 7

Author: S. Gouy, J. Goetgheluck, C. Uzan, J. Du...

Publish Year: 2012

## Article: Cytoreductive surgery in ovarian cancer

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2151328>

Dec 17, 2007 · Unfortunately ovarian cancers are usually discovered at an advanced stage (FIGO stages III and IV) resulting in the poor overall prognosis of this condition. The 5-year survival rate of advanced ovarian cancer is 20–30% in advanced stage disease and the treatment involves surgery and platinum-based chemotherapy.

Cited by: 25

Author: Christophe Pomel, Arjun Jeyarajah, David...

Publish Year: 2007

## Aortic lymphadenectomy in cervical cancer: Anatomy ...

[https://www.gynecologiconcology-online.net/article/S0090-8258\(07\)00500-8/fulltext](https://www.gynecologiconcology-online.net/article/S0090-8258(07)00500-8/fulltext)

Feb 19, 2014 · The left ureter is identified at the pelvic brim, separated from the ovarian pedicle, and displaced with a retractor out of the field of dissection. At this point, paraaortic lymphadenectomy can start. Removal of the most cranial nodes, which are located both behind and under the left renal vein is the most challenging part of aortic lymphadenectomy.

Cited by: 5

Author: Pierluigi Benedetti Panici, Marco Calcagn...

Publish Year: 2007

## Primary Surgery or Neoadjuvant Chemotherapy in Advanced ...

**Name of Journal:** *World Journal of Surgical Procedures*

**Manuscript NO:** 50267

**Manuscript Type:** LETTER TO THE EDITOR

**Regional lymphadenectomy in advanced ovarian cancer: Enigma continues**

Pankaj Kumar Garg, Sandipan Chowdhury, Rajkumar K Seenivasagam, Dharma Ram Poonia

**Abstract**

The role of regional lymphadenectomy has always been a matter of discussion in the surgical management of solid tumors – Pelvic and para-aortic lymphadenectomy in ovarian cancer is one such issue. A recently published randomized trial suggested that regional lymphadenectomy in advanced ovarian cancer is unlikely to offer a survival

Match Overview

There are no matching sources for this report.



83,200 Results

Any time ▾

## Update on lymphadenectomy in early and advanced ovarian ...

[https://www.researchgate.net/publication/5654536\\_Update\\_on\\_lymphadenectomy\\_in\\_early...](https://www.researchgate.net/publication/5654536_Update_on_lymphadenectomy_in_early...)

Conclusion: Lymphadenectomy is associated with greater 5-year overall survival in patients with early and advanced stage ovarian cancer, but an effect on progression-free survival and recurrence ...

## The Futility of Systematic Lymphadenectomy in Early-Stage ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5984865>

Lymphadenectomy is considered an integral part of comprehensive surgical staging of endometrial cancer but debate on the value of lymphadenectomy continues in early-stage endometrial cancer. The aim of our study was to determine the number of node positive patients in clinically early-stage low-grade (G1-G2) endometrioid endometrial cancer.

Cited by: 1

Author: Nidhi Nayyar, Prerna Lakhwani, Ashish G...

Publish Year: 2018

## Ovarian cancer - The Lancet

[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(13\)62146-7/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(13)62146-7/fulltext)

Epithelial ovarian cancer is the commonest cause of gynaecological cancer-associated death. The disease typically presents in postmenopausal women, with a few months of abdominal pain and distension. Most women have advanced disease (International Federation of Gynecology and Obstetrics [FIGO] stage III), for which the standard of care remains surgery and platinum-based cytotoxic chemotherapy.

## Prognostic factors for and prognostic value of mesenteric ...

<https://www.sciencedirect.com/science/article/pii/S0748798311006494>

Unfortunately, in most cases, invasive epithelial ovarian cancer continues to be discovered at an advanced-stage.

Cited by: 7

Author: S. Gouy, J. Goetgheluck, C. Uzan, J. Du...

Publish Year: 2012

## Primary Surgery or Neoadjuvant Chemotherapy in Advanced ...

[https://ascopubs.org/doi/full/10.1200/EDBK\\_160624](https://ascopubs.org/doi/full/10.1200/EDBK_160624)

Oct 29, 2018 · According to the National Comprehensive Cancer Network, the current primary treatment of advanced epithelial ovarian cancer is optimal cytoreductive surgery followed by six to eight cycles of dual-agent platinum- and taxane-based chemotherapy. 5 Cytoreductive surgery (also called "debulking")