

Match Overview



1	Internet 16 words crawled on 26-Aug-2016 mdanderson.influent.utsystem.edu	1%
2	Crossref 14 words Laco, J. "Tumor-to-tumor metastasis: Warthin tumor as a recipient of lung carcinoma and of renal carcinoma - Rep...	1%
3	Crossref 12 words Hadeel Jawad, Peter McCarthy, Gerard O'Leary, Cynthia C. Heffron. "Presentation of Chronic Lymphocytic Leukemia/S	<1%

1 **Name of Journal:** *World Journal of Clinical Cases*

2 **Manuscript NO:** 50555

3 **Manuscript Type:** CASE REPORT

4

5 **Diffuse large B-cell lymphoma arising from follicular lymphoma with**
6 **warthin's tumor of the parotid gland - immunophenotypic and genetic**
7 **features: A case report**

8

9 Wang CS *et al.* DLBCL with WT

10

11 Chang-Song Wang, Xia Chu, Di Yang, Lei Ren, Nian-Long Meng, Xue-Xia Lv,
12 Tian Yun, Yan-Sha Cao

13



国内版

国际版

Diffuse large B-cell lymphoma arising from follicular lymphoma with warthin's tum



All

Images

Videos

关闭取词

9,340 Results

Any time ▼

Characterization of the lymphoid stroma in Warthin's tumor ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2780035>

Apr 05, 2009 · It has been reported that Warthin's tumor is rarely associated with malignant lymphoma. Most of these were low-grade lymphomas, including MALT-type lymphoma and follicular lymphoma. They are also associated with diffuse large B cell lymphoma, Hodgkin's disease, and T-cell lymphoma [2–17]. This relationship between Warthin's tumor and lymphomas gave the impression that these lymphomas might arise in the lymphoid stroma of Warthin's ...

Cited by: 6

Author: Kunchang Song, James D. Cotelingam, ...

Publish Year: 2009

Diffuse Large B-Cell Lymphoma Arising in Warthin's Tumor ...

https://www.researchgate.net/publication/259722619_Diffuse_Large_B-Cell_Lymphoma...

The patient was diagnosed with "Warthin's tumor and diffuse large B-cell lymphoma with expression of CD30." The histopathologic and clinical features are discussed

Follicular lymphoma frequently originates in the salivary ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/j.1440-1827.2006.02011.x>

Sep 20, 2006 · A total of 27 primary salivary gland lymphomas were examined: 6 FL (five, grade 1; one, grade 2); 19 MALT lymphomas; and two diffuse large B-cell lymphomas. The FL patients ranged in age from 24 to 73 years, with a mean of 49 years, which was younger than that of MALT patients (mean: 64 years; $P < 0.05$). Four of the six FL arose from the submandibular gland, which ...

Cited by: 24

Author: Satoko Nakamura, Koichi Ichimura, Yumi...

Publish Year: 2006

[PDF] Malignant lymphoma on parotid gland: a clinical case

<https://synapse.koreamed.org/Synapse/Data/PDFData/3070JKAOMS/jkaoms-43-138.pdf>

Parotid gland lymphoma is most likely to be B-cell, non-Hodgkin's lymphoma of one of three types, which include follicular, marginal zone, and diffuse large B-cell, although other histologic patterns have been described.

[国内版](#)[国际版](#)[All](#)[Images](#)[Videos](#)[关闭取词](#)

9,350 Results

Any time ▾

Characterization of the lymphoid stroma in Warthin's tumor ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2780035>

Apr 05, 2009 · Warthin's tumor is rarely associated with malignant lymphoma. Only 18 cases were reported in the literature so far. In most cases the latter is a low grade process, including Marginal zone/Mucosa associated lymphoid tissue (MALT) type lymphoma, follicular lymphoma, and rarely diffuse large cell lymphoma which may arise de novo or secondary to low grade lymphoma.

Cited by: 6

Author: Kunchang Song, James D. Cotelingam, ...

Publish Year: 2009

Diffuse Large B-Cell Lymphoma Arising in Warthin's Tumor ...

https://www.researchgate.net/publication/259722619_Diffuse_Large_B-Cell_Lymphoma...

Diffuse Large B-Cell Lymphoma Arising in Warthin's Tumor: Case Study and Review of the Literature Article (PDF Available) in The Korean Journal of Pathology 47(6):579-82 · December 2013 with 126 ...

Follicular lymphoma frequently originates in the salivary ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/j.1440-1827.2006.02011.x>

Sep 20, 2006 · In two (33%) of the six FL patients, the tumor had originated in the parotid gland, and in four patients (67%), in the submandibular gland. In 14 patients (74%) with MALT lymphoma, the tumor had originated in the parotid gland, and in five patients (26%), in the submandibular gland.

Cited by: 24

Author: Satoko Nakamura, Koichi Ichimura, Yumi...

Publish Year: 2006

Malignant lymphoma involving a warthin's tumor: A case ...

<https://www.sciencedirect.com/science/article/pii/S0046817790901825>

Case Studies MALIGNANT LYMPHOMA INVOLVING A WARTHIN'S TUMOR: A CASE WITH IMMUNOPHENOTYPIC AND GENE REARRANGEMENT ANALYSIS L. JEFFREY MEDEIROS, MD, RITA Rizzi, MD, PILAR LARDELLI, MD, AND ELAINE S. JAFFE, MD We describe a Warthin's tumor which was involved by malignant lymphoma.

Cited by: 25

Author: L. Jeffrey Medeiros, Rita Rizzi, Pilar Lard...

Publish Year: 1990

[PDF] Malignant lymphoma on parotid gland: a clinical case

<https://synapse.koreamed.org/Synapse/Data/PDFData/3070JKAOMS/jkaoms-43-138.pdf>

examining parotid swellings and masses. Parotid gland lymphoma is most likely to be B-cell, non-Hodgkin's lymphoma of one of three types, which include follicular, marginal zone, and diffuse large