

검색결과 약 26,000,000개 (0.53초)

## Clinical characteristics and treatment outcomes in patients with liver cirrhosis and lymphoma에 대한 학술자료

Clinical features and outcome of patients with non- ... - Shaw - 1844회 인용

Hepatitis B infection in patients with lymphomas - Liang - 150회 인용

Clinical presentation, course, and prognostic factors in ... - Diehl - 372회 인용

## Clinical features and treatment outcomes in patients with ...

<https://www.ncbi.nlm.nih.gov/articles/PMC3974951> - 이 페이지 번역하기

BW Kang 저술 - 2014 - 13회 인용 - 관련 학술자료

2014. 3. 24. - **Clinical features and characteristics** A total of 105 **patients** (80.1%) had stage III or IV MCL at diagnosis. ... According to the simplified IPI, 18 **patients** (13.7%) were at high risk. B **symptoms** were noted in 34 **patients** (26.0%). The overall incidence of extranodal involvement was 69.5%.

## (PDF) Clinical features and treatment outcomes in patients ...

<https://www.researchgate.net/publication/2616056...> - 이 페이지 번역하기

Hide. Abstract. We investigated the **clinical features and treatment outcomes** of **patients** with mantle cell **lymphoma** (MCL) in Korea. .... **lymphoma** in Korea: Study by the Consortium for Improving Survival .... sweats, and weight loss), presence of extranodal **disease**, ..... extranodal sites included the lungs, **liver**, and spleen, al-.

## Clinical characteristics and outcome of pediatric patients with ...

1

**Name of Journal:** *World Journal of Hepatology*

**Manuscript NO:** 51349

**Manuscript Type:** ORIGINAL ARTICLE

*Case Control Study*

**Clinical** characteristics and treatment outcomes in patients with liver cirrhosis and lymphoma

González-Regueiro JA *et al.* Outcomes in liver cirrhosis and lymphoma

José A González-Regueiro, Astrid Ruiz-Margáin, Mariana Cruz-Contreras, Ana M Montaña-Duclaud, Andrea Cavazos-Gómez, Roberta Demichelis-Gómez, Ricardo U Macías-Rodríguez

**Abstract**

**BACKGROUND**

A significant number of patients with liver cirrhosis develop concomitantly some type of solid or hematological cancer, including lymphoma. Treatment of patients with lymphoma and cirrhosis is challenging for physicians, due to the clinical characteristics related to cirrhosis, including biochemical and functional abnormalities, as well as portal hypertension and lack of scientific evidence, limiting the use of chemotherapy. Currently, experts recommend only offering oncological treatment to patients with compensated cirrhosis.

Match Overview

1

**Internet** 31 words  
crawled on 06-Oct-2019  
[f6publishing.blob.core.windows.net](http://f6publishing.blob.core.windows.net)

1%

2

**Internet** 14 words  
crawled on 21-Sep-2017  
[journals.plos.org](http://journals.plos.org)

<1%

3

**Crossref** 12 words  
"Poster Session 3: Clinical and Translational Fibrosis Rese...  
ch; Encephalopathy and Other Complications of Cirrhosis",

<1%





All

Images

Videos

关闭取词

2,550,000 Results

Any time ▾

## Clinical characteristics of liver failure with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6544643>

May 31, 2019 - **Liver** failure with hemophagocytic lymphohistiocytosis (HLH) is a life-threatening syndrome with high mortality. The aim of this study was to decipher **clinical** and laboratory **characteristics** of hemophagocytic lymphohistiocytosis after definite diagnosis of **liver** failure and to provide clues for early diagnosis **and treatment** of HLH in **patients with liver** failure.

## Clinical Characteristics and Outcome of Hepatic ...

<https://www.ncbi.nlm.nih.gov/pubmed/28872150>

**Clinical Characteristics and Outcome of Hepatic Sarcoidosis**: A Population-Based Study 1976-2013. Ungprasert P(1)(2), Crowson CS(1)(3), Simonetto DA(4), Matteson EL(1)(5). Author information: (1)Division of **Rheumatology**, Department of **Internal Medicine**, Mayo Clinic College of **Medicine** and Science, Mayo **Clinic**, Rochester, Minnesota, USA.

## [PDF] Primary lymphoma of the liver: clinical features and ...

<https://jbuon.com/archive/10-4-505.pdf>

**B-type lymphoma** (Table 1). Most **patients** (5/9) were found with stage IV dis-ease. Only 2 **patients** (No. 2 and 6) presented with stage I disease (**solitary liver lesion**). Seven **patients** were **treated** with **CHOP** (**cyclophosphamide**, doxorubicin, vincristine, prednisone) combination chemotherapy. Complete and good partial remissions were achieved

## Non-Hodgkin's Lymphoma Involving the Liver: Clinical and ...

<https://www.sciencedirect.com/science/article/pii/S1557919011704024>

**Clinical Lymphoma** & Myeloma January 2006 273 Introduction Non-Hodgkin's **lymphoma** (NHL) represents the fifth most common cancer type in the United States, with an incidence that has been increasing rapidly in Western countries. 1 Lymphomas are a diverse collection of different diseases with varying histologies and **clinical** presentations.

**Cited by:** 68**Author:** J. Stuart Salmon, Mary Ann Thompson, R...**Publish Year:** 2006

## Lymphoma of the liver: Clinicopathological features of 19