



All

Images

Videos

关闭取词

25,200 Results

Any time ▾

Neoadjuvant Chemoradiation Therapy Using Concurrent S-1 ...

<https://www.sciencedirect.com/science/article/pii/S0360301614002958>

Jul 01, 2014 · NCRT with concurrent S-1/irinotecan produced high response rates and excellent long-term survival, with acceptable adverse effects in patients with rectal cancer. ypN2 is a strong predictor of dismal outcomes, and a combination of ypT and tumor location can identify high-risk patients among those with ypN0/1 disease.

Cited by: 23

Author: Takatoshi Nakamura, Keishi Yamashita, T...

Publish Year: 2014

A multicenter phase I study of preoperative ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5013752>

In a phase II trial performed by Sato et al., chemoradiotherapy with S-1 plus irinotecan had a histopathological complete response (pCR) rate of 34.7% in patients with locally advanced rectal cancer. However, the clinical target volume for the primary tumor used typically included the ...

Cited by: 5

Author: Takeo Sato, Kazushige Hayakawa, Naohi...

Publish Year: 2016

Preoperative Concurrent Chemoradiotherapy for Locally ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3394859>

Purpose. This study was designed to investigate the long-term oncologic outcomes for locally advanced rectal cancer patients after treatment with preoperative concurrent chemoradiotherapy followed by total mesorectal excision, and to identify prognostic ...

Cited by: 15

Author: Moonkyoo Kong, Seong Eon Hong, Woo ...

Publish Year: 2012

A Review of Neoadjuvant Chemoradiotherapy for Locally ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4971740>

Jul 17, 2016 · Preoperative concurrent radiochemotherapy (nCRT) have become the standard treatment for resectable stage II ~ III patients. For unresectable locally advanced rectal cancer, preoperative concurrent chemoradiotherapy is the only standard treatment, and most of these patients become resectable after the nCRT.

Cited by: 25

Author: Yi Li, Ji Wang, Xiaowei Ma, Li Tan, Yanli ...

Publish Year: 2016

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 51571

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Impact of preoperative chemoradiotherapy using concurrent S-1 and CPT-11 on long-term clinical outcomes in locally advanced rectal cancer

Kimura K *et al.* S-1, CPT-11, and CRT for rectal cancer

Kei Kimura, Naohito Beppu, Hiroshi Doi, Kozo Kataoka, Tomoki Yamano, Motoi

Match Overview

Match Number	Source Type	Words	Similarity
1	Crossref	74 words	2%
	Kei Kimura, Tomoki Yamano, Masataka Igeta, Ayako Imada et al. "polymorphisms in rectal cancer associated v ..."		
2	Internet	26 words crawled on 26-Jul-2019 www.ncbi.nlm.nih.gov	1%
3	Internet	21 words crawled on 05-Nov-2019 academic.oup.com	1%
4	Internet	20 words crawled on 20-Nov-2017 ash.confex.com	1%
5	Crossref	17 words	1%
	Sato, Takeo, Kazushige Hayakawa, Naohiro Tomita, Masafumi Noda, Norihiko Kamikonya, Toshiaki Watanabe, D		
6	Crossref	17 words	1%
	Naohito Beppu, Masayoshi Kobayashi, Nagahide Matsubara, Masashi Noda et al. "Comparison of the pathologi		
7	Internet	17 words crawled on 05-Dec-2019 f6publishing.blob.core.windows.net	1%
8	Crossref	15 words	<1%
	Kazushige Kawai, Eiji Sunami, Keisuke Hata, Toshiaki Tanaka, Takeshi Nishikawa, Kensuke Otani, Kazuhito Sa		
9	Crossref	14 words	<1%
	T. Akiyoshi, S. Toda, T. Tominaga, K. Oba et al. "Prognostic impact of residual lateral lymph node metastasis afi ..."		
10	Internet	14 words crawled on 11-Dec-2019 www.nejm.org	<1%
11	Internet	14 words crawled on 18-Nov-2019 www.jstage.jst.go.jp	<1%

国内版

国际版

Impact of preoperative chemoradiotherapy using concurrent S



ALL

IMAGES

VIDEOS

关闭取词

22,900 Results

Any time ▾

Preoperative Chemoradiotherapy Using Concurrent ...

<https://ascopubs.org/doi/10.1200/JCO.2010.29.7697>

Jan 24, 2011 · To assess long-term clinical outcomes of preoperative chemoradiotherapy of magnetic resonance imaging (MRI)-defined locally advanced rectal adenocarcinoma using concurrent irinotecan and capecitabine.

Cited by: 32

Author: Simon Gollins, Arthur Sun Myint, Brian H...

Publish Year: 2011

Phase I study of preoperative chemoradiotherapy with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6403498>

Feb 08, 2019 · The present study designed a novel preoperative chemoradiotherapy (CRT) with sequential oxaliplatin and irinotecan with S-1 for locally advanced rectal cancer (LARC). This phase I study evaluated the maximum tolerated dose and recommended dose (RD) of oxaliplatin following irinotecan with S-1.

Author: Hiroyuki Fujikawa, Yuji Toiyama, Yasu... Publish Year: 2019

Neoadjuvant Chemoradiation Therapy Using Concurrent S-1 ...

<https://www.sciencedirect.com/science/article/pii/S0360301614002958>

Jul 01, 2014 · NCRT with concurrent S-1/irinotecan produced high response rates and excellent long-term survival, with acceptable adverse effects in patients with rectal cancer. ypN2 is a strong predictor of dismal outcomes, and a combination of ypT and tumor location can identify high-risk patients among those with ypN0/1 disease.

Cited by: 23

Author: Takatoshi Nakamura, Keishi Yamashita, T...

Publish Year: 2014

UGT1A1 polymorphisms in rectal cancer associated with the ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6272094>

Oct 22, 2018 · Our results indicate that UGT 1A1 polymorphism is a predictive factor to determine the clinical efficacy of preoperative chemoradiotherapy and hematological toxicity induced by chemoradiotherapy using irinotecan in locally advanced rectal cancer patients.



ALL

IMAGES

VIDEOS

关闭取词

23,000 Results

Any time ▾

Preoperative Chemoradiotherapy Using Concurrent ...

<https://ascopubs.org/doi/10.1200/JCO.2010.29.7697>

Jan 24, 2011 · To assess long-term clinical outcomes of **preoperative chemoradiotherapy of magnetic resonance imaging (MRI)-defined locally advanced rectal adenocarcinoma using concurrent irinotecan and capecitabine.**

Cited by: 32

Author: Simon Gollins, Arthur Sun Myint, Brian H...

Publish Year: 2011

Neoadjuvant Chemoradiation Therapy Using Concurrent S-1 ...

<https://www.sciencedirect.com/science/article/pii/S0360301614002958>

Jul 01, 2014 · **NCRT with concurrent S-1/irinotecan produced high response rates and excellent long-term survival, with acceptable adverse effects in patients with rectal cancer.** ypN2 is a strong predictor of dismal outcomes, and a combination of ypT and tumor location can identify high-risk patients among those with ypN0/1 disease.

Cited by: 23

Author: Takatoshi Nakamura, Keishi Yamashita, T...

Publish Year: 2014

Phase I study of preoperative chemoradiotherapy with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6403498>

The present study designed a novel **preoperative chemoradiotherapy (CRT) with sequential oxaliplatin and irinotecan with S-1** for locally advanced rectal cancer (LARC). This phase I study evaluated the maximum tolerated dose and recommended dose (RD) of oxaliplatin following irinotecan with S-1.

Author: Hiroyuki Fujikawa, Yuji Toiyama, Yasu... Publish Year: 2019

Clinicopathological outcomes of preoperative ...

<https://link.springer.com/article/10.1007/s00595-015-1250-z>

Sep 12, 2015 · To investigate the clinicopathological outcomes of patients with T4 lower rectal cancer treated using **preoperative chemoradiotherapy with S-1 plus Irinotecan.** Methods Between 2005 and 2011, 35 patients with T4M0 lower rectal cancer, diagnosed initially as T4a in 12 and as T4b in 23, were treated with 45 Gy of radiotherapy concomitantly with S-1 ...

Cited by: 3

Author: Naohito Beppu, Hidenori Yoshie, Fumihik...

Publish Year: 2016

(PDF) Preoperative Chemoradiotherapy Using Concurrent