

Match Overview

1	Internet 55 words crawled on 09-Jul-2017 www.cancerindex.org	2%
2	Internet 34 words crawled on 06-Oct-2019 f6publishing.blob.core.windows.net	1%
3	Crossref 16 words Helga Gimbel. "Total or subtotal hysterectomy for benign ... erine diseases? A meta-analysis", Acta Obstetrica Et Gyne	1%
4	Internet 13 words crawled on 06-Sep-2016 www.diagnosticimaging.com	<1%
5	Crossref 13 words &NA; "Abstracts", Journal of Pelvic Medicine and Surgery, 0 3/2007	<1%
6	Internet 12 words crawled on 09-Apr-2019 link.springer.com	<1%

2
Name of journal: *World Journal of Clinical Cases*

Manuscript NO: 52002

Manuscript Type: CASE REPORT

1
Large pelvic mass arising from the cervical stump: A case report

Kai Zhang, Jing-Hong Jiang, Jia-Li Hu, Yu-Lin Liu, Xu-Hong Zhang, Ying-Mei
Wang, Feng-Xia Xue

Abstract

BACKGROUND

A large cervical cyst with a cervical high squamous intraepithelial lesion arising from the cervical stump is rare. After supracervical hysterectomy, there is a risk of various lesions occurring in the cervical stump. We review the types and



All

Images

Videos

关闭取词

10,300 Results

Any time ▾

Gastrointestinal stromal tumor mimicking gynecological ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3861168>

Case report. A 53-year-old woman presented with **upper abdominal pain**. In her medical history, she had subtotal hysterectomy and bilateral salpingo-oophorectomy operation due to **myoma uteri**, and as a prophylactic oophorectomy. During **abdominal ultrasonography** for ruling out cholelithiasis, a pelvic mass was found.

Author: Eyup Yayci, Omer Tolga Guler, Gamz... Publish Year: 2012

Serous carcinoma arising from uterine adenomyosis ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4893213>

Jun 04, 2016 · Case 3. The **cervical stump** measured 5 × 5 × 4.5 cm. The **mucosa** looked a little bit rough, but no tumor was grossly seen in the **cervical canal**. A 3 × 1.5 × 1.5-cm **mass** was recognizable in the **myometrium**. The **mass** had 3 **small cysts** containing brownish, viscous fluid. The diameters of these **cysts** varied between 0.5-1.0 cm.

Cited by: 3 Author: Bingjian Lu, Qin Chen, Xiaofei Zhang, Lili ...

Publish Year: 2016

Novel management of a giant cervical myoma in a ...

<https://casereports.bmj.com/content/2017/bcr-2017-221408> ▾

Oct 07, 2017 · Mean size in reported series was 13 cm (range 5–24 cm). 7–15 Differentials for a large pelvic mass originating from the uterus in a premenopausal female patient will include: **Pregnancy Uterine corpus lesions**: benign (**leiomyomas**, **adenomyosis** and **polyps**) or malignant (**leiomyosarcoma**, **endometrial stromal sarcomas**, **carcinosarcoma** and **endometrial carcinoma**)

Author: Jolene Wong, Grace Hwei Ching Tan, ... Publish Year: 2017

Myoma arising in a Cervical Stump | OBGYN.Net

<https://www.obgyn.net/laparoscopy/myoma-arising-cervical-stump> ▾

Jun 29, 2011 · A **myoma** arising in the **cervical stump** measuring 8 cm in width was diagnosed both by the **pelvic examination** and vaginal ultrasound. The **myoma** appeared to be fixed in the **pelvic cavity**, with absence of **lateral movements**.

[PDF] Cervical Carcinosarcoma Occurring after Subtotal ...

[https://www.gynecologiconcology-online.net/article/S0090-8258\(97\)94872-1/pdf](https://www.gynecologiconcology-online.net/article/S0090-8258(97)94872-1/pdf)

present a case of **carcinosarcoma** arising in the **cervical remnant** of a patient 30 years following a



All

Images

Videos

开启取词

10,400 Results

Any time ▾

Gastrointestinal stromal tumor mimicking gynecological ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3861168>

During abdominal ultrasonography for ruling out cholelithiasis, a pelvic mass was found. On vaginal examination, cervical stump was seen as normal and with bimanual examination an ovoid shaped mass was found. It was 8 × 5 cm in size as a mobile mass.

Author: Eyup Yayci, Omer Tolga Guler, Gamz... Publish Year: 2012

Novel management of a giant cervical myoma in a ...

<https://casereports.bmj.com/content/2017/bcr-2017-221408> ▾

Oct 07, 2017 · Mean size in reported series was 13 cm (range 5–24 cm). 7–15 Differentials for a large pelvic mass originating from the uterus in a premenopausal female patient will include: **Pregnancy** **Uterine corpus lesions:** benign (leiomyomas, adenomyosis and polyps) or malignant (leiomyosarcoma, endometrial stromal sarcomas, carcinosarcoma and endometrial carcinoma)

Author: Jolene Wong, Grace Hwei Ching Tan, ... Publish Year: 2017

Serous carcinoma arising from uterine adenomyosis ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4893213>

Jun 04, 2016 · Scanty serous carcinoma was present in the endometrium without evidence of myometrial invasion. The eutopic endometrium in both case 1 and 2 had no evidence of neoplastic changes after complete examination. Case 3 had 3 microscopic serous EICs in the adenomyotic cysts of the cervical stump.

Cited by: 3 Author: Bingjian Lu, Qin Chen, Xiaofei Zhang, Lili ...

Publish Year: 2016

Leiomyoma Recurrent at the Cervical Stump: Report of Two ...

<https://www.sciencedirect.com/science/article/pii/S1553465011012830>

Case Reports. In light of the concern for malignancy, the patient was counseled about exploratory laparotomy and resection of the mass. At the time of her procedure a 15-cm solid mass was visualized arising from the cervical stump. The mass was mobilized from the anterior and lateral pelvic walls and resected along with the cervix.

Cited by: 3 Author: Christine M. Chu, Uchenna C. Acholonu, ...

Publish Year: 2012

[PDF] Case Cervical Stump Fibroidpost Supra-cervical ...

journal.fk.unpad.ac.id/index.php/ijhs/article/download/1257/1210