



26,400 Results

Any time ▾

## Metastatic renal cell carcinoma without evidence of a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4059817>

Although metastases are common in patients with renal cell carcinoma (rcc), it is extremely rare for patients to present with metastatic rcc (m rcc) without evidence of a primary mass in the kidney. Two cases of m rcc with no detectable primary renal mass are reported here. Both patients had bilateral native kidneys in situ and no significant prior urologic history.

Cited by: 6

Author: R.M. Kumar, T. Aziz, H. Jamshaid, J. Gill...

Publish Year: 2014

## Renal Cell Carcinoma With Retroperitoneal Lymph Nodes ...

<https://www.sciencedirect.com/science/article/pii/S0022534705635379>

A, 32-year-old male who presented with hematuria and nonspecific abdominal pain had small, central right renal cell carcinoma, bulky retroperitoneal adenopathy and no evidence of distant metastatic disease. B, microscopic foci (arrows) of clear cell carcinoma within lymph node surrounding renal pedicle. Reduced from  $\times 40$ .

Cited by: 366

Author: Allan J. Pantuck, Amnon Zisman, Fredric...

Publish Year: 2003

## Renal cell carcinoma with retroperitoneal lymph nodes ...

<https://onlinelibrary.wiley.com/doi/full/10.1002/cncr.11422>

The disease-specific survival for patients with metastatic renal cell carcinoma with regional lymphadenopathy was equivalent regardless of the extent of lymph node disease. N1 indicates metastasis in a single lymph node and N2 indicates metastases in more than one lymph node.

Cited by: 233

Author: Allan J. Pantuck, Amnon Zisman, Frederi...

Publish Year: 2003

## Renal cell carcinoma with retroperitoneal lymph nodes ...

[onlinelibrary.wiley.com/doi/10.1002/cncr.11422/full](https://onlinelibrary.wiley.com/doi/10.1002/cncr.11422/full)

May 29, 2003 - The disease-specific survival for patients with metastatic renal cell carcinoma with regional lymphadenopathy was equivalent regardless of the extent of lymph node disease. N1 indicates metastasis in a single lymph node and N2 indicates metastases in more than one lymph node.

**Name of Journal:** *World Journal of Clinical Oncology*

**Manuscript NO:** 52054

**Manuscript Type:** CASE REPORT

**Metastatic clear cell renal cell carcinoma in isolated retroperitoneal lymph node without evidence of primary tumor in kidneys: A case report**

Shields LB *et al.* CCRCC Lymph Node without primary

Lisa BE Shields, Arash Rezazadeh Kalebasty

**Abstract**

**BACKGROUND**

Retroperitoneal lymph node dissection (RPLND) plays a diagnostic, therapeutic, and

### Match Overview

1	Internet 39 words crawled on 25-Nov-2019 <a href="http://pubs.rsna.org">pubs.rsna.org</a>	2%
2	Internet 35 words crawled on 10-Sep-2018 <a href="http://www.spandidos-publications.com">www.spandidos-publications.com</a>	2%
3	Crossref 33 words Hasan Khosravi, Andressa L. Akabane, Allireza Alloo, Rosa Lynn M. Nazarian, Genevieve M. Boland. "Metastatic melar..."	2%
4	Crossref 29 words Manoj P. Menon, Keith D. Eaton. "Spontaneous Regressi... of Non-Small-Cell Lung Cancer in AIDS After Immune Rec..."	1%
5	Crossref 13 words Karmali, Riaz J., Hiroo Suami, Christopher G. Wood, and J ose A. Karam. "Lymphatic drainage in renal cell carcinom..."	1%



All

Images

Videos

关闭取词

26,600 Results

Any time ▾

## Renal Cell Carcinoma With Retroperitoneal Lymph Nodes ...

<https://www.sciencedirect.com/science/article/pii/S0022534705635379>

A, 32-year-old male who presented with hematuria and nonspecific abdominal pain had small, central right **renal cell carcinoma**, bulky **retroperitoneal** adenopathy and no **evidence** of distant **metastatic** disease. B, microscopic foci (arrows) of **clear cell carcinoma** within **lymph node** surrounding **renal** pedicle. Reduced from ×40.

**Cited by:** 365

**Author:** Allan J. Pantuck, Amnon Zisman, Fredrick...

**Publish Year:** 2003

## Metastatic renal cell carcinoma without evidence of a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4059817>

Although metastases are common in patients with **renal cell carcinoma** (rcc), it is extremely rare for patients to present with **metastatic** rcc (m rcc) **without evidence of a primary** mass in the kidney. Two cases of m rcc with no detectable **primary renal** mass are reported here. Both patients had bilateral native **kidneys** in situ and no significant prior urologic history.

**Cited by:** 6

**Author:** R.M. Kumar, T. Aziz, H. Jamshaid, J. Gill, ...

**Publish Year:** 2014

## Renal cell carcinoma with retroperitoneal lymph nodes ...

<https://onlinelibrary.wiley.com/doi/full/10.1002/cncr.11422>

The disease-specific survival for patients with **metastatic renal cell carcinoma** with regional lymphadenopathy was equivalent regardless of the extent of **lymph node** disease. N1 indicates metastasis in a single **lymph node** and N2 indicates metastases in more than one **lymph node**.

**Cited by:** 232

**Author:** Allan J. Pantuck, Amnon Zisman, Frederic...

**Publish Year:** 2003

## Renal cell carcinoma with retroperitoneal lymph nodes ...

[onlinelibrary.wiley.com/doi/10.1002/cncr.11422/full](https://onlinelibrary.wiley.com/doi/10.1002/cncr.11422/full)

May 29, 2003 · The disease-specific survival for patients with **metastatic renal cell carcinoma** with