

Supplementary table 1. Adjusted hazard ratios of middle-age and elderly Crohn's disease for other variables not depicted in the main figure

			Adjusted Hazard Ratio (95% Confidence Interval)	
			Middle-age	Elderly
Lowest income quintile	No		1(reference)	1(reference)
	Yes		0.95(0.82,1.10)	1.14(0.91,1.41)
Diabetes mellitus	No		1(reference)	1(reference)
	Yes		0.92 (0.74,1.15)	1.03(0.81,1.31)
Hypertension	No		1(reference)	1(reference)
	Yes		0.94(0.81,1.09)	1.14(0.93,1.39)
Metabolic syndrome	No		1(reference)	1(reference)
	Yes		0.87(0.76,1.01)	1.09(0.894,1.335)
Fasting glucose*	Q1		1(reference)	1(reference)
	Q2		0.90(0.76,1.06)	0.83(0.63,1.09)
	Q3		0.86(0.72,1.03)	0.87(0.67,1.14)
	Q4		0.76(0.62,0.93)	0.73(0.52,1.02)
Total cholesterol*	Q1		1(reference)	1(reference)
	Q2		0.98(0.82,1.18)	1.23(0.93,1.62)
	Q3		0.95(0.79,1.15)	0.91(0.70,1.20)
	Q4		0.81(0.65,1.01)	0.99(0.75,1.32)

* Fasting glucose (mg/dL) and total cholesterol (mg/dL) levels were sub-divided into quartiles.