

World Journal of *Gastroenterology*

World J Gastroenterol 2020 March 14; 26(10): 995-1106



OPINION REVIEW

- 995 Global whole family based-*Helicobacter pylori* eradication strategy to prevent its related diseases and gastric cancer
Ding SZ

REVIEW

- 1005 Role of spleen tyrosine kinase in liver diseases
Kurniawan DW, Storm G, Prakash J, Bansal R

MINIREVIEWS

- 1020 Abnormal liver function tests associated with severe rhabdomyolysis
Lim AKH

ORIGINAL ARTICLE**Basic Study**

- 1029 Mesencephalic astrocyte-derived neurotrophic factor ameliorates steatosis in HepG2 cells by regulating hepatic lipid metabolism
He M, Wang C, Long XH, Peng JJ, Liu DF, Yang GY, Jensen MD, Zhang LL

Retrospective Cohort Study

- 1042 Prognostic factors and predictors of postoperative adjuvant transcatheter arterial chemoembolization benefit in patients with resected hepatocellular carcinoma
Chen MY, Juengpanich S, Hu JH, Topatana W, Cao JS, Tong CH, Lin J, Cai XJ

Retrospective Study

- 1056 Double-balloon endoscopic retrograde cholangiopancreatography for patients who underwent liver operation: A retrospective study
Nishio R, Kawashima H, Nakamura M, Ohno E, Ishikawa T, Yamamura T, Maeda K, Sawada T, Tanaka H, Sakai D, Miyahara R, Ishigami M, Hirooka Y, Fujishiro M
- 1067 Serum N-glycan markers for diagnosing liver fibrosis induced by hepatitis B virus
Cao X, Shang QH, Chi XL, Zhang W, Xiao HM, Sun MM, Chen G, An Y, Lv CL, Wang L, Nan YM, Chen CY, Tan ZN, Liu XE, Zhuang H
- 1080 Predictors of outcomes of endoscopic balloon dilatation in strictures after esophageal atresia repair: A retrospective study
Dai DL, Zhang CX, Zou YG, Yang QH, Zou Y, Wen FQ

Prospective Study

- 1088** Technetium-99m-labeled macroaggregated albumin lung perfusion scan for diagnosis of hepatopulmonary syndrome: A prospective study comparing brain uptake and whole-body uptake
Zhao H, Tsao J, Zhang XW, Ma HY, Weng NN, Tang GS, Li X

META-ANALYSIS

- 1098** Is aggressive intravenous fluid resuscitation beneficial in acute pancreatitis? A meta-analysis of randomized control trials and cohort studies
Gad MM, Simons-Linares CR

ABOUT COVER

Associate Editor of *World Journal of Gastroenterology*, Nahum Mendez-Sanchez, FACG, MD, PhD, Doctor, Professor, Liver Research Unit, Medica Sur Clinic & Foundation, Mexico City 14050, Mexico

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (*WJG*, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2019 edition of Journal Citation Report® cites the 2018 impact factor for *WJG* as 3.411 (5-year impact factor: 3.579), ranking *WJG* as 35th among 84 journals in gastroenterology and hepatology (quartile in category Q2). CiteScore (2018): 3.43.

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Yan-Liang Zhang*
 Proofing Production Department Director: *Yun-Xiaojuan Wu*

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Subrata Ghosh, Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE

Ze-Mao Gong, Director

PUBLICATION DATE

March 14, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Basic Study

Mesencephalic astrocyte-derived neurotrophic factor ameliorates steatosis in HepG2 cells by regulating hepatic lipid metabolism

Miao He, Cong Wang, Xiao-Hong Long, Jia-Jia Peng, Dong-Fang Liu, Gang-Yi Yang, Michael D Jensen, Li-Li Zhang

ORCID number: Miao He (0000-0002-9055-7042); Cong Wang (0000-0002-1209-4978); Xiao-Hong Long (0000-0002-2412-0053); Jia-Jia Peng (0000-0001-9423-6847); Dong-Fang Liu (0000-0002-8801-2190); Gang-Yi Yang (0000-0002-6458-6747); Michael D Jensen (0000-0001-5589-8389); Li-Li Zhang (0000-0001-9007-5281).

Author contributions: He M and Wang C performed the majority of the experiments and contributed to the analysis of the data and the drafting of the manuscript; Long XH and Peng JJ contributed to the analysis of the data; Liu DF and Yang GY revised and approved the final version of the manuscript; Zhang LL contributed to the study design, the drafting of the manuscript, and critical discussion and approved the final version of the manuscript; Jensen MD revised, discussed, and approved the final version of the manuscript.

Supported by National Natural Science Foundation of China, No. 81300702 and No. 81501199; Natural Science Foundation Project of Chongqing CSTC, No. cstc2018jcyjAX0210 and No. cstc2017jcyjAX0016; and the Kuanren Talents Program of the Second Affiliated Hospital of Chongqing Medical University.

Institutional animal care and use committee statement: This study protocol was reviewed and approved by the Animal Care and Use Ethics Committee of Chongqing Medical University, Chongqing, China (protocol No.

Miao He, Cong Wang, Xiao-Hong Long, Jia-Jia Peng, Dong-Fang Liu, Gang-Yi Yang, Li-Li Zhang, Department of Endocrinology, the Second Affiliated Hospital, Chongqing Medical University, Chongqing 400010, China

Michael D Jensen, Endocrine Research Unit, Mayo Clinic, Rochester, MN 55905, United States

Corresponding author: Li-Li Zhang, MD, PhD, Associate Professor, Doctor, Postdoc, Research Fellow, Department of Endocrinology, the Second Affiliated Hospital, Chongqing Medical University, 74 Linjiang Road, Yuzhong District, Chongqing 400010, China. zhanglili.jl@fox-mail.com

Abstract**BACKGROUND**

Nonalcoholic fatty liver disease (NAFLD) is a global metabolism-associated liver disease. Mesencephalic astrocyte-derived neurotrophic factor (MANF) is a newly discovered secreted protein that is involved in metabolic homeostasis. However, much remains to be discovered about its function in hepatic lipid metabolism; thus, we assessed whether MANF could regulate hepatic metabolism.

AIM

To establish *in vivo* and *in vitro* NAFLD models to explore the role of MANF in hepatic lipid metabolism.

METHODS

HepG2 cells treated with free fatty acids (FFAs) and ob/ob mice were used as NAFLD models. Liver tissues collected from wild type and ob/ob mice were used to detect MANF expression. Cells were treated with FFAs for different durations. Moreover, we used lentiviral constructs to establish overexpression and knockdown cell models in order to interfere with MANF expression levels and observe whether MANF influences hepatic steatosis. Western blot analysis and quantitative real-time PCR were used to detect protein and gene expression, and oil red O staining was used to visualize intracellular lipid droplets.

RESULTS

Hepatic MANF protein and mRNA expression in wild type mice were 10-fold and 2-fold higher, respectively, than those in ob/ob mice. The MANF protein was temporarily increased by 1.3-fold after stimulation with FFAs for 24 h and gradually decreased to 0.66-fold that of the control at the 72 h time point in HepG2 cells. MANF deficiency upregulated the expression of genes involved in

2019-48). All animal experiments conformed to the accepted principles for the care and use of laboratory animals (license No. SYXK 2018-0003).

Conflict-of-interest statement: All the authors have no conflicts of interest related to the manuscript.

Data sharing statement: No additional data are available.

ARRIVE guidelines statement: The authors have read the ARRIVE guidelines, and the manuscript was prepared and revised according to the ARRIVE guidelines.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: November 21, 2019

Peer-review started: November 21, 2019

First decision: January 7, 2020

Revised: February 15, 2020

Accepted: February 21, 2020

Article in press: February 21, 2020

Published online: March 14, 2020

P-Reviewer: Fatkhudinov T, Aureliano M, Hosomi R

S-Editor: Zhou JJ

L-Editor: Wang TQ

E-Editor: Zhang YL



fatty acid synthesis, cholesterol synthesis, and fatty acid uptake and aggravated HepG2 cell steatosis, while MANF overexpression inhibited fatty acid synthesis and uptake and cholesterol synthesis, and rescued HepG2 cells from FFAs-induced steatosis. Furthermore, a significant decrease in triglyceride levels was observed in the MANF overexpression group compared with the control group (0.4288 ± 0.0081 mmol/g vs 0.3746 ± 0.0121 mmol/g, $P < 0.05$) upon FFAs treatment. There was also a 17% decrease in intracellular total cholesterol levels between the MANF overexpression group and the control group (0.1301 ± 0.0059 mmol/g vs 0.1088 ± 0.0009 mmol/g, $P < 0.05$) upon FFAs treatment. Moreover, MANF suppressed lipid deposition in HepG2 cells.

CONCLUSION

Our findings indicate that MANF improves the phenotype of liver cell steatosis and may be a potential therapeutic target in hepatic steatosis processes.

Key words: Mesencephalic astrocyte-derived neurotrophic factor; Nonalcoholic fatty liver disease; Hepatic steatosis; Lipogenesis; *In vitro*; HepG2

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: We first uncovered an important function of mesencephalic astrocyte-derived neurotrophic factor (MANF) in the pathogenesis of nonalcoholic fatty liver disease. We found that MANF exerts a significant effect on hepatic fatty metabolism. This study suggests for the first time that MANF expression was increased at an early stage and gradually decreased afterward under high free fatty acids stimulation in HepG2 cells. Moreover, the results from gain-and-loss functional experiments showed that loss of MANF accelerated lipogenesis and aggravated HepG2 cell steatosis, while MANF overexpression inhibited lipogenesis and rescued HepG2 cell steatosis from free fatty acids treatment.

Citation: He M, Wang C, Long XH, Peng JJ, Liu DF, Yang GY, Jensen MD, Zhang LL. Mesencephalic astrocyte-derived neurotrophic factor ameliorates steatosis in HepG2 cells by regulating hepatic lipid metabolism. *World J Gastroenterol* 2020; 26(10): 1029-1041

URL: <https://www.wjnet.com/1007-9327/full/v26/i10/1029.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v26.i10.1029>

INTRODUCTION

Nonalcoholic fatty liver disease (NAFLD) is a clinicopathological condition characterized by the intracellular accumulation of lipids in hepatocytes followed by steatosis and inflammation in the liver. This multifactorial disease affects approximately 25% of the average population and 70%-90% of overweight/obese patients^[1]. The pathogenic mechanisms of NAFLD are known to include lipid accumulation, inflammation, and oxidative stress. However, the precise mechanisms behind the pathogenesis of NAFLD are largely unknown. Given the high prevalence and severe prognosis of NAFLD, understanding the underlying mechanisms has become a priority.

Mesencephalic astrocyte-derived neurotrophic factor (MANF) is a newly discovered secreted protein that was originally isolated from a rat mesencephalic type-1 astrocyte cell line^[2]. The structure of MANF includes two important domains. The N-terminal domain is homologous with saposin-like proteins, which are a large family of small, cysteine-rich proteins that interact with lipids and membranes^[3]. The C-terminal domain is homologous with SAF-A/B, Acinus, and PIAS proteins and may have reductase or disulfide isomerase activity^[4]. MANF is widely expressed in neuronal and nonneuronal tissues, particularly secretory tissues such as the liver and pancreas^[5]. Studies have shown that MANF could exert protective functions in multiple diseases, such as intracerebral hemorrhage and Parkinson's disease^[6,7]. Both *in vivo* and *in vitro* studies indicated that the cytoprotective effect of MANF is not limited to the nervous system, and it can also protect pancreatic beta cells from inflammatory stress-induced cell death^[8] and improve pancreatic β cell proliferation and survival, while MANF knockout mice display a severe diabetic phenotype^[9].

Furthermore, human studies have shown that circulating MANF levels are increased in patients with type 1 diabetes, newly diagnosed prediabetes, and type 2 diabetes, suggesting a potential role for MANF in metabolic diseases^[9-12].

More interestingly, a recent study showed that MANF plays an increasingly significant role in obesity and participates in regulating food intake and body weight^[13], protecting the liver against aging and metabolic stress-related damage, and that nonalcoholic steatohepatitis patients showed decreased levels of circulating MANF^[14]. However, much remains to be explored about the role of MANF in hepatic lipid homeostasis. Regarding the potential role of MANF in metabolic diseases and its protective effect against metabolic stress, we proposed that MANF may play a potential protective role in NAFLD and have beneficial effects on lipid stress-induced liver cell steatosis.

MATERIALS AND METHODS

Animal studies

All animal experiments were approved by the Animal Care and Use Ethics Committee of Chongqing Medical University (approved protocol No. 2019-48). The animal protocol was designed to minimize pain or discomfort to the animals. Adult male mice [C57BL/6J, 8-week-old, 18 ± 3 g, wild-type (WT) mice, $n = 4$] and 8-week-old male ob/ob mice ($n = 4$) were maintained in an environment with a 12 h light/dark cycle at 21-25 °C with free access to water and food. Genetically obese mice (ob/ob, Model Animal Research Center of Nanjing University, Nanjing, Jiangsu, China) fed a normal control chow diet were used for the fatty liver model, and WT mice (Chongqing Medical University, Chongqing, China) fed a normal control chow diet served as controls. After 12 wk of feeding, mice were euthanized with 10 g/L pentobarbital sodium, and liver tissues were collected and immediately stored at -80 °C or fixed in 40 g/L paraformaldehyde for further analyses.

Cell culture and treatment

Human HepG2 cells were obtained from Procell Life Science and Technology (Wuhan, Hubei, China). The cells were incubated in Dulbecco's Modified Eagle's Medium (HyClone, Logan, United States) with 100 g/L fetal bovine serum (ExCell Bio, Shanghai, China) and 10 g/L penicillin-streptomycin solution (Beyotime, Shanghai, China). All cells were placed in a clean humidified incubator containing 50 mL/L CO₂ at 37 °C. For the MANF knockdown experiment, HepG2 cells were transfected with lentivirus (Lv) expressing short hairpin RNA-targeted MANF, and the lentiviral vector alone was used as the control. For the MANF overexpression experiment, HepG2 cells were transfected with lentivirus expressing MANF, and lentiviral vector alone was used as a control. All lentiviruses were designed, synthesized, and packed by HanBio (Hanbio Biotechnology Co. Ltd., Shanghai, China). To construct a nonalcoholic fatty liver model *in vitro*, HepG2 cells were cultured in medium with 0.5 mmol/L FFAs (oleate:palmitate = 2:1, Sigma-Aldrich, St. Louis, MO, United States) containing 10 g/L low fatty acid bovine serum albumin (BSA, Sigma) as previously reported^[15], and cells treated with BSA served as controls.

Western blot analysis

Total protein was extracted from mouse liver tissues or HepG2 cells with RIPA lysis buffer (Beyotime) containing phosphatase inhibitor cocktail A (Beyotime) and protease inhibitor (Roche, Basel, Switzerland), and the protein concentration was measured with a BCA Protein Assay kit (Beyotime). Protein samples (30 µg) were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (120 mL/L gel) and transferred to a 0.22 µm polyvinylidene difluoride membrane (Merck Millipore, Darmstadt, Germany). Then, the membrane was blocked with 50 g/L BSA for 2 h at room temperature, followed by incubation with primary antibody at 4 °C for 12-16 h and then secondary antibody for 1-2 h at room temperature. The protein expression level was detected using the chemiluminescence method, and the images were captured by a computer. GAPDH served as a control. The antibodies included anti-MANF (Abcam, Cambridge, MA, United States, ab67271, 1:1000), anti-GAPDH (Proteintech Group, Wuhan, Hubei, China, 60004-1-Ig, 1:5000) and HRP-coupled goat anti-rabbit or goat anti-mouse IgG secondary antibody (Zhongshan Co., Beijing, China, ZB-2301, ZB-2305, 1:5000).

Quantitative RT-PCR

Total RNA was extracted from mouse liver tissues and cells with TRIzol Reagent (Ambion, Carlsbad, CA, United States) according to the manufacturer's instructions. RNA concentration and quality were measured with a Nanodrop2000 (Thermo

Scientific). RNA was then reverse transcribed into cDNA using a PrimeScriptRT reagent kit (Takara Bio, Japan) according to the manufacturer's protocol. Quantitative PCR was performed using a TB Green™ Premix Ex Taq™ II kit (Takara) and an ABI StepOnePlus Real-Time PCR System (Life Technologies, CA, United States). The expression levels were normalized to the level of β -actin mRNA in the same samples. The data were analyzed by the $2^{-\Delta\Delta Ct}$ method. The primers used are listed in Table 1.

Oil red O staining

At the end of treatment, HepG2 cells on 6-well plates were washed with phosphate-buffered saline (PBS, HyClone), fixed with 40 g/L paraformaldehyde fixing solution (Boster Biological Technology Co., Ltd, Wuhan, Hubei, China) for 30 min at room temperature, washed with PBS, and dipped in 600 mL/L isopropanol for 1 min. Subsequently, the cells were stained with 5 g/L oil red O solution (Sigma) in 600 mL/L isopropanol for 15 min and then washed with distilled water three times. Finally, cells were counterstained with hematoxylin staining solution (Zhongshan Co.) for 5 s, washed with distilled water, and mounted with glycerol jelly mounting medium (Beyotime). The cells were then visualized under a light microscope (Olympus, Tokyo, Japan), and images were captured. Staining was quantified using ImageJ (National Institute of Health, Bethesda, MD, United States) and analyzed as previously described^[16,17].

Measurements of intracellular lipids

Intracellular triglyceride (TG) and total cholesterol (TC) contents were quantified with a triglyceride assay kit and a total cholesterol assay kit (Nanjing Jiancheng Bioengineering Institute, Nanjing, Jiangsu, China) according to the manufacturer's instructions.

Statistical analysis

Statistical analyses were performed using SPSS statistical software (Version 21.0, SPSS Inc., Chicago, IL, United States). Data distributions were assessed using the Shapiro-Wilk test. Differences in normally distributed data were analyzed by independent-samples *t*-tests (two groups) or one-way ANOVA (three or more groups) followed by Tukey's test for data that were homoscedastic or Tamhane's T2 analysis for data that were heteroscedastic. For all statistical tests, $P < 0.05$ was considered indicative of statistical significance. All the data are presented as the mean \pm SE, and the graphics were plotted by GraphPad Prism 5.0 (GraphPad Software, La Jolla, CA, United States).

RESULTS

Hepatic MANF expression is decreased in NAFLD

To explore the expression of MANF in fatty liver, we detected its expression in liver samples from ob/ob mice and WT mice (animals analyzed: 4/4). Hepatic MANF protein and mRNA expression in WT mice were 10-fold ($P < 0.01$) and 2-fold ($P < 0.05$) higher, respectively, than those in ob/ob mice (Figure 1A and B). Interestingly, the MANF protein expression was firstly raised to 1.3 times ($P < 0.01$) upon FFAs treatment for 24 h and subsequently reduced by 47% ($P < 0.05$) at the 72 h (Figure 1C). The MANF mRNA level was temporarily increased by 2 times ($P < 0.001$) with FFAs stimulation for 24 h and gradually reduced by 25% ($P < 0.01$) at the 72 h time point in HepG2 cells (Figure 1D). Furthermore, to determine the expression of MANF under lipid stress *in vitro*, HepG2 cells were treated with BSA or 0.5 mmol/L FFAs, and oil red O staining showed that intracellular lipid droplets were significantly increased in HepG2 cells under FFAs stimulation (Figure 1E).

MANF overexpression alleviates lipid deposition in HepG2 cells

To further explore the function of MANF in lipid-induced stress, we used lentiviral vectors to overexpress MANF and screened the stably transfected cell line (Figure 2A). The mRNA and protein expression were assessed by quantitative real-time PCR and Western blot, respectively (Figure 2B). As shown in Figure 2C, without FFAs treatment, MANF overexpression did not impact the lipid droplet area, while upon FFAs treatment, MANF overexpression significantly diminished the lipid droplet content in HepG2 cells ($P < 0.01$).

MANF overexpression suppresses lipogenesis

Next, we determined the intracellular content of TG and TC in HepG2 cells. The results showed a significant decrease in TG levels (0.3746 ± 0.0121 mmol/g) in the MANF over-expression group compared with those in the control group ($0.4288 \pm$

Table 1 Primers used for quantitative real-time PCR experiments

Gene	Species	Primers
MANF	<i>Homo sapiens</i>	F: 5'-CGTTGTGCTACTATATCGGGG-3' R: 5'-GGCCAGAGGCTTTGATACCT-3'
ACCa	<i>Homo sapiens</i>	F: 5'-ATGTCCTGGCTTGACCTAGTA-3' R: 5'-CCCCAAAGCGAGTAACAAATTCT-3'
FAS	<i>Homo sapiens</i>	F: 5'-AGATTGTGTGATGAAGGACATGG-3' R: 5'-TGTTGCTGGTGAGTGTGCATT-3'
SREBP-1C	<i>Homo sapiens</i>	F: 5'-CGGAACCATCTTGGCAACAGT-3' R: 5'-CGCTTCTCAATGGCGTTGT-3'
HMGCR	<i>Homo sapiens</i>	F: 5'-TGATTGACCTTCCAGAGCAAG-3' R: 5'-CTAAAATTGCCATTCCACGAGC-3'
CD36	<i>Homo sapiens</i>	F: 5'-GGCTGTGACCGAACGTG-3' R: 5'-AGGTCCTCAACTGGCATTAGAA-3'
FABP-1	<i>Homo sapiens</i>	F: 5'-GTGTCGGAAATCGTGCAGAAT-3' R: 5'-GACTTCTCCCTGTCTATTGTC-3'
β -actin	<i>Homo sapiens</i>	F: 5'-GCCGACAGGATGCAGAAGG-3' R: 5'-TGGAAGGTGGACAGCGAGG-3'
MANF	<i>Mus musculus</i>	F: 5'-TCTGGGACGATTTTACCAGGA-3' R: 5'-TCTTGCTTCACGGCAAAACITTA-3'
β -actin	<i>Mus musculus</i>	F: 5'-TCACTGTCCACCTCCAGCAGATG-3' R: 5'-CTCAGTAACAGTCCGCCTAGAAGC-3'

MANF: Mesencephalic astrocyte-derived neurotrophic factor; ACC α : Acetyl-CoA carboxylase α ; FAS: Fatty acid synthase; SREBP-1C: Sterol regulatory element binding transcription factor 1C; HMGCR: 3-hydroxy3-methylglutaryl coenzyme A reductase; FABP-1: Fatty acid binding protein-1.

0.0081 mmol/g) upon FFAs treatment ($P < 0.05$ Figure 3A). Furthermore, there was also a 17% decrease in intracellular TC levels (Figure 3B) in the MANF overexpression group (0.1088 ± 0.0009 mmol/g) compared with those in the control group (0.1301 ± 0.0059 mmol/g) after FFAs treatment ($P < 0.05$). Thus, to investigate the mechanisms of MANF responsible for suppressing lipogenesis, we detected the mRNA levels of lipid metabolism genes related to fatty acid synthesis (acetyl-CoA carboxylase α [ACC α], fatty acid synthase [FAS], sterol regulatory element binding transcription factor 1C [SREBP-1C]), cholesterol synthesis (3-hydroxy3-methylglutaryl [HMG] coenzyme A [CoA] reductase [HMGCR]), and fatty acid uptake (fatty acid translocase/CD36, fatty acid binding protein-1 [FABP-1]) in HepG2 cells.

The results showed that the expression of ACC α ($P < 0.05$) and SREBP-1C ($P < 0.01$) was lowered by 25% in the MANF overexpression group compared with that in the control group under FFAs treatment. Furthermore, there was a 57% reduction in FAS expression ($P < 0.001$) and a 18% decrease in HMGCR expression ($P < 0.05$) in the MANF overexpression group following FFAs stimulation (Figure 3C and D). After overexpression of MANF, the expression levels of CD36 and FABP-1 decreased by 71% and 33%, respectively (all $P < 0.001$, Figure 3E and F).

MANF deficiency exacerbates lipid accumulation in HepG2 cells

We then constructed a stable MANF-downregulated HepG2 cell line and explored the role of MANF deficiency in hepatic lipid accumulation (Figure 4A). The interference efficiency was also determined by quantitative real-time PCR and Western blot (Figure 4B). Oil red O staining showed no significant differences in lipid droplets between the MANF knockdown group and the control group in the absence of lipid stress. Additionally, under FFAs stimulation, the lipid droplet area in the MANF knockdown group was 37% greater than that in the control group ($P < 0.05$, Figure 4C).

MANF deficiency accelerates lipogenesis

Without FFAs treatment, MANF deficiency has little impact on intracellular TG levels. Interestingly, the intracellular TG levels were significantly increased in the MANF short hairpin RNA group (0.4294 ± 0.0072 mmol/g) compared with the control group (0.3946 ± 0.0041 mmol/g) under stimulation with FFAs ($P < 0.01$, Figure 5A). However, the intracellular TC levels were not significantly different between the




Figure 1 Mesencephalic astrocyte-derived neurotrophic factor expression is downregulated in both *in vitro* and *in vivo* nonalcoholic fatty liver disease models. A: Mesencephalic astrocyte-derived neurotrophic factor (MANF) protein expression in the liver of ob/ob mice and WT mice ($n = 4$), $^eP < 0.001$ vs WT; B: MANF mRNA expression in the liver of ob/ob mice and WT mice ($n = 3$), $^aP < 0.05$ vs WT; C and D: MANF protein (C) and mRNA (D) expression in HepG2 cells after treatment with bovine serum albumin or 0.5 mM free fatty acids at different time points, $^aP < 0.05$, $^bP < 0.01$, $^eP < 0.001$ vs 0 h; E: Representative oil red O staining of HepG2 cells (Scale bar = 50 μ m). Data are presented as the mean \pm SE. All cell experiments were repeated three times. MANF: Mesencephalic astrocyte-derived neurotrophic factor; GAPDH: Glyceraldehyde 3-phosphate dehydrogenase; BSA: Bovine serum albumin; FFAs: Free fatty acids; WT: Wild type.

MANF short hairpin RNA group and the control group with or without FFAs treatment (Figure 5B). In addition, the expression of genes related to fatty acid synthesis, cholesterol synthesis (Figure 5C and D), and fatty acid uptake (Figure 5E and F) was examined. The results showed that the expression of ACC α , FAS, SREBP-1C, and HMGCR in the MANF knockdown group was 2-fold ($P < 0.01$), 5.7-fold ($P < 0.001$), 4.8-fold ($P < 0.05$), and 1.6-fold ($P < 0.05$) higher, respectively, than that in the control group after FFAs treatment. The mRNA levels of CD36 and FABP-1 in the MANF knockdown group were 1.3-fold higher than those in the control group ($P < 0.01$).

DISCUSSION

Our study uncovered an important function of MANF in the pathogenesis of NAFLD. In the present study, both *in vivo* and *in vitro* studies provided basic pathological insights into a potential correlation between MANF and hepatic steatosis, which was functionally validated by up- and down-regulated MANF expression *in vitro*. Notably, MANF expression was increased in the early stage and gradually decreased under high FFAs stimulation. MANF deficiency increased lipogenesis and aggravated HepG2 cell steatosis, while MANF overexpression inhibited lipogenesis and rescued HepG2 cells from FFAs-induced steatosis. Our results demonstrated that MANF is a potential therapeutic target in hepatic steatosis processes.




Figure 3 Mesencephalic astrocyte-derived neurotrophic factor overexpression suppresses lipogenesis. A and B: The intracellular levels of TG (A) and TC (B) in HepG2 cells treated with bovine serum albumin or free fatty acids (FFAs) for 72 h, ^b*P* < 0.01, ^e*P* < 0.001 vs Lv-GFP; ^c*P* < 0.05 vs Lv-GFP + FFAs; C and D: The mRNA levels of genes involved in fatty acid synthesis (ACC α , FAS, SREBP-1C, and HMGCR) without FFAs treatment (C) and upon FFAs treatment (D), ^e*P* < 0.001 vs Lv-GFP; ^c*P* < 0.05, ^d*P* < 0.01, ^f*P* < 0.001 vs Lv-GFP + FFAs; E and F: The mRNA levels of genes related to lipid uptake (CD36, FABP-1) without FFAs treatment (E) and upon FFAs treatment (F), ^a*P* < 0.05, ^e*P* < 0.001 vs Lv-GFP; ^f*P* < 0.001 vs Lv-GFP + FFAs. Data represent the mean \pm SE, and all experiments were repeated three times. MANF: Mesencephalic astrocyte-derived neurotrophic factor; TG: Triglyceride; TC: Total cholesterol; Lv: Lentivirus; BSA: Bovine serum albumin; FFAs: Free fatty acids; ACC α : Acetyl-CoA carboxylase α ; FAS: Fatty acid synthase; SREBP-1C: Sterol regulatory element binding transcription factor 1C; HMGCR: 3-hydroxy3-methylglutaryl coenzyme A reductase; FABP-1: Fatty acid binding protein-1.

Moreover, our clinical studies (data not shown) showed that plasma MANF levels were negatively associated with blood glucose and TG levels and positively associated with high-density lipoprotein cholesterol levels, which further indicated that MANF may play a protective role in glucolipid metabolism. Moreover, a high circulating TG level is the key feature of insulin resistance and dyslipidemia, and TG levels are significantly different between patients with or without NAFLD^[18]. The accumulation of lipids, especially in the form of TGs, in the liver has been well accepted to set the stage for the progression of fatty liver disease and seems to be the hallmark of NAFLD^[19]. Most importantly, the lipid accumulation and steatosis of HepG2 cells were worsened by MANF deficiency and attenuated by upregulating MANF, especially under stimulation with FFAs. NAFLD is frequently initiated and progresses when the susceptible genotype interacts with environmental factors such




Figure 5 Mesencephalic astrocyte-derived neurotrophic factor knockdown increases lipogenesis. A and B: The intracellular TG (A) and TC (B) levels of HepG2 cells treated with bovine serum albumin (BSA) or free fatty acids (FFAs) for 72 h, ^a*P* < 0.05, ^b*P* < 0.001 vs Lv-GFP; ^d*P* < 0.01 vs Lv-GFP + FFAs; C and D: The mRNA expression levels of ACC α , FAS, SREBP-1C, and HMGCR in HepG2 cells treated with BSA (C) and with FFAs (D), ^b*P* < 0.01, ^c*P* < 0.001 vs Lv-GFP; ^e*P* < 0.05, ^f*P* < 0.01, ^g*P* < 0.001 vs Lv-GFP + FFAs; E and F: The mRNA expression of CD36 and FABP-1 in HepG2 cells treated with BSA (E) and with FFAs (F), ^b*P* < 0.01 vs Lv-GFP; ^h*P* < 0.01 vs Lv-GFP + FFAs. All data are presented as the mean \pm SE. All experiments were repeated three times. MANF: Mesencephalic astrocyte-derived neurotrophic factor; TG: Triglyceride; TC: Total cholesterol; Lv: Lentivirus; Lv-shMANF: Lentivirus expressing short hairpin RNA-targeted MANF; BSA: Bovine serum albumin; FFAs: Free fatty acids; ACC α : Acetyl-CoA carboxylase α ; FAS: Fatty acid synthase; SREBP-1C: Sterol regulatory element binding transcription factor 1C; HMGCR: 3-hydroxy3-methylglutaryl coenzyme A reductase; FABP-1: Fatty acid binding protein-1.

related to the fatty acid metabolism in the liver and SREBP-1c overexpression raised the hepatic TG accumulation. We have found that overexpression of MANF could downregulate this gene while MANF knockdown could increase SREBP-1c expression, which implies that MANF could affect SREBP-1c expression and thus have a potential role in the regulation of lipogenesis. Our results demonstrated that the downregulation of MANF could enhance these genes and result in the accumulation of lipids, while the upregulation of MANF could significantly downregulate these genes and attenuate HepG2 cell steatosis, showing improved accumulation of lipids, especially TGs. Moreover, HMGCR, a key enzyme responsible for cholesterol synthesis^[29], was reduced by MANF overexpression, indicating the advantageous effect of MANF in excessive cholesterol deposition. We detected genes

involved in fatty acid uptake in addition to those involved in fatty acid synthesis. Hepatic fatty acid uptake mainly depends on the FABP family and the scavenger receptor CD36. CD36 expression is much lower in normal hepatocytes but is greatly induced by high-fat diets, leading to TG accumulation and hepatic cell steatosis^[30-35], while disruption of its expression could attenuate NAFLD^[36]. FABP-1 is a liver-specific FABP, and moderate inhibition of FABP-1 function could ameliorate lipid accumulation in the liver^[37]. Most interestingly, modulating MANF expression could definitely affect fatty acid uptake by regulating CD36, and overexpression of MANF could inhibit FABP-1 expression, while MANF deficiency can increase FABP-1 expression. These results definitely showed that regulating MANF may offer an attractive therapeutic strategy for NAFLD.

There are a few limitations in this study. First, NAFLD encompasses simple steatosis to nonalcoholic steatohepatitis as well as further progression to cirrhosis and liver carcinoma, and we did not detect MANF expression in various stages of NAFLD. However, we verified the tendency for the first time in an *in vitro* study, and the gain and loss of function of the gene gave definite causal implications for MANF in the development of liver cell steatosis. Second, gain- and loss-of-function experiments involving MANF in an animal model of NAFLD still need to be performed. Nevertheless, the results from our experiments are sufficient to demonstrate a novel function of MANF in the pathogenesis of hepatic cell steatosis.

In conclusion, our study revealed a novel role of MANF in regulating hepatic lipid metabolism and steatosis in an *in vitro* model of NAFLD. This study is the first to indicate that hepatic MANF expression was induced upon FFAs overload and gradually decreased thereafter. MANF deficiency could significantly inhibit genes involved in lipogenesis and lipid uptake, potentially leading to attenuation of liver cell steatosis. Moreover, rescuing hepatic MANF may be a promising therapeutic target for the treatment of NAFLD, and additional intracellular mechanisms remain to be explored to fully understand the mechanism.

ARTICLE HIGHLIGHTS

Research background

Nonalcoholic fatty liver disease (NAFLD) is a global metabolism-associated liver disease. Hepatic steatosis, inflammation, and insulin resistance are the primary pathologic changes of NAFLD. Although NAFLD has become a serious threat to human public health, its pathogenic mechanisms are largely unknown.

Research motivation

Mesencephalic astrocyte-derived neurotrophic factor (MANF) is a newly discovered conserved neurotrophic factor. Recent studies found that MANF is involved in diabetes, obesity, and metabolic homeostasis. However, much remains to be discovered about its function in hepatic lipid metabolism; thus, we detected whether MANF could regulate hepatic metabolism.

Research objectives

This study aimed to detect MANF expression in both *in vivo* and *in vitro* NAFLD models and explore the role of MANF in hepatic lipid metabolism.

Research methods

HepG2 cells treated with free fatty acids and ob/ob mice were used as NAFLD models. Liver tissues collected from WT and ob/ob mice were used to detect MANF expression. Cells were treated with free fatty acids for different durations. Moreover, we established overexpression and knockdown cell models with lentiviruses to interfere with MANF expression levels in order to observe whether MANF influences hepatic steatosis.

Research results

Hepatic MANF expression was obviously decreased in ob/ob mice compared with wild type mice. The MANF level was elevated temporarily and gradually decreased in HepG2 cells as the duration of treatment with high free fatty acids increased. MANF deficiency accelerated lipogenesis and aggravated HepG2 cell steatosis, while MANF overexpression inhibited lipogenesis and rescued HepG2 cells from free fatty acids-induced steatosis.

Research conclusions

MANF alleviated lipid deposition and suppressed lipogenesis, showing a potential protective role in NAFLD.

Research perspectives

MANF may be a potential therapeutic target in hepatic steatosis processes.

REFERENCES

- 1 **Sahini N**, Borlak J. Recent insights into the molecular pathophysiology of lipid droplet formation in hepatocytes. *Prog Lipid Res* 2014; **54**: 86-112 [PMID: 24607340 DOI: 10.1016/j.plipres.2014.02.002]
- 2 **Petrova P**, Raibekas A, Pevsner J, Vigo N, Anafi M, Moore MK, Peaire AE, Shridhar V, Smith DI, Kelly J, Durocher Y, Commissiong JW. MANF: a new mesencephalic, astrocyte-derived neurotrophic factor with selectivity for dopaminergic neurons. *J Mol Neurosci* 2003; **20**: 173-188 [PMID: 12794311 DOI: 10.1385/jmn.20.2.173]
- 3 **Parkash V**, Lindholm P, Peränen J, Kalkkinen N, Oksanen E, Saarma M, Leppänen VM, Goldman A. The structure of the conserved neurotrophic factors MANF and CDFN explains why they are bifunctional. *Protein Eng Des Sel* 2009; **22**: 233-241 [PMID: 19258449 DOI: 10.1093/protein/gzn080]
- 4 **Lindahl M**, Saarma M, Lindholm P. Unconventional neurotrophic factors CDFN and MANF: Structure, physiological functions and therapeutic potential. *Neurobiol Dis* 2017; **97**: 90-102 [PMID: 27425895 DOI: 10.1016/j.nbd.2016.07.009]
- 5 **Lindholm P**, Peränen J, Andressoo JO, Kalkkinen N, Kokaia Z, Lindvall O, Timmusk T, Saarma M. MANF is widely expressed in mammalian tissues and differentially regulated after ischemic and epileptic insults in rodent brain. *Mol Cell Neurosci* 2008; **39**: 356-371 [PMID: 18718866 DOI: 10.1016/j.mcn.2008.07.016]
- 6 **Xu W**, Gao L, Li T, Zheng J, Shao A, Zhang J. Mesencephalic Astrocyte-Derived Neurotrophic Factor (MANF) Protects Against Neuronal Apoptosis via Activation of Akt/MDM2/p53 Signaling Pathway in a Rat Model of Intracerebral Hemorrhage. *Front Mol Neurosci* 2018; **11**: 176 [PMID: 29896089 DOI: 10.3389/fnmol.2018.00176]
- 7 **Zhang Z**, Shen Y, Luo H, Zhang F, Peng D, Jing L, Wu Y, Xia X, Song Y, Li W, Jin L. MANF protects dopamine neurons and locomotion defects from a human α -synuclein induced Parkinson's disease model in *C. elegans* by regulating ER stress and autophagy pathways. *Exp Neurol* 2018; **308**: 59-71 [PMID: 29959908 DOI: 10.1016/j.expneurol.2018.06.016]
- 8 **Hakonen E**, Chandra V, Fogarty CL, Yu NY, Ustinov J, Katayama S, Galli E, Danilova T, Lindholm P, Vartiainen A, Einarsdottir E, Krjutškov K, Kere J, Saarma M, Lindahl M, Otonkoski T. MANF protects human pancreatic beta cells against stress-induced cell death. *Diabetologia* 2018; **61**: 2202-2214 [PMID: 30032427 DOI: 10.1007/s00125-018-4687-y]
- 9 **Lindahl M**, Danilova T, Palm E, Lindholm P, Vöikar V, Hakonen E, Ustinov J, Andressoo JO, Harvey BK, Otonkoski T, Rossi J, Saarma M. MANF is indispensable for the proliferation and survival of pancreatic β cells. *Cell Rep* 2014; **7**: 366-375 [PMID: 24726366 DOI: 10.1016/j.celrep.2014.03.023]
- 10 **Wu T**, Zhang F, Yang Q, Zhang Y, Liu Q, Jiang W, Cao H, Li D, Xie S, Tong N, He J. Circulating mesencephalic astrocyte-derived neurotrophic factor is increased in newly diagnosed prediabetic and diabetic patients, and is associated with insulin resistance. *Endocr J* 2017; **64**: 403-410 [PMID: 28216543 DOI: 10.1507/endocrj.EJ16-0472]
- 11 **Galli E**, Härkönen T, Sainio MT, Ustav M, Toots U, Urtti A, Yliperttula M, Lindahl M, Knip M, Saarma M, Lindholm P. Increased circulating concentrations of mesencephalic astrocyte-derived neurotrophic factor in children with type 1 diabetes. *Sci Rep* 2016; **6**: 29058 [PMID: 27356471 DOI: 10.1038/srep29058]
- 12 **Yavarna T**, Al-Dewik N, Al-Mureikhi M, Ali R, Al-Mesaifri F, Mahmoud L, Shahbeck N, Lakhani S, AlMulla M, Nawaz Z, Vitazka P, Alkuraya FS, Ben-Omran T. High diagnostic yield of clinical exome sequencing in Middle Eastern patients with Mendelian disorders. *Hum Genet* 2015; **134**: 967-980 [PMID: 26077850 DOI: 10.1007/s00439-015-1575-0]
- 13 **Yang S**, Yang H, Chang R, Yin P, Yang Y, Yang W, Huang S, Gaertig MA, Li S, Li XJ. MANF regulates hypothalamic control of food intake and body weight. *Nat Commun* 2017; **8**: 579 [PMID: 28924165 DOI: 10.1038/s41467-017-00750-x]
- 14 **Sousa-Victor P**, Neves J, Cedron-Craft W, Ventura PB, Liao CY, Riley RR, Soifer I, van Bruggen N, Kolumam GA, Villeda SA, Lamba DA, Jasper H. MANF regulates metabolic and immune homeostasis in ageing and protects against liver damage. *Nat Metab* 2019; **1**: 276-290 [PMID: 31489403 DOI: 10.1038/s42255-018-0023-6]
- 15 **Feldstein AE**, Werneburg NW, Canbay A, Guicciardi ME, Bronk SF, Rydzewski R, Burgart LJ, Gores GJ. Free fatty acids promote hepatic lipotoxicity by stimulating TNF- α expression via a lysosomal pathway. *Hepatology* 2004; **40**: 185-194 [PMID: 15239102 DOI: 10.1002/hep.20283]
- 16 **Wen J**, Lin T, Wu W, Yang Y, Luo C, Zhou C, Wan J, Liu S, Wang D, Wang P, Li J. Tiaopi huxin recipe improved endothelial dysfunction and attenuated atherosclerosis by decreasing the expression of caveolin-1 in ApoE-deficient mice. *J Cell Physiol* 2019; **234**: 234 [PMID: 30729525 DOI: 10.1002/jcp.28184]
- 17 **Zhang L**, Yang M, Ren H, Hu H, Boden G, Li L, Yang G. GLP-1 analogue prevents NAFLD in ApoE KO mice with diet and Acrp30 knockdown by inhibiting c-JNK. *Liver Int* 2013; **33**: 794-804 [PMID: 23432843 DOI: 10.1111/liv.12120]
- 18 **Trojak A**, Waluś-Miarka M, Woźniakiewicz E, Malecki MT, Idzior-Waluś B. Nonalcoholic fatty liver disease is associated with low HDL cholesterol and coronary angioplasty in patients with type 2 diabetes. *Med Sci Monit* 2013; **19**: 1167-1172 [PMID: 24336007 DOI: 10.12659/MSM.889649]
- 19 **Day CP**, James OF. Steatohepatitis: a tale of two "hits"? *Gastroenterology* 1998; **114**: 842-845 [PMID: 9547102 DOI: 10.1016/s0016-5085(98)70599-2]
- 20 **Anstee QM**, Day CP. The genetics of NAFLD. *Nat Rev Gastroenterol Hepatol* 2013; **10**: 645-655 [PMID: 24061205 DOI: 10.1038/nrgastro.2013.182]
- 21 **Anstee QM**, Seth D, Day CP. Genetic Factors That Affect Risk of Alcoholic and Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2016; **150**: 1728-1744.e7 [PMID: 26873399 DOI: 10.1053/j.gastro.2016.01.037]
- 22 **Brownsey RW**, Zhande R, Boone AN. Isoforms of acetyl-CoA carboxylase: structures, regulatory properties and metabolic functions. *Biochem Soc Trans* 1997; **25**: 1232-1238 [PMID: 9449982 DOI: 10.1042/bst0251232]
- 23 **Goedeke L**, Bates J, Vatner DF, Perry RJ, Wang T, Ramirez R, Li L, Ellis MW, Zhang D, Wong KE, Beyens C, Cline GW, Ray AS, Shulman GI. Acetyl-CoA Carboxylase Inhibition Reverses NAFLD and Hepatic Insulin Resistance but Promotes Hypertriglyceridemia in Rodents. *Hepatology* 2018; **68**: 2197-2211 [PMID: 29790582 DOI: 10.1002/hep.30097]
- 24 **Shimano H**. Sterol regulatory element-binding proteins (SREBPs): transcriptional regulators of lipid synthetic genes. *Prog Lipid Res* 2001; **40**: 439-452 [PMID: 11591434 DOI: 10.1016/s0163-7827(01)00010-8]

- 25 **Horton JD**, Goldstein JL, Brown MS. SREBPs: activators of the complete program of cholesterol and fatty acid synthesis in the liver. *J Clin Invest* 2002; **109**: 1125-1131 [PMID: [11994399](#) DOI: [10.1172/JCI15593](#)]
- 26 **Nakamuta M**, Kohjima M, Morizono S, Kotoh K, Yoshimoto T, Miyagi I, Enjoji M. Evaluation of fatty acid metabolism-related gene expression in nonalcoholic fatty liver disease. *Int J Mol Med* 2005; **16**: 631-635 [PMID: [16142397](#) DOI: [10.3892/ijmm.16.4.631](#)]
- 27 **Kohjima M**, Enjoji M, Higuchi N, Kato M, Kotoh K, Yoshimoto T, Fujino T, Yada M, Yada R, Harada N, Takayanagi R, Nakamuta M. Re-evaluation of fatty acid metabolism-related gene expression in nonalcoholic fatty liver disease. *Int J Mol Med* 2007; **20**: 351-358 [PMID: [17671740](#) DOI: [10.3892/ijmm.20.3.351](#)]
- 28 **Horton JD**, Bashmakov Y, Shimomura I, Shimano H. Regulation of sterol regulatory element binding proteins in livers of fasted and refed mice. *Proc Natl Acad Sci USA* 1998; **95**: 5987-5992 [PMID: [9600904](#) DOI: [10.1073/pnas.95.11.5987](#)]
- 29 **Ma S**, Sun W, Gao L, Liu S. Therapeutic targets of hypercholesterolemia: HMGR and LDLR. *Diabetes Metab Syndr Obes* 2019; **12**: 1543-1553 [PMID: [31686875](#) DOI: [10.2147/DMSO.S219013](#)]
- 30 **Bechmann LP**, Gieseler RK, Sowa JP, Kahraman A, Erhard J, Wedemeyer I, Emons B, Jochum C, Feldkamp T, Gerken G, Canbay A. Apoptosis is associated with CD36/fatty acid translocase upregulation in non-alcoholic steatohepatitis. *Liver Int* 2010; **30**: 850-859 [PMID: [20408954](#) DOI: [10.1111/j.1478-3231.2010.02248.x](#)]
- 31 **Miquilena-Colina ME**, Lima-Cabello E, Sánchez-Campos S, García-Mediavilla MV, Fernández-Bermejo M, Lozano-Rodríguez T, Vargas-Castrillón J, Buqué X, Ochoa B, Aspichueta P, González-Gallego J, García-Monzón C. Hepatic fatty acid translocase CD36 upregulation is associated with insulin resistance, hyperinsulinaemia and increased steatosis in non-alcoholic steatohepatitis and chronic hepatitis C. *Gut* 2011; **60**: 1394-1402 [PMID: [21270117](#) DOI: [10.1136/gut.2010.222844](#)]
- 32 **Greco D**, Kotronen A, Westerbacka J, Puig O, Arkkila P, Kiviluoto T, Laitinen S, Kolak M, Fisher RM, Hamsten A, Auvinen P, Yki-Järvinen H. Gene expression in human NAFLD. *Am J Physiol Gastrointest Liver Physiol* 2008; **294**: G1281-G1287 [PMID: [18388185](#) DOI: [10.1152/ajpgi.00074.2008](#)]
- 33 **Su X**, Abumrad NA. Cellular fatty acid uptake: a pathway under construction. *Trends Endocrinol Metab* 2009; **20**: 72-77 [PMID: [19185504](#) DOI: [10.1016/j.tem.2008.11.001](#)]
- 34 **Koonen DP**, Jacobs RL, Febbraio M, Young ME, Soltys CL, Ong H, Vance DE, Dyck JR. Increased hepatic CD36 expression contributes to dyslipidemia associated with diet-induced obesity. *Diabetes* 2007; **56**: 2863-2871 [PMID: [17728375](#) DOI: [10.2337/db07-0907](#)]
- 35 **Zhou J**, Febbraio M, Wada T, Zhai Y, Kuruba R, He J, Lee JH, Khadem S, Ren S, Li S, Silverstein RL, Xie W. Hepatic fatty acid transporter Cd36 is a common target of LXR, PXR, and PPARgamma in promoting steatosis. *Gastroenterology* 2008; **134**: 556-567 [PMID: [18242221](#) DOI: [10.1053/j.gastro.2007.11.037](#)]
- 36 **Wilson CG**, Tran JL, Erion DM, Vera NB, Febbraio M, Weiss EJ. Hepatocyte-Specific Disruption of CD36 Attenuates Fatty Liver and Improves Insulin Sensitivity in HFD-Fed Mice. *Endocrinology* 2016; **157**: 570-585 [PMID: [26650570](#) DOI: [10.1210/en.2015-1866](#)]
- 37 **Mukai T**, Egawa M, Takeuchi T, Yamashita H, Kusudo T. Silencing of FABP1 ameliorates hepatic steatosis, inflammation, and oxidative stress in mice with nonalcoholic fatty liver disease. *FEBS Open Bio* 2017; **7**: 1009-1016 [PMID: [28680813](#) DOI: [10.1002/2211-5463.12240](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

