

Preoperative albumin levels predict prolonged postoperative i



ALL

IMAGES

VIDEOS

关闭取词

30,300 Results

Any time ▾

Implications of preoperative ... - PubMed Central (PMC)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4872063>

May 27, 2016 · **Prolonged postoperative ileus symptoms** include **nausea and vomiting**, inability to tolerate an oral diet, abdominal distention, and delayed passage of flatus and stool[44,45]. **Decreasing preoperative albumin** was a **predictor for prolonged postoperative ileus** (OR = 1.11 per gram per litre unit change; 95%CI: 1.02-1.22; P = 0.047).

Cited by: 17**Author:** Adam Truong, Mark H Hanna, Zhobin Mo...**Publish Year:** 2016

Preoperative Serum Albumin Level as a Predictor of ...

<https://jamanetwork.com/journals/jamasurgery/fullarticle/390180> ▾

Both **serum albumin levels and weight loss** appear in the mortality model for **thoracic surgery**, indicating that **albumin level detects risk** not associated with weight **loss** of greater than 10% of body weight in the 6 months prior to **surgery**.

Cited by: 1005**Author:** James Gibbs, William Cull, William Hend...**Publish Year:** 1999

Postoperative decrease of serum albumin predicts short ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5526768>

Jul 21, 2017 · Core tip: In this work, we investigated whether postoperative decrease of serum **albumin** can **predict** short-term complications following gastric cancer resection. Results indicate that the decrease of serum **albumin** could be more accurate than postoperative C-reactive protein in predicting complications after gastrectomy.

Cited by: 3**Author:** Zhi-Jian Liu, Xiao-Long Ge, Shi-Chao Ai, ...**Publish Year:** 2017



53473-Review.docx

Quotes Excluded
Bibliography Excluded

4%
SIMILAR

Retrospective Cohort Study

Preoperative albumin levels predict prolonged postoperative ileus in gastrointestinal surgery

Liang WQ *et al.* Preoperative albumin predicts PPOI in GI surgery

² Wen-Quan Liang, Ke-Cheng Zhang, Hua Li, Jian-Xin Cui, Hong-Qing Xi, Ji-Yang Li, Ai-Zhen Cai, Yu-Hua Liu, Wang Zhang, Lan Zhang, Bo Wei, Lin Chen

Abstract

BACKGROUND

Prolonged postoperative ileus (PPOI) is a prolonged status of “pathological” gastrointestinal (GI) tract dysmotility. There are relatively few studies

Match Overview

1	Internet 28 words crawled on 26-Jan-2019 peerj.com	1%
2	Internet 24 words crawled on 29-Oct-2019 www.wjgnet.com	1%
3	Crossref 22 words Derek B. Hennessey, John P. Burke, Tara Ni-Dhonochu, Connor Shields, Desmond C. Winter, Kenneth Mealy. "Preoper	1%
4	Internet 20 words crawled on 16-Feb-2016 prognosismethods.cochrane.org	1%
5	Crossref 19 words Demars, Sandra M.. "Postoperative Ileus", The Perioperative Medicine Consult Handbook, 2015.	1%
6	Internet 18 words crawled on 31-Jul-2018 medlineplus.gov	1%

Preoperative albumin levels predict prolonged postoperative i



ALL

IMAGES

VIDEOS

关闭取词

30,800 Results

Any time ▼

Preoperative Serum Albumin Level as a Predictor of ...

<https://jamanetwork.com/journals/jamasurgery/fullarticle/390180> ▼

Both **serum albumin levels** and **weight loss** appear in the mortality model for **thoracic surgery**, indicating that **albumin level detects risk** not associated with **weight loss** of greater than 10% of body weight in the 6 months prior to **surgery**.

Cited by: 1004

Author: James Gibbs, William Cull, William Hend...

Publish Year: 1999

Implications of preoperative hypoalbuminemia in colorectal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4872063>

May 27, 2016 · **Prolonged postoperative ileus symptoms** include **nausea and vomiting**, inability to tolerate an oral diet, abdominal distention, and delayed passage of flatus and stool[44,45]. **Decreasing preoperative albumin** was a **predictor for prolonged postoperative ileus** (OR = 1.11 per gram per litre unit change; 95%CI: 1.02-1.22; P = 0.047).

Cited by: 17

Author: Adam Truong, Mark H Hanna, Zhobin Mo...

Publish Year: 2016

Postoperative ileus: Pathophysiology, incidence, and



国内版

国际版

Preoperative albumin levels predict prolonged postopera



Sign in



ALL

IMAGES

VIDEOS

关闭取词

Add Bing Firefox extension >

30,800 Results

Any time ▾

[Implications of preoperative ... - PubMed Central \(PMC\)](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4872063>

May 27, 2016 · **Prolonged postoperative ileus symptoms** include **nausea and vomiting**, inability to tolerate an oral diet, abdominal distention, and delayed passage of flatus and stool[44,45]. **Decreasing preoperative albumin** was a **predictor for prolonged postoperative ileus** (OR = 1.11 per gram per litre unit change; 95%CI: 1.02-1.22; P = 0.047).

Cited by: 17

Author: Adam Truong, Mark H Hanna, Zhobin Mo...

Publish Year: 2016

[Preoperative Serum Albumin Level as a Predictor of ...](#)

<https://jamanetwork.com/journals/jamasurgery/fullarticle/390180> ▾

Both **serum albumin levels and weight loss** appear in the mortality model for **thoracic surgery**, indicating that **albumin level detects risk** not associated with weight loss of greater than 10% of body weight in the 6 months prior to **surgery**.

Cited by: 1004

Author: James Gibbs, William Cull, William Hend...

Publish Year: 1999

[Cytokine levels in abdominal exudate predict prolonged](#)