


Match Overview 

1	Crossref 53 words G. Scozzari, A. Arezzo, M. Morino. "Enterovesical fistulas: diagnosis and management", <i>Techniques in Coloproctology</i> , 2010	1%
2	Crossref 42 words Hechenbleikner, Elizabeth M., Jill C. Buckley, and Elizabeth C. Wick. "Acquired Rectourethral Fistulas in Adults : A Systematic Review"	1%
3	Internet 12 words crawled on 01-May-2020 worldwidescience.org	<1%

Name of Journal: *World Journal of Gastrointestinal Surgery*

Manuscript NO: 53919

Manuscript Type: REVIEW

When the bowel meets the bladder: Optimal management of colorectal pathology with urological involvement

Abstract

Fistulae between the gastrointestinal and urinary systems are rare but becoming increasingly more common in current surgical practice. They are a heterogeneous group of pathological entities that are uncommon complications of both benign and malignant processes. As the incidence of complicated diverticular disease and



When the bowel meets the bladder: Optimal management of



ALL

IMAGES

VIDEOS

关闭取词

15,400 Results

Any time ▾

Do the urinary bladder and large bowel interact, in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3309116>

On the other hand, treatment of overactive bladder with bladder relaxing drugs can cause constipation.

10 The same was found in patients with **neurogenic** pathology. 11 Urinary tract infections are quite common in patients with LUT dysfunctions and treatment with antibiotics can cause bowel problems. 12 The use of bowel segments for the treatment of urinary disorders such as bladder enlargement or replacement ...

Cited by: 31

Author: Anna P. Malykhina, Jean-Jacques Wynda...

Publish Year: 2012

Bladder and Bowel Dysfunction | Johns Hopkins Medicine

<https://www.hopkinsmedicine.org/health/conditions-and-diseases/bladder-and-bowel...> ▾

Bladder and **bowel** problems often originate with nerve or muscle dysfunction, as these systems control the flow of urine and the release of stool. Other health issues may cause **bladder** and/or **bowel** dysfunction, including medicinal side effects, stress, neurologic diseases, diabetes, hemorrhoids and pelvic floor disorders.

Constipation - 5 "Little" Health Issues That May Be Bigger Than You Think

Physiology and Management of Bladder and Bowel ...

<https://www.o-wm.com/content/physiology-and...> ▾

Dec 12, 2007 · An **effective management program requires the advanced knowledge and understanding of the physiology of normal bowel and bladder function as well as common alterations in function the continence care clinician can provide.** 9 The Wound Ostomy Continence Nurses (WOCN) Society and the Society of Urologic Nurses and Associates (SUNA) offer an avenue ...



国内版 国际版

When the bowel meets the bladder: Optimal managemen



Sign in



ALL

IMAGES

VIDEOS

Add the Give with Bing extension >

15,700 Results

Any time ▾

Do the urinary bladder and large bowel interact, in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3309116>

On the other hand, treatment of overactive bladder with bladder **relaxing drugs can cause constipation**. 10 The same was found in patients with neurogenic pathology. 11 Urinary tract infections are quite common in patients with LUT dysfunctions and treatment with antibiotics can cause bowel problems. 12 The use of bowel segments for the treatment of urinary disorders such as bladder ...

Cited by: 31

Author: Anna P. Malykhina, Jean-Jacques Wynda...

Publish Year: 2012

Endometriosis and Vesico-Sphincteral Disorders

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4476201>

In another study performed by Lapasse et al. concerning 12 patients with deep **endometriosis** of the posterior **pelvic wall**, the anamnesis highlighted vesico-sphincteral symptoms among 50% of the patients (day-time and night-time pollakiuria, Urge **incontinence**, **Bladder** voiding by thirst, lower back pain, dysuria, full **bladder cramps** or pain, cramps or pain during or at end of urination, **stress incontinence**, decreased **bladder** ...

Cited by: 9

Author: Anis Fadhlaoui, Tessa Gillon, Issam Lebb...

Publish Year: 2015

Management of Lower Urinary Tract Dysfunction in Patients ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3459999>

Sep 19, 2012 · The **lower urinary tract** consists of the bladder, bladder outlet, and urethra. It serves two major functions: 1) storage of **urine at low pressure with normal sensation and perfect continence**, and 2) periodic and complete voluntary elimination of urine [1,2]. The proper performance of these





When the bowel meets the bladder: Optimal manageme



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

15,500 Results

Any time ▾

[Do the urinary bladder and large bowel interact, in ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3309116>

On the other hand, treatment of overactive bladder with bladder **relaxing drugs can cause constipation**. 10 The same was found in patients with neurogenic pathology. 11 Urinary tract infections are quite common in patients with LUT dysfunctions and treatment with antibiotics can cause bowel problems. 12 The use of bowel segments for the treatment of urinary disorders such as bladder ...

Cited by: 31

Author: Anna P. Malykhina, Jean-Jacques Wynda...

Publish Year: 2012

[Physiology and Management of Bladder and Bowel Continence ...](#)

<https://www.o-wm.com/content/physiology-and...>

An effective management program requires the advanced knowledge and understanding of **the physiology of normal bowel and bladder function as well as common alterations in function the continence care clinician can provide**. 9 The Wound Ostomy Continence Nurses (WOCN) Society and the Society of Urologic Nurses and Associates (SUNA) offer an avenue for specialty certification in ...

[Neurogenic bowel dysfunction in patients with spinal cord ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3235587>

Dec 14, 2011 · EPIDEMIOLOGY. Traumatic SCI represents a significant public health problem worldwide[.Each year, 11 000 individuals are estimated to have SCI in the United States] with a mortality rate of 27.4 per million people.An annual incidence of 33.6 per million is reported in Greece and 19.5 per million in Sweden[, while in Denmark the number of SCI patients is about 3000.

Cited by: 63

Author: Richard A Awad

Publish Year: 2011

[Evaluation and Treatment of Neurogenic Bowel Dysfunction ...](#)

<https://touchneurology.com/evaluation-and...>

A stepwise approach to bowel management, based on a systematic assessment is **effective in managing the intrusive gut symptoms that afflict the majority of patients with central neurological**