

17

Name of Journal: *World Journal of Gastroenterology***Manuscript NO:** 54448**Manuscript Type:** ORIGINAL ARTICLE*Retrospective Cohort Study***Medications in type-2 diabetics and their association with liver fibrosis**Siddiqui MT *et al.* T2D medications associated with liver fibrosis

Mohamed Tausif Siddiqui, Hina Amin, Rajat Garg, Pravallika Chadalavada, Wael Al-Yaman, Rocio Lopez, Amandeep Singh

Abstract**BACKGROUND**

The prevalence of nonalcoholic fatty liver disease (NAFLD) is significantly rising worldwide. Type-2 diabetes (T2D) is major risk factor for NAFLD progression.

AIM

10

To assess the association of nonalcoholic fatty liver disease and fibrosis (AF) in

Match Overview

1	Internet 78 words crawled on 03-Mar-2013 www.hindawi.com	2%
2	Crossref 61 words Kathleen E. Corey. "The effect of angiotensin-blocking a... nts on liver fibrosis in patients with hepatitis C", <i>Liver Inter</i>	2%
3	Internet 58 words crawled on 11-May-2020 doaj.org	2%
4	Internet 54 words crawled on 20-Dec-2019 link.springer.com	2%
5	Internet 46 words crawled on 18-Jan-2020 journals.sagepub.com	1%
6	Internet 40 words crawled on 13-Jun-2017 www.wjgnet.com	1%
7	Internet 35 words crawled on 28-Apr-2020 research.birmingham.ac.uk	1%
8	Internet 34 words crawled on 04-Jul-2019 www.nature.com	1%
9	Crossref 28 words Kaveh Hajifathalian, Monica Saumoy, Enad Dawod, Shaw n L. Shah et al. "Sa1885 - Weight Loss and Improvem..."	1%
10	Crossref 27 words "Posters (Abstracts 301-2389)", <i>Hepatology</i> , 2018	1%
11	Internet 25 words crawled on 13-Apr-2016 www.ajol.info	1%
12	Internet 22 words crawled on 23-May-2020	1%



Medications in Type-2 Diabetics and their association with



ALL

IMAGES

VIDEOS

209,000 Results

Any time ▾

Statins, antidiabetic medications and liver histology in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4838660>

Mar 18, 2016 · **Non-alcoholic fatty liver disease (NAFLD)** covers a spectrum of liver injury strongly associated with **type-2 diabetes mellitus (T2DM)** as the prevalence of **NAFLD in diabetics** ranges from 40% to 70%. 1 2 Importantly, **NAFLD** is also associated with cardiovascular disease, and this **association** is believed to be independent of traditional cardiovascular risk factors. 3 **NAFLD ...**

Cited by: 52

Author: Fabio Nascimbeni, Judith Aron-Wisne...

Publish Year: 2016

Diabetes drugs for nonalcoholic fatty liver disease: a ...

<https://systematicreviewsjournal.biomedcentral.com/articles/10.1186/s13643-019-1200-8>

Nov 29, 2019 · Background. **Fatty liver** is **associated** with obesity, **type 2 diabetes**, hyperlipidemia, hypertension, and metabolic syndrome. While there are no approved **drugs** for the treatment of **nonalcoholic fatty liver disease (NAFLD)** or **nonalcoholic** steatohepatitis, strategies to ameliorate **fatty liver** often target these **related** diseases.

Author: Ian Blazina, Shelley Selph **Publish Year:** 2019

Non-alcoholic fatty liver disease and diabetes

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4943559>

Jul 30, 2015 · Non-alcoholic **fatty liver disease (NAFLD)** and **type 2 diabetes (T2DM)** are common conditions that regularly co-exist and can act synergistically to drive adverse outcomes. The presence of both **NAFLD** and **T2DM** increases the likelihood of the development of ...

Cited by: 160

Author: Jonathan M. Hazlehurst, Conor Woods...

Publish Year: 2016

Medications in type-2 diabetics and their association with liver fi



ALL

IMAGES

VIDEOS

20,600,000 Results

Any time ▾

Including results for **medications in type-2 diabetes** and their association with liver fibrosis.

Do you want results only for **Medications in type-2 diabetics and their association with liver fibrosis**?

Diabetes Medications Have Different Effects on the Liver

<https://www.medscape.com> > [viewarticle](#) > [887648](#)

Apr 30, 2020 · WASHINGTON, DC — For people with **advanced liver fibrosis** due to **nonalcoholic fatty liver disease (NAFLD)** who also have **type 2 diabetes**, outcomes are better with **metformin** than with **sulfonylureas**,...

Medications for Type 2 Diabetes - WebMD

<https://www.webmd.com> > [diabetes](#) > [story](#) > [type-2-treatments](#)

It's unlikely your doctor would choose insulin first as a treatment for your **type 2 diabetes**. The timing of doses can be tricky. But it can be a key player when blood sugar is hard to control.

Drug reverses signs of liver disease in people living with ...

<https://www.nih.gov> > [news-events](#) > [news-releases](#) > [drug...](#)

Oct 15, 2019 · Obesity and **type 2 diabetes** raise the risk of developing NAFLD regardless of HIV status, and people with HIV are at increased risk of NAFLD because some HIV **medications** and HIV itself are **associated** with gaining abdominal fat and may contribute to **liver** fat build-up. This research was supported through NIAID grant U01 AI115711.

PEOPLE ALSO ASK

Is type 2 diabetes a risk factor for liver fibrosis? ▾

Is there discrimination in between Type 2 diabetes and hepatogenous diabetes? ▾



ALL IMAGES VIDEOS MAPS NEWS SHOPPING

20,500,000 Results Any time ▾

Including results for medications in type-2 **diabetes** and their association with liver fibrosis.

Do you want results only for Medications in type-2 diabetics and their association with liver fibrosis?

Diabetes Medications Have Different Effects on the Liver

<https://www.medscape.com> › [viewarticle](#) › [887648](#)

Apr 30, 2020 · WASHINGTON, DC — For people with **advanced liver fibrosis** due to **nonalcoholic fatty liver disease (NAFLD)** who also have **type 2 diabetes**, outcomes are better with **metformin** than with **sulfonylureas**,...

Statins, antidiabetic medications and liver histology in ...

<https://www.ncbi.nlm.nih.gov> › [pmc](#) › [articles](#) › [PMC4838660](#)

Model 1 included a single drug among **statins**, **metformin**, **sulfonylureas** or **insulin**; model 2 included a number of antidiabetic drugs and a single drug among **statins**, **metformin**, **sulfonylureas** or **insulin**; model 3 included **statins** and **antidiabetic** (**metformin**, **sulfonylureas** and **insulin**...

Cited by: 59 **Author:** Fabio Nascimbeni, Judith Aron-Wisnewsk...

Publish Year: 2016

Medications for Type 2 Diabetes - WebMD

<https://www.webmd.com> › [diabetes](#) › [story](#) › [type-2-treatments](#)

It's unlikely your doctor would choose insulin first as a treatment for your type 2 **diabetes**. The timing of doses can be tricky. But it can be a key player when blood sugar is hard to control.

PEOPLE ALSO ASK

Is type 2 diabetes a risk factor for liver fibrosis? ▾

What medications do people with Type 2 diabetes take? ▾

Can diabetes medicine help with fatty liver disease? ▾

See results for

Fatty Liver (Medical Condition)

Excessive build-up of fat in the liver.