

6

Name of Journal: *World Journal of Clinical Cases***Manuscript NO:** 54783**Manuscript Type:** CASE REPORT

Diagnosis of an actively bleeding brachial artery hematoma by contrast-enhanced ultrasound

Jiao-Jiao Ma, Bo Zhang

Abstract

BACKGROUND

Match Overview

1	Internet 47 words crawled on 07-May-2020 www.wjgnet.com	3%
2	Internet 34 words crawled on 03-May-2020 www.ncbi.nlm.nih.gov	2%
3	Internet 25 words crawled on 28-Feb-2020 www.mayoclinic.org	2%
4	Internet 23 words crawled on 23-May-2020 www.frontiersin.org	1%
5	Crossref 21 words Alexandra A. Roman, Andre J. Arsenault, Kenneth D. Jackson, John M. Price. "Novel Onset of a Posttraumatic Super..."	1%
6	Internet 15 words crawled on 15-Feb-2020 f6publishing.blob.core.windows.net	1%
7	Crossref 13 words Daobao Zhang, Zhiyong Liu, Jianguo Xu. "A giant pseudoaneurysm from superficial temporal artery after trauma", World Journal of Clinical Cases	1%
8	Crossref 12 words Jiao-Jiao Ma, Bo Zhang. "A giant pseudoaneurysm from superficial temporal artery after trauma", World Journal of Clinical Cases	1%

Diagnosis of an actively bleeding brachial artery hematoma by cont



ALL

IMAGES

VIDEOS

31,100 Results

Any time ▾

Iatrogenic Percutaneous Vascular Injuries: Clinical ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4447873>

Hematoma and Retroperitoneal **Hemorrhage**. Postprocedural **bleeding** or **hematoma** occurs at a rate of 2 to 12%, but those that are significant enough to require transfusion occur in fewer than 1% of cases. 10 11 Hematomas most often occur at the arteriotomy site, resulting in induration and discoloration. Treatment includes local compression to achieve hemostasis and reversal of precipitating ...

Cited by: 9

Author: Benjamin H. Ge, Alexander Copelan, Do...

Publish Year: 2015

[PDF] Case Report A Giant Brachial Artery Pseudoaneurysm in ...

<https://www.scireslit.com/VascularMedicine/AJVM-ID23.pdf>

suggested that the vascular lesion was compatible with an **artery** pseudoaneurysm. Color duplex **ultrasound** and **contrast-enhanced** Computed Tomography Angiography (CTA) demonstrated a 3.0 × 2.8 × 3.0 cm mass arising from the left **brachial artery** (Figure 2). The left **brachial artery** pseudoaneurysm was surgically resected,

Rectus Sheath Haematoma Associated with Low Molecular ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1964713>

Arteriogram performed via left **brachial artery** (femoral **artery** obscured by haematoma) showing successful embolisation of the superior epigastric **artery** (3 × 3 mm coil). Despite cessation of **bleeding** and a stable haemoglobin, respiratory pressures continued to increase, inotropic support escalated and the patient became cyanotic and acidotic.

Cited by: 62

Publish Year: 2007



Download the new Microsoft Edge to get the best of the web.



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

31,000 Results

Any time ▾

Iatrogenic Percutaneous Vascular Injuries: Clinical ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4447873>

Hematoma and Retroperitoneal **Hemorrhage**. Postprocedural **bleeding** or **hematoma** occurs at a rate of 2 to 12%, but those that are significant enough to require transfusion occur in fewer than 1% of cases. 10 11 Hematomas most often occur at the arteriotomy site, resulting in induration and discoloration. Treatment includes local compression to achieve hemostasis and reversal of precipitating ...

Cited by: 9

Author: Benjamin H. Ge, Alexander Copelan, Do...

Publish Year: 2015

[PDF] Case Report A Giant Brachial Artery Pseudoaneurysm in an ...

<https://www.scireslit.com/VascularMedicine/AJVM-ID23.pdf>

suggested that the vascular lesion was compatible with an **artery** pseudoaneurysm. Color duplex **ultrasound** and **contrast-enhanced** Computed Tomography Angiography (CTA) demonstrated a 3.0 × 2.8 × 3.0 cm mass arising from the left **brachial artery** (Figure 2). The left **brachial artery** pseudoaneurysm was surgically resected,

Rectus Sheath Haematoma Associated with Low Molecular ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1964713>

Arteriogram performed via left **brachial artery** (femoral **artery** obscured by haematoma) showing successful embolisation of the superior epigastric **artery** (3 × 3 mm coil). Despite cessation of **bleeding** and a stable haemoglobin, respiratory pressures continued to increase, inotropic support escalated and the patient became oliguric and acidotic.

Cited by: 64

Author: J Donaldson, CH Knowles, SK Clark, I R...

Publish Year: 2007

Ultrasound imaging of the arterial system - Gupta ...

cdt.amegroups.com/article/view/24428/24957 ▾

Abstract: **Ultrasound** (US) and Doppler are often the first imaging to be carried out for arterial disease, and with good reason. US is noninvasive, free of radiation exposure and crucial for follow up imaging. The review that follows aims to highlight the various applications of US in imaging of the arterial system.

Cited by: 2

Author: Prashant Gupta, Shannon Lyons, Sandee...

Publish Year: 2019