

World Journal of *Clinical Cases*

World J Clin Cases 2020 August 6; 8(15): 3136-3376



OPINION REVIEW

- 3136 Impacts and challenges of United States medical students during the COVID-19 pandemic
Rolak S, Keefe AM, Davidson EL, Aryal P, Parajuli S
- 3142 Recent advances in the management of gastrointestinal stromal tumor
Ahmed M
- 3156 Medical research during the COVID-19 pandemic
AlNaamani K, AlSinani S, Barkun AN

REVIEW

- 3164 Progress of intravoxel incoherent motion diffusion-weighted imaging in liver diseases
Tao YY, Zhou Y, Wang R, Gong XQ, Zheng J, Yang C, Yang L, Zhang XM

MINIREVIEWS

- 3177 Typical and atypical COVID-19 computed tomography findings
Caruso D, Polidori T, Guido G, Nicolai M, Bracci B, Cremona A, Zerunian M, Polici M, Pucciarelli F, Rucci C, Dominici CD, Girolamo MD, Argento G, Sergi D, Laghi A
- 3188 Review of possible psychological impacts of COVID-19 on frontline medical staff and reduction strategies
Fu XW, Wu LN, Shan L

ORIGINAL ARTICLE**Clinical and Translational Research**

- 3197 Overexpression of AMPD2 indicates poor prognosis in colorectal cancer patients *via* the Notch3 signaling pathway
Gao QZ, Qin Y, Wang WJ, Fei BJ, Han WF, Jin JQ, Gao X

Case Control Study

- 3209 Effect of motivational interviewing on postoperative weight control in patients with obstructive sleep apnea-hypopnea syndrome
Sun XH, Xue PS, Qi XX, Fan L

Retrospective Study

- 3218 Thalidomide for refractory gastrointestinal bleeding from vascular malformations in patients with significant comorbidities
Bayudan AM, Chen CH

- 3230** Colorectal adenocarcinoma patients with M1a diseases gain more clinical benefits from palliative primary tumor resection than those with M1b diseases: A propensity score matching analysis

Li CL, Tang DR, Ji J, Zang B, Chen C, Zhao JQ

- 3240** Surgical outcomes of bladder augmentation: A comparison of three different augmentation procedures

Sun XG, Wang RY, Xu JL, Li DG, Chen WX, Li JL, Wang J, Li AW

Clinical Trials Study

- 3249** Comparison of measurements of anterior chamber angle *via* anterior segment optical coherence tomography and ultrasound biomicroscopy

Yu ZY, Huang T, Lu L, Qu B

Observational Study

- 3259** Dydrogesterone treatment for menstrual-cycle regularization in abnormal uterine bleeding – ovulation dysfunction patients

Wang L, Guan HY, Xia HX, Chen XY, Zhang W

CASE REPORT

- 3267** Multi-organ IgG4-related disease continues to mislead clinicians: A case report and literature review

Strainiene S, Sarlauskas L, Savlan I, Liakina V, Stundiene I, Valantinas J

- 3280** *Campylobacter jejuni* enterocolitis presenting with testicular pain: A case report

Sanagawa M, Kenzaka T, Kato S, Yamaoka I, Fujimoto S

- 3284** Natural killer/T-cell lymphoma with intracranial infiltration and Epstein-Barr virus infection: A case report

Li N, Wang YZ, Zhang Y, Zhang WL, Zhou Y, Huang DS

- 3291** Successful management of tubular colonic duplication using a laparoscopic approach: A case report and review of the literature

Li GB, Han JG, Wang ZJ, Zhai ZW, Tao Y

- 3299** Hypothyroidism with elevated pancreatic amylase and lipase without clinical symptoms: A case report

Xu YW, Li R, Xu SC

- 3305** Two mechanically ventilated cases of COVID-19 successfully managed with a sequential ventilation weaning protocol: Two case reports

Peng M, Ren D, Liu YF, Meng X, Wu M, Chen RL, Yu BJ, Tao LC, Chen L, Lai ZQ

- 3314** Adult duodenal intussusception with horizontal adenoma: A rare case report

Wang KP, Jiang H, Kong C, Wang LZ, Wang GY, Mo JG, Jin C

- 3320** Isolated metachronous splenic multiple metastases after colon cancer surgery: A case report and literature review

Hu L, Zhu JY, Fang L, Yu XC, Yan ZL

- 3329** Imaging of hemorrhagic primary central nervous system lymphoma: A case report
Wu YW, Zheng J, Liu LL, Cai JH, Yuan H, Ye J
- 3334** Coexistence of ovarian serous papillary cystadenofibroma and type A insulin resistance syndrome in a 14-year-old girl: A case report
Yan FF, Huang BK, Chen YL, Zhuang YZ, You XY, Liu CQ, Li XJ
- 3341** Acute suppurative oesophagitis with fever and cough: A case report
Men CJ, Singh SK, Zhang GL, Wang Y, Liu CW
- 3349** Computed tomography, magnetic resonance imaging, and F-deoxyglucose positron emission computed tomography/computed tomography findings of alveolar soft part sarcoma with calcification in the thigh: A case report
Wu ZJ, Bian TT, Zhan XH, Dong C, Wang YL, Xu WJ
- 3355** COVID-19 with asthma: A case report
Liu AL, Xu N, Li AJ
- 3365** Total laparoscopic segmental gastrectomy for gastrointestinal stromal tumors: A case report
Ren YX, He M, Ye PC, Wei SJ
- 3372** Facial and bilateral lower extremity edema due to drug-drug interactions in a patient with hepatitis C virus infection and benign prostate hypertrophy: A case report
Li YP, Yang Y, Wang MQ, Zhang X, Wang WJ, Li M, Wu FP, Dang SS

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Dr. Romano is Professor of Medicine-Gastroenterology at the University of Campania “Luigi Vanvitelli” in Naples, Italy. Dr. Romano received his MD degree cum Laude at the University Federico II in Naples, Italy in 1980 and, after 4 year of Post-Graduate course, he became Specialist in Gastroenterology and Gastrointestinal Endoscopy. Dr. Romano’s research interest was on the cross-talk between H. pylori and gastric epithelial cells, and presently is mainly focused on H. pylori eradication therapy and on the role of nutraceuticals in gastrointestinal diseases. Dr. Romano is presently the Chief of the Endoscopy and Chronic Inflammatory Gastrointestinal Disorders Unit, and Teacher at the University of Campania “Luigi Vanvitelli” in Naples, Italy.

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases (WJCC, World J Clin Cases)* is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJCC* as 1.013; IF without journal self cites: 0.991; Ranking: 120 among 165 journals in medicine, general and internal; and Quartile category: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Electronic Editor: Yan-Xia Xing, Production Department Director: Yun-Xiaojuan Wu, Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Semimonthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

August 6, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Hypothyroidism with elevated pancreatic amylase and lipase without clinical symptoms: A case report

Yong-Wei Xu, Ran Li, Shu-Chang Xu

ORCID number: Yong-Wei Xu 0000-0002-7416-4281; Ran Li 0000-0003-2304-6282; Shu-Chang Xu 0000-0002-4841-462X.

Author contributions: Xu YW drafted the manuscript; Xu YW and Li R analyzed the data; Xu SC revised the article critically; all authors read and approved the final manuscript.

Supported by Shanghai Shenkang Hospital Development Center, No. SHDC12016109.

Informed consent statement: Written informed consent was obtained from the patient for publication of this case report.

Conflict-of-interest statement: No potential conflicts of interest relevant to this article were reported.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution

Yong-Wei Xu, Shu-Chang Xu, Department of Gastroenterology, Tongji Hospital, Tongji University School of Medicine, Shanghai 200065, China

Ran Li, Department of Endocrinology and Metabolism, Tongji Hospital, Tongji University School of Medicine, Shanghai 200065, China

Corresponding author: Shu-Chang Xu, PhD, MD, Chief Doctor, Professor, Department of Gastroenterology, Tongji Hospital, Tongji University School of Medicine, No. 389 Xincun Road, Putuo District, Shanghai 200065, China. xschang@163.com

Abstract

BACKGROUND

Hypothyroidism is an endocrine disorder that has worldwide prevalence and can affect multiple organ systems. We report a case of hypothyroidism with elevated pancreatic amylase and trypsin without acute pancreatitis. No such case has been previously reported.

CASE SUMMARY

A 29-year-old woman did not pay much attention to a fever 4 d prior. During this time, she experienced anorexia and only drank a small amount of water every day. She did not present with abdominal distension, postprandial nausea, vomiting, cough or expectoration. After physical and laboratory examinations, the patient was diagnosed with hypothyroidism. During the course of the disease, hypothyroidism was generally accompanied by constantly increased pancreatic amylase and trypsin. After admission, the possible etiology of the patient was excluded and the concentrations of pancreatic lipase and amylase in serum were > 2000U/L (reference range 23-300 U/L) and 410 U/L (reference range 30-110 U/L), respectively. So we highly suspected that it may be acute pancreatitis. Interestingly, she never developed any complications associated with acute pancreatitis despite high levels of serum pancreatic amylase and trypsin, and she reported no symptoms of abdominal pain. Serum amylase and lipase decreased gradually after active thyroxine supplementation, and the patient was discharged from the hospital after active treatment.

CONCLUSION

This case suggests that clinicians should pay attention to hypothyroidism with elevated pancreatic amylase and trypsin, even if no complications of acute pancreatitis are reported.

NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: March 21, 2020

Peer-review started: March 21, 2020

First decision: April 24, 2020

Revised: April 30, 2020

Accepted: July 15, 2020

Article in press: July 15, 2020

Published online: August 6, 2020

P-Reviewer: Eleazu C, Inal V

S-Editor: Ma YJ

L-Editor: Filipodia

E-Editor: Li JH



Key words: Hypothyroidism; Amylase; Trypsin; Acute pancreatitis; Clinical symptoms; Case report

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Hypothyroidism is an endocrine disorder that has worldwide prevalence and can affect multiple organ systems. However, even when pancreatic amylase and lipase are elevated, hypothyroidism without acute pancreatitis rarely causes clinical symptoms. Here, we report a case of hypothyroidism leading to elevated trypsin without acute pancreatitis in a patient who had no clinical symptoms. This case report can serve as a reminder to gastroenterologists, emergency physicians, and surgeons who encounter these cases in clinical practice.

Citation: Xu YW, Li R, Xu SC. Hypothyroidism with elevated pancreatic amylase and lipase without clinical symptoms: A case report. *World J Clin Cases* 2020; 8(15): 3299-3304

URL: <https://www.wjgnet.com/2307-8960/full/v8/i15/3299.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v8.i15.3299>

INTRODUCTION

Hypothyroidism is one of the most common endocrine disorders, affecting as much as 10% of the global population^[1]. Hypothyroidism is the result of an insufficient amount of circulating thyroid hormones; thus, it can affect multiple systems^[2]. In primary hypothyroidism, due to thyroid gland failure, serum thyroid hormone levels, including circulating levels of serum total thyroxine (T4) and free (FT4) thyroxine, total (T3) and free (FT3) triiodothyronine, decrease below the lower limit of the reference range, and pituitary thyrotropin (TSH) levels are elevated due to lack of feed-back inhibition the relationship of the hypothalamic-pituitary-thyroid axis. Therefore, the diagnosis of hypothyroidism is mainly based on serum parameters.

Thyroid hormones play an important role in the body and affect every organ system. The symptoms of hypothyroidism are not only nonspecific but also overlap with complaints of normal healthy people^[3]. The thyroid is an important endocrine organ in the body. The secretion of thyroid hormone can regulate the metabolism of sugar, protein, and fat in the human body; affect the growth and development of the body; and to some extent affect nervous, endocrine, cardiovascular and reproductive system functioning in the body. Hypothyroidism leads to decrease in the secretion of thyroid hormones, which loses the feedback effect of thyrotropin-releasing hormone (TRH) inhibition, causing thyroid epithelial cells to proliferate, leading to thyroid enlargement. This is a common clinical phenomenon. Of course, there are some rare phenomena. Hypothyroidism can be accompanied by pericardial effusion and pleural effusion in the cardiovascular system. It also leads to anorexia, abdominal distention of the digestive system, female menorrhagia, chronic amenorrhea, male impotence, and decreased libido. Hypothyroidism can also lead to elevated serum pancreatic amylase and trypsin without acute pancreatitis; however, there have been no previous reports on this phenomenon.

CASE PRESENTATION

Chief complaints

A 29-year-old woman was admitted to our hospital with anorexia, nausea, vomiting, apathy, and chills.

History of present illness

A young woman with anorexia, nausea, vomiting, apathy, chills, and fever drank plenty of water at home and did not take her symptoms seriously. However, her temperature did not decrease. Moreover, her fatigue was aggravated, and she started experiencing chest pain. Then the patient was admitted to our hospital.

History of past illness

She denied hepatitis, metabolic disorders, heart disease, cerebrovascular disease, mental illness, and immunodeficiency diseases.

Personal and family history

The patient was married and had two sons and one daughter. Her menstrual cycle was regular, the menstrual quantity was moderate, and the color was normal. No additional family history was obtained.

Physical examination upon admission

Physical examination after admission showed that the patient's body temperature was 39.8 °C, her respiratory rate was 20 breaths per min, and her blood pressure was 75/72 mmHg. The results of general physical examination were not good, and she was in poor spirits and expressionless. The patient's sclera were not stained yellow, and superficial lymph nodes were not enlarged. However, after a physical examination of the patient's thyroid, it was not an obvious goiter, and maybe only a Grade 1 goiter. This was because the patient's condition was short and the time frame was short, so the patient's goiter did not show. An abdominal examination showed that her abdomen was flat and soft, and there was no obvious tenderness in her abdomen.

Laboratory examinations

After admission to the hospital, the patient underwent a routine blood test (white blood cell count of $7.54 \times 10^9/L$, reference range $3.5 \times 10^9/L$ - $9.5 \times 10^9/L$; neutrophil count of $9.5 \times 10^9/L$, reference range $1.8 \times 10^9/L$ - $6.3 \times 10^9/L$; pancreatic amylase at 33 U/L, reference range 30-110 U/L and lipase at 363 U/L, reference range 23-300 U/L). The serum concentrations of TSH (0.09 mIU/L, reference range 0.3-5.5 mIU/L), total thyroxine (TT4 64.9 nmol/L, reference range 58.1-140.6 nmol/L, and free thyroxine FT4 10.06 pmol/L, reference range 11.5-22.7 pmol/L), and total triiodothyronine (TT3 0.63 nmol/L, reference range 0.92-2.79 nmol/L, and free FT3 1.04 pmol/L, reference range 0.92-2.79 pmol/L) were measured. The results of these tests were measured on the first day after the patient was admitted to the hospital. After treatment, the results were significantly improved (Table 1).

Imaging examinations

The patient was examined after admission, and a computed tomography (CT) scan of the chest showed no obvious abnormalities. The abdominal CT scan showed a small kidney stone in the left kidney, hepatic adipose infiltration and a small amount of ascites. Abdominal imaging indicated no acute pancreatitis. There was no significant difference between the results after treatment and the results before treatment (Figure 1).

FINAL DIAGNOSIS

The patient was diagnosed with septic shock, multiple organ dysfunction syndrome, and hypothyroidism.

TREATMENT

Based on the results of the examinations and after discussion with the treatment group, it was suggested that some measures be taken, including gastrointestinal decompression and rapid anti-inflammatory treatment with antibiotics. After active anti-inflammatory treatment, the inflammation indicators in the patient's blood, such as white blood cells and neutrophils, gradually decreased (Figure 2). At the same time, the patient was treated with thyroxine (75 µg) for hypothyroidism on the sixth day after admission, and the thyroid function was monitored. After supplementation with levothyroxine sodium tablets, pancreatic amylase and lipase also slowly decreased (Figure 3).

Table 1 Test results of thyroid hormone in serum

Parameter	Second day of admission	Sixth day of admission	tenth day of admission	Reference ranges
Total thyroxine, TT4	64.9	62.4	79.4	58.1-140.6 nmol/L
Free thyroxine, FT4	10.06	10.7	11.4	11.5-22.7 pmol/L
Total triiodothyronine, TT3	0.63	0.59	0.86	0.92-2.79 nmol/L
Free triiodothyronine, FT3	1.04	2.05	2.21	0.92-2.79 pmol/L
Thyrotropin, thyroid stimulating hormone, TSH	0.09	0.03	0.77	0.3-5.5 mIU/L




Figure 1 Abdominal computed tomography. A: Abdominal computed tomography (CT) examination with no manifestation of pancreatitis on admission; B: Results of abdominal CT examination after admission.




Figure 2 After the patient was admitted to the hospital, the changes in white blood cells and neutrophils in the blood were checked. White blood cell count of reference range $3.5 \times 10^9/L$ - $9.5 \times 10^9/L$; neutrophil cell count of reference range $1.8 \times 10^9/L$ - $6.3 \times 10^9/L$. X-axis: Days after admission; Y-axis: Units of measured value.

OUTCOME AND FOLLOW-UP

After active treatment, the patient’s symptoms gradually improved. The patient recovered and was discharged from the hospital after antibacterial treatment, supplementation with levothyroxine sodium tablets and nutritional support. The patient continued thyroxine supplementation after discharge from the hospital. After the patient was discharged, after re-examination on the eighteenth day, the serum amylase and lipase gradually became normal (Figure 3). No acute pancreatitis was observed in the patients during the course of the disease, not only lipase and amylase in the blood gradually increase, but also with it slowly decreased to normal ranges after the patient was discharged from the hospital.




Figure 3 After the patient was admitted to the hospital, the changes in amylase and middle lipase in the patient's blood were examined. Pancreatic amylase reference range 30-110 U/L and pancreatic lipase reference range 23-300 U/L. X-axis: Days after admission; Y-axis: Units of measured value.

DISCUSSION

Hypothyroidism is an endocrine disease with significant morbidity and worldwide prevalence, which can affect multiple organ systems. It can be asymptomatic and subclinical and has the potential to be complicated by fatal pathologies. Hypothyroidism, characterized by the absence of functional thyroid hormone, is diagnosed when decreased levels of thyroid hormones cause an increase in the secretion of TSH^[4]. Hypothyroidism, which occurs in not only the thyroid itself, can cause systemic clinical symptoms because thyroid hormone is dispersed throughout the whole body.

The clinical symptoms of hypothyroidism vary and can include clinical and subclinical symptoms. Clinical manifestations can vary based on the age of the patient at the time of diagnosis and the severity of hormone deficiency. Weight gain, constipation, cold sensitivity, fatigue, and dry skin are the most commonly reported symptoms of hypothyroidism^[5,6]. Hypothyroidism can also cause a variety of common diseases, including chronic nephritis, anemia, and edema in the serous cavity. Additionally, Mirwais *et al*^[7] reported that hypothyroidism caused pericardial effusion, and Rishi Ramtahal reported that subclinical hypothyroidism caused hypertension in pregnancy^[8], preeclampsia-like syndrome^[9] and paralytic ileus and acute kidney injuries^[10].

We began to suspect that the action of inflammatory factors caused pancreatic lipase and amylase to increase, however, after active control of inflammation, trypsin remained high and gradually increasing. Leukocytes and neutrophils were gradually reduced on the fourth day after admission, after active anti-inflammatory treatment (Figure 2). There were no clinical symptoms of acute pancreatitis, but after the control of inflammatory, amylase and lipase in the pancreas were still higher than normal. On the basis of the results of the examination, we considered that the increase in trypsin may be due to the decrease in thyroxine, maybe not due to the effect of inflammation. Therefore, we initiated thyroxine supplementation. On the fourth day after admission, the patient began to add levothyroxine on the sixth day, the thyroxine level in the patient began to gradually increase (Table 1). After with the thyroid hormone supplementation, pancreatic amylase eventually decreased (Figure 3).

Hypothyroidism has not been previously reported to lead to elevated pancreatic amylase and lipase without clinical symptoms. A previous study by Blanco-Molina A found that dissociated acini in rats with hypothyroidism showed higher bethanechol-induced amylase secretion than acini in control animals, and the experimental rates were also more sensitive to the inhibitory TRH effect than the controls^[11]. The animals with hypothyroidism were more sensitive to the inhibitory effect of TRH than the control animals. It was previously shown that TRH inhibited amylase pancreatic secretion by a direct effect on acinar preparations. When TRH was decreased, the inhibition of pancreatic amylase was lost. These findings suggest that TRH could be an inhibitor of pancreatic amylase secretion, resulting in elevated, amylase and lipase levels in hypothyroidism.

This case occurred in a patient admitted to the hospital with fever. During the course of the disease, amylase and lipase were elevated, but no acute pancreatitis was observed. Hypothyroidism was finally diagnosed considering the patient's medical history and the examination results. It was finally found that the thyroid function of

the patient was reduced, and serum amylase and lipase decreased gradually after thyroid hormone supplementation. The patient was finally discharged safely. According to the literature, the mechanism of this phenomenon is not clear. In the future, we will analyze additional cases to study its pathogenesis.

CONCLUSION

Hypothyroidism with elevated pancreatic amylase and lipase without clinical symptoms is a relatively rare occurrence in the clinical setting. In future clinical practice, we should pay attention to the complications of hypothyroidism. We expect that a detailed description of this case will provide valuable resources for future diagnosis. However, additional cases are needed to confirm our findings, and research is needed to further elucidate the mechanism.

REFERENCES

- 1 **Niazi AK**, Kalra S, Irfan A, Islam A. Thyroidology over the ages. *Indian J Endocrinol Metab* 2011; **15**: S121-S126 [PMID: 21966648 DOI: 10.4103/2230-8210.83347]
- 2 **Gomes-Lima C**, Wartofsky L, Burman K. Can Reverse T3 Assay Be Employed to Guide T4 vs. T4/T3 Therapy in Hypothyroidism? *Front Endocrinol (Lausanne)* 2019; **10**: 856 [PMID: 31920970 DOI: 10.3389/fendo.2019.00856]
- 3 **Biondi B**, Cooper DS. Thyroid hormone therapy for hypothyroidism. *Endocrine* 2019; **66**: 18-26 [PMID: 31372822 DOI: 10.1007/s12020-019-02023-7]
- 4 **Udovic M**, Pena RH, Patham B, Tabatabai L, Kansara A. Hypothyroidism and the Heart. *Methodist Debakey Cardiovasc J* 2017; **13**: 55-59 [PMID: 28740582 DOI: 10.14797/mdcj-13-2-55]
- 5 **Carlé A**, Pedersen IB, Knudsen N, Perrild H, Ovesen L, Laurberg P. Hypothyroid symptoms and the likelihood of overt thyroid failure: a population-based case-control study. *Eur J Endocrinol* 2014; **171**: 593-602 [PMID: 25305308 DOI: 10.1530/EJE-14-0481]
- 6 **Diaz A**, Lipman Diaz EG. Hypothyroidism. *Pediatr Rev* 2014; **35**: 336-47; quiz 348-9 [PMID: 25086165 DOI: 10.1542/pir.35-8-336]
- 7 **Mirwais S**, Kazmi SH, Hussain SI, Mirwais M, Sharma A. Hypothyroidism Causing Pericardial Effusion: A Case Report. *Cureus* 2019; **11**: e6393 [PMID: 31938670 DOI: 10.7759/cureus.6393]
- 8 **Ramtahal R**, Dhanoo A. Subclinical hypothyroidism causing hypertension in pregnancy. *J Am Soc Hypertens* 2016; **10**: 691-693 [PMID: 27506728 DOI: 10.1016/j.jash.2016.06.039]
- 9 **Inversetti A**, Serafini A, Manzoni MF, Dolcetta Capuzzo A, Valsecchi L, Candiani M. Severe hypothyroidism causing pre-eclampsia-like syndrome. *Case Rep Endocrinol* 2012; **2012**: 586056 [PMID: 22937297 DOI: 10.1155/2012/586056]
- 10 **Rodrigo C**, Gamakaranage CS, Epa DS, Gnanathanan A, Rajapakse S. Hypothyroidism causing paralytic ileus and acute kidney injury - case report. *Thyroid Res* 2011; **4**: 7 [PMID: 21303532 DOI: 10.1186/1756-6614-4-7]
- 11 **Blanco-Molina A**, Lopez Segura F, Vara E, Lopez Miranda J, Torre-Cisneros J, Perez-Jimenez F. Different secretory response of pancreatic isolated lobules and dissociated acini from hypothyroid rats to exogen TRH. *Exp Clin Endocrinol* 1991; **98**: 15-22 [PMID: 1718766 DOI: 10.1055/s-0029-1211095]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

