

2 Name of Journal: *World Journal of Orthopedics*

Manuscript NO: 55983

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Improved patient reported outcomes with functional articulating spacers in two-stage revision of the infected hip **3**

Ewout S Veltman, Dirk Jan F Moojen, Rudolf W Poolman

Match Overview

1	Crossref 76 words "EHS Congress 2018 - abstract book", HIP International, 2018	2%
2	Internet 51 words crawled on 30-Jan-2020 f6publishing.blob.core.windows.net	1%
3	Internet 16 words crawled on 30-Sep-2020 journals.sagepub.com	<1%
4	Internet 12 words crawled on 06-Jul-2020 www.hindawi.com	<1%



ALL

IMAGES

VIDEOS

27,400 Results

Any time ▾

Suppl 2: Revision in Cemented and Cementless Infected Hip ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3722547>

TWO-STAGE REVISION. Two-stage revision has been reported to produce the best results, with an infection eradication rate higher than 90% . The principles of two-stage revision are the removal of all prosthetic components, including cement, with radical debridement of infected tissue and bone.

Cited by: 9

Author: Paolo Cherubino, Marco Puricelli, Fabio ...

Publish Year: 2013

Outcome of single-stage versus two-stage exchange for ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6719605>

Aug 02, 2019 · The eradication rate was higher in the two-stage exchange group: 89.8% with 40 months of follow-up versus 81.9% with 44 months of follow-up in the single-stage exchange group. In cases of two-stage revision, they recommended using an articulating spacer to improve the eradication rate: 91.2% versus 87.0% in cases of static spacer.

Cited by: 8

Author: Corentin Pangaud, Matthieu Ollivier, Jean-...

Publish Year: 2019

PEOPLE ALSO ASK

What is the use of antibiotic loaded articulating spacers? ▾

Is total hip arthroplasty a serious complication? ▾

What is treatment for PJIs after shoulder arthroplasty? ▾

What antibiotic is used for a type 1 infection? ▾

Feedback

Two-Stage Revision Total Hip Arthroplasty With a Specific ...

<https://www.sciencedirect.com/science/article/pii/S0883540318307204>

Dec 01, 2018 · In their series of 99 hips treated with the same specific articulating spacer design for 2-stage hip exchange, Biring et al reported an 89% rate of long-term infection eradication.

Furthermore, in this cohort, only 3 patients (2%) required a 2-stage revision for PJI at a minimum of 2 years of follow-up after reimplantation.

国内版

国际版

Improved patient reported outcomes with functional articulating sp



ALL

IMAGES

VIDEOS

22,100 Results

Any time ▾

[Suppl 2: Revision in Cemented and Cementless Infected Hip ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3722547>

TWO-STAGE REVISION Two-stage revision has been reported to produce the best results, with an infection eradication rate higher than 90%. The principles of **two-stage revision** are the removal of all prosthetic components, including cement, with radical debridement of infected tissue and bone.

Cited by: 9

Author: Paolo Cherubino, Marco Puricelli, Fabio ...

Publish Year: 2013

[Two-Stage Revision Total Hip Arthroplasty With a Specific ...](#)

<https://www.sciencedirect.com/science/article/pii/S0883540318307204>

Dec 01, 2018 · **Two-stage revision** utilizing **spacers** loaded with **high-dose antibiotic cement** prior to reimplantation remains the gold standard for treatment of periprosthetic joint infections (PJI) in **total hip arthroplasty** (THA) in North America, but there is a paucity of data on **mid-term outcomes**.

Cited by: 9

Author: Brian P. Chalmers, Tad M. Mabry, Matthe...

Publish Year: 2018

[Are Preformed Articulating Spacers Superior To Surgeon ...](#)



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

39,200 Results

Any time ▾

[Suppl 2: Revision in Cemented and Cementless Infected Hip ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3722547>

TWO-STAGE REVISION **Two-stage revision** has been **reported** to produce the best results, with an infection eradication rate higher than 90%. The principles of **two-stage revision** are the removal of all prosthetic components, including cement, with radical debridement of infected tissue and bone.

Cited by: 9**Author:** Paolo Cherubino, Marco Puricelli, Fabio ...**Publish Year:** 2013

[Two-Stage Revision Arthroplasty for Periprosthetic Hip ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4429212>

The **two-stage revision hip arthroplasty protocol** offers the greatest chance for eradication of infection, with **reported success rates** of greater than 90% in various studies [5, 6]. Although it has been the gold standard for the treatment of **patients** with chronic PJI, few studies have **reported outcomes** beyond short-term **follow-up**.

Cited by: 18**Author:** Szu-Yuan Chen, Chi-Chien Hu, Chun-Chi...**Publish Year:** 2015

[Two-Stage Revision Total Hip Arthroplasty With a Specific ...](#)

<https://www.sciencedirect.com/science/article/pii/S0883540318307204>

Dec 01, 2018 · **Two-stage revision** utilizing **spacers** loaded with **high-dose antibiotic cement** prior to reimplantation remains the gold standard for treatment of periprosthetic joint infections (PJI) in **total hip arthroplasty** (THA) in North America, but there is a paucity of data on **mid-term outcomes**.

Cited by: 9**Author:** Brian P. Chalmers, Tad M. Mabry, Matthe...**Publish Year:** 2018

[\(PDF\) An Articulating Antibiotic Spacer Controls Infection ...](#)

<https://www.researchgate.net/publication/51076875...>

Apr 26, 2011 · **Two-stage** surgery with **revision total hip arthroplasty** and **spacer** implantation [10,11] has been **reported** to result in a high rate of infection control but also a ...