



PEER-REVIEW REPORT

Name of journal: World Journal of Clinical Cases

Manuscript NO: 56702

Title: Retroperitoneal vs transperitoneal laparoscopic lithotripsy of 20-40 mm renal stones within horseshoe kidneys

Reviewer's code: 02507819

Position: Editorial Board

Academic degree: FRCS (Gen Surg), MBBS, MD, MNAMS

Professional title: Associate Professor

Reviewer's Country/Territory: India

Author's Country/Territory: China

Manuscript submission date: 2020-05-21

Reviewer chosen by: AI Technique

Reviewer accepted review: 2020-05-23 15:24

Reviewer performed review: 2020-05-28 04:22

Review time: 4 Days and 12 Hours

Scientific quality	<input type="checkbox"/> Grade A: Excellent <input checked="" type="checkbox"/> Grade B: Very good <input type="checkbox"/> Grade C: Good <input type="checkbox"/> Grade D: Fair <input type="checkbox"/> Grade E: Do not publish
Language quality	<input checked="" type="checkbox"/> Grade A: Priority publishing <input type="checkbox"/> Grade B: Minor language polishing <input type="checkbox"/> Grade C: A great deal of language polishing <input type="checkbox"/> Grade D: Rejection
Conclusion	<input type="checkbox"/> Accept (High priority) <input checked="" type="checkbox"/> Accept (General priority) <input type="checkbox"/> Minor revision <input type="checkbox"/> Major revision <input type="checkbox"/> Rejection
Re-review	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
Peer-reviewer statements	Peer-Review: <input checked="" type="checkbox"/> Anonymous <input type="checkbox"/> Onymous Conflicts-of-Interest: <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No



**Baishideng
Publishing
Group**

7041 Koll Center Parkway, Suite
160, Pleasanton, CA 94566, USA
Telephone: +1-925-399-1568
E-mail: bpgoffice@wjgnet.com
https://www.wjgnet.com

SPECIFIC COMMENTS TO AUTHORS

This retrospective study compares retroperitoneal and transperitoneal approaches of laparoscopic pyelolithotomy in horse shoe kidney. According to this study, there is no difference in two approaches other than mean post-operative fasting time. I want to know whether authors encountered any intra-renal pelvis and if yes, what were the problems faced and how did they manage? What was the size of double J ureteral stent used?