

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 56892

Manuscript Type: CASE REPORT

Uterine rupture in patients with a history of multiple curettages: Report of two cases

Uterine rupture in patients with a history of multiple curettages

Abstract

BACKGROUND

Uterine rupture is a serious obstetric emergency, a severe event, and a serious threat to maternal and foetal life. It is a rare and not well characterized by the fact that multiple operations of uterine cavities contribute to uterine rupture during pregnancy. Atypical uterine rupture is easily misdiagnosed as other obstetric or surgical diseases. In current guidelines, abdominocentesis is a contraindication for late pregnancy. Therefore, these cases provide new ideas for clinical diagnosis and treatment of uterine rupture.

CASE SUMMARY

Case 1

A 34-year-old woman (gravida 5, para 2), 32 wk+4 days of gestation, presented with acute upper abdominal pain for 8 h with nausea and vomiting. CT revealed pelvic and abdominal effusion. We extracted 3 mL of unclotted blood from her abdominal cavity. An emergency caesarean section was performed. A uterine rupture was found, and the

Match Overview

- | Rank | Source | Words | Similarity |
|------|--|-----------|------------|
| 1 | Crossref | 122 words | 4% |
| | Yan Sun, Jing Huang, Hong-Fang Kong. "Spontaneous rupture of unscarred uterus in the third trimester after in vitro fertilization ..." | | |
| 2 | Crossref | 24 words | 1% |
| | Shu-Han You, Yao-Lung Chang, Chih-Feng Yen. "Rupture of the scarred and unscarred gravid uterus: Outcomes and risk factors ..." | | |
| 3 | Crossref | 22 words | 1% |
| | Meher B Ali, Mashal B Ali. "Late Presentation of Uterine Rupture: A Case Report", Cureus, 2019 | | |
| 4 | Internet | 20 words | 1% |
| | crawled on 24-Jun-2020
journals.lww.com | | |



Uterine rupture in patients with a history of multiple curettages: Report o

ALL

IMAGES

VIDEOS

18,500 Results

Any time ▾

Uterine rupture revisited: Predisposing factors, clinical ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3809304>

Methodology: A 14-year retrospective analysis of 61 gravid (>20 weeks of gestation) **uterine rupture** cases between January 1998 to March 2012 was carried out. Results: The incidence of ruptured uteri was calculated to be 0.116%. Persistence for vaginal delivery after cesarean was the most common cause of **uterine rupture** (31.1%).

Cited by: 10

Author: Abdulkadir Turgut, Ali Ozler, Mehmet Sid...

Publish Year: 2013

(PDF) Spontaneous uterine rupture: Report of two cases

<https://www.researchgate.net/publication/223984703...>

Uterine rupture is a rare entity with an incidence of 0.07% and is a cause of perinatal mortality. Dehiscence of previous **uterine** scar is the most important cause, but other causes have been ...

Uterine Rupture: A Seven Year Review at a Tertiary Care ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4746953>

Mar 26, 2014 · The incidence of **rupture** was 0.318% in women with **history** of prior **uterine** surgery and was 0.02% in women without **history** of any prior **uterine** surgery. Among the cases with **history** of prior **uterine** surgery, the vast majority of **patients** had prior low transverse cesarean section (84.8%). There was only one **patient** with prior hysterotomy.

Cited by: 7

Author: Maruti Sinha, Ridhima Gupta, Pushpende...

Publish Year: 2016

PEOPLE ALSO ASK

What is the risk of uterine rupture? ▾

Can uterine rupture be predicted? ▾

What are the symptoms of uterine pain? ▾

Feedback



Uterine rupture in patients with a history of multiple curettages: Rep



ALL

IMAGES

VIDEOS

37,700 Results

Any time ▼

(PDF) Spontaneous uterine rupture: Report of two cases

<https://www.researchgate.net/publication/223984703...>

Uterine rupture is a rare entity with an incidence of 0.07% and is a cause of perinatal mortality. Dehiscence of previous **uterine** scar is the most important cause, but other causes have been ...

An unusual presentation of recurrent uterine rupture ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4469860>

Uterine rupture is a rare but hazardous obstetric complication. High levels of caution should be exercised in patients with a history of prior **uterine rupture**, as they may present with atypical symptoms. Ultrasonography could provide valuable information in such **cases** where there is an elevated risk of **uterine rupture** at the previous **rupture** site.

Cited by: 1

Author: Shu Qi Tan, Edwin Wee Hong Thia, Chee Se...

Publish Year: 2015

Uterine rupture revisited: Predisposing factors, clinical ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3809304>

Methodology: A 14-year retrospective analysis of 61 gravid (>20 weeks of gestation) **uterine rupture cases** between January 1998 to March 2012 was carried out. Results: The incidence of ruptured uteri was calculated to be 0.116%. Persistence for vaginal delivery after cesarean was the most common cause of **uterine rupture** (31.1%).

Cited by: 10

Author: Abdulkadir Turgut, Ali Ozler, Mehmet Siddik...

Publish Year: 2013

PEOPLE ALSO ASK

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

42,000 Results

Any time ▼

Case Report: Silent uterine rupture in scarred uterus

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5747674>

Case presentation. **Uterine rupture** is an uncommon but one of the most serious obstetric complications, with an increased risk of maternal and perinatal morbidity, and mortality. 1 **Uterine rupture** is reported in 0.5%–1% of **patients** attempting vaginal births after caesarean section. **Uterine rupture** occurs when a full-thickness disruption of the **uterine** wall that also involves the **uterine** ...

Author: Ream Langhe, Umme Farwa Shah, At... **Publish Year:** 2017

Case Report: An unusual cause for epigastric pain in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3948010>

Mar 05, 2014 · The incidence of **uterine rupture** as a result of previous perforation is unclear with little published data and few **case reports**. Cases of **uterine rupture** after perforation following hysteroscopic resection of fibroids, **uterine** septum are well published but the authors found no known previous cases related to laparoscopy.

Cited by: 2 **Author:** Asmaa Al-Kufaishi, Kim Erasmus, David ...

Publish Year: 2014

Uterine Rupture after Laparoscopic Myomectomy in Two Cases ...

<https://www.hindawi.com/journals/crig/2017/1404815> ▼

We describe **two** cases of **uterine rupture** in pregnancy after laparoscopic myomectomy and analyze all the aetiological factors involved in this circumstance according to the recent literature, focusing above all on the surgical procedures and the characteristics of the excised myomas. The **two** cases of **uterine rupture** in pregnancy following laparoscopic myomectomy occurred at 36 and 18 weeks of ...

Cited by: 3 **Author:** Antonella Vimercati, Vittoria Del Vecchio, ...

Publish Year: 2017