

Kyoto classification in patients who developed multiple gastric carcinomas after *Helicobacter pylori* eradication

Sakitani K *et al.* Kyoto classification in gastric carcinomas

Kosuke Sakitani, Toshihiro Nishizawa, Akira Toyoshima, Shuntaro Yoshida, Tatsuya Matsuno, Tomoharu Yamada, Masatoshi Irokawa, Yoshiyuki Takahashi, Yousuke Nakai, Osamu Toyoshima, Kazuhiko Koike

Kosuke Sakitani, Shuntaro Yoshida, Tatsuya Matsuno, Masatoshi Irokawa, Yoshiyuki Takahashi, Osamu Toyoshima, Department of Gastroenterology, Toyoshima Endoscopy Clinic, Tokyo 157-0066, Japan

Toshihiro Nishizawa, Department of Gastroenterology, International University of Health and Welfare, Narita Hospital, Narita 2868520, Chiba, Japan

Akira Toyoshima, Department of Colorectal Surgery, Japanese Red Cross Medical Center, Tokyo 1508935, Japan

Match Overview

1	Crossref 25 words Kosuke Sakitani, Toshihiro Nishizawa, Masahide Arita, Shu... aro Yoshida et al. " Early detection of gastric cancer after era	1%
2	Internet 13 words crawled on 01-Aug-2020 atm.amegroups.com	<1%
3	Internet 13 words crawled on 21-Jul-2020 www.ndafp.org	<1%
4	Internet 12 words crawled on 06-Jul-2020 www.thieme-connect.com	<1%

Kyoto classification in patients who developed multiple gastric car



ALL

IMAGES

VIDEOS

30,900 Results

Any time ▾

Helicobacter pylori eradication improved the Kyoto ...

<https://onlinelibrary.wiley.com/doi/full/10.1002/jgh3.12360>

Abstract Background and Aim Endoscopy-based **Kyoto classification** predicts the risk of Helicobacter **pylori** infection and **gastric** cancer; however, the change in score following **H. pylori eradication** ...

Author: Osamu Toyoshima, Toshihiro Nishiza... **Publish Year:** 2020

Kyoto global consensus report on Helicobacter pylori ...

<https://gut.bmj.com/content/64/9/1353> ▾

Sep 01, 2015 · Objective To present results of the **Kyoto** Global Consensus Meeting, which was convened to develop global consensus on (1) **classification** of chronic gastritis and duodenitis, (2) clinical distinction of dyspepsia caused by Helicobacter **pylori** from functional dyspepsia, (3) appropriate diagnostic assessment of gastritis and (4) when, whom and how to treat **H. pylori** ...

Cited by: 971 **Author:** Kentaro Sugano, Jan Tack, Ernst J Kuip...

Publish Year: 2015

Endoscopic Kyoto classification of Helicobacter pylori ...

https://www.researchgate.net/publication/339105073_Endoscopic_Kyoto_classification_of...

A Kyoto classification score of 2 or more indicates H. **pylori** infection. Kyoto classification scores of patients with and without gastric cancer were 4.8 and 3.8, respectively. A Kyoto...

Can image-enhanced endoscopy improve the diagnosis of ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/den.13540>

Sep 24, 2019 · Recently, the Kyoto classification of gastritis was developed based on the endoscopic characteristics of H. **pylori** infection-associated gastritis to diagnose H. pylori infection accurately and identify the risk factors of gastric carcinoma. 9 This classification represents 19 endoscopic findings that divides patients into three groups: patients without H. pylori infection ...



32,100 Results Any time ▾

Endoscopic Kyoto classification of Helicobacter pylori ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7015719>

Feb 07, 2020 · The Kyoto classification organized endoscopic findings related to H. pylori infection. A Kyoto classification score ≥ 2 indicates H. pylori infection. An H. pylori test is essential for such cases with no history of H. pylori eradication. A Kyoto classification score ≥ 4 might indicate gastric cancer risk. Such cases need careful follow-up.

Cited by: 3 **Author:** Osamu Toyoshima, Toshihiro Nishizawa, K...

Publish Year: 2020

Kyoto global consensus report on Helicobacter pylori ...

<https://gut.bmj.com/content/64/9/1353> ▾

Sep 01, 2015 · Objective To present results of the **Kyoto** Global Consensus Meeting, which was convened to develop global consensus on (1) **classification** of chronic gastritis and duodenitis, (2) clinical distinction of dyspepsia caused by **Helicobacter pylori** from functional dyspepsia, (3) appropriate diagnostic assessment of gastritis and (4) when, whom and how to treat H. **pylori** gastritis.

Cited by: 988 **Author:** Kentaro Sugano, Jan Tack, Ernst J Kuipers, ...

Publish Year: 2015

Endoscopic Kyoto classification of Helicobacter pylori ...

<https://www.researchgate.net/publication/339105073...>

A **Kyoto classification** score of **0** indicates **no H. pylori** infection. A Kyoto classification **score** of **2** or more indicates **H. pylori** infection. Kyoto classification scores of patients with and without...

(PDF) Efficacy of the Kyoto Classification of Gastritis in ...

Search Tools

Turn off Hover Translation (关闭取词)



Kyoto classification in patients who developed multiple gastric carci



ALL IMAGES VIDEOS MAPS NEWS SHOPPING

31,800 Results Any time ▾

Endoscopic Kyoto classification of Helicobacter pylori ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7015719>

Feb 07, 2020 · The Kyoto classification organized endoscopic findings related to H. pylori infection. A Kyoto classification score ≥ 2 indicates H. pylori infection. An H. pylori test is essential for such cases with no history of H. pylori eradication. A Kyoto classification score ≥ 4 might indicate gastric cancer risk. Such cases need careful follow-up.

Cited by: 3 **Author:** Osamu Toyoshima, Toshihiro Nishizawa, ...

Publish Year: 2020

Kyoto global consensus report on Helicobacter pylori ...

<https://gut.bmj.com/content/64/9/1353> ▾

Sep 01, 2015 · Objective To present results of the **Kyoto** Global Consensus Meeting, which was convened to develop global consensus on (1) **classification** of chronic gastritis and duodenitis, (2) clinical distinction of dyspepsia caused by **Helicobacter pylori** from functional dyspepsia, (3) appropriate diagnostic assessment of gastritis and (4) when, whom and how to treat H. **pylori** gastritis.

Cited by: 988 **Author:** Kentaro Sugano, Jan Tack, Ernst J Kuipe...

Publish Year: 2015

Endoscopic Kyoto classification of Helicobacter pylori ...

<https://www.researchgate.net/publication/339105073...>

A **Kyoto classification** score of 0 indicates no H. pylori infection. A Kyoto classification score of 2 or more indicates H. pylori infection. Kyoto classification scores of patients with and without...

(PDF) Efficacy of the Kyoto Classification of Gastritis in ...

https://www.researchgate.net/publication/315318697_Efficacy_of_the_Kyoto...

Objective The Kyoto gastritis classification categorizes the **endoscopic** characteristics of Helicobacter pylori (H. pylori) infection-associated gastritis and identifies patterns associated with a...

Epstein–Barr virus-associated gastric carcinomas developed ...

<https://link.springer.com/article/10.1007/s12328-020-01094-8> ▾

Jan 22, 2020 · Epstein–Barr virus-associated **gastric carcinoma** (EBVaGC) can develop in **gastric**

global.bing.com/?FORM=Z9FD1 ed by **Helicobacter pylori** (H. **pylori**)-induced atrophic gastritis. However, it is unclear

Helicobacter pylori

Helicobacter pylori, previously known as Campylobacter pylori, is a gram-negative, helically-shaped, microaerophilic bacterium usually found in the stomach. Its helical shape is thought to have evolved in order to penetrate the mucoid lining of the stomach and thereby establish infection. The bacterium was first identified in 1982 by Australian doctors Barry Marshall and Robin Warren, who found that it was present in a person with chronic gastritis and gastric ulcers, conditions not previously believed to have a microbial cause. H. pylori has been associated with the mucosa-associated lymphoid tissue in the stomach, esophagus, colon, rectum, or tissues around the eye, and of lymphoid tissue in the stomach.

Wikipedia

People also search for

Gastritis

Stomach Cancer

Irritable Bowel Syndrome

Epstein–Barr virus

Typhoid

See more ▾

Data from: Wikipedia

Suggest an edit