



Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 57709

Manuscript Type: META-ANALYSIS

Diagnostic efficacy of Japan NBI Expert Team and pit pattern classifications for colorectal lesions: A meta-analysis

Zhang Y *et al.* JNET and Pit pattern classifications

Yu Zhang, Hui-Yan Chen, Xiao-Lu Zhou, Wen-Sheng Pan, Xin-Xin Zhou, Hang-Hai Pan

Match Overview

1	Internet 206 words crawled on 08-Aug-2020 www.wjgnet.com	5%
2	Internet 69 words crawled on 28-Nov-2016 pdfs.semanticscholar.org	2%
3	Internet 62 words www.ncbi.nlm.nih.gov	1%
4	Crossref 60 words Ming Li. "Kudo's pit pattern classification for colorectal n ... oplasms: A meta-analysis", World Journal of Gastroentero	1%
5	Internet 43 words crawled on 14-Mar-2020 worldwidescience.org	1%
6	Internet 41 words crawled on 30-Jul-2019 ir.lib.hiroshima-u.ac.jp	1%

Crossref 41 words

激活 Windows

转到“设置”以激活 Windows。 Text-Only Report

Diagnostic efficacy of Japan NBI Expert Team (JNET) and Pit pattern



ALL

IMAGES

VIDEOS

5,490 Results

Any time ▾

[Diagnostic performance of Japan NBI Expert Team ...](#)

<https://www.sciencedirect.com/science/article/pii/S0016510717301207>

Oct 01, 2017 · The **Japan NBI Expert Team (JNET)** classification is the first universal **narrow-band imaging** magnifying endoscopic classification of **colorectal** tumors. Considering each type in this classification, the **diagnostic** ability of Type 2B is the weakest. Generally, clinical behavior is believed to be different in each gross type of **colorectal** tumor.

Cited by: 16**Author:** Kyoku Sumimoto, Shinji Tanaka, Kenjiro...**Publish Year:** 2017

[Diagnostic yield of the Japan NBI Expert Team \(JNET ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6683640>

Apr 26, 2019 · Magnifying **NBI**, in particular, is a reliable method for differentiating neoplastic from non-neoplastic **lesions**. 4 Several magnifying **NBI classifications** for optical diagnoses of neoplastic and non-neoplastic **lesions** have been developed. 5 –10 Recently, a unified magnifying **NBI** classification called the **Japan NBI Expert Team (JNET)** ...

Cited by: 1**Author:** Shunsuke Kobayashi, Masayoshi Yama...**Publish Year:** 2019

[Diagnostic yield of the Japan NBI Expert Team \(JNET ...](#)

<https://journals.sagepub.com/doi/abs/10.1177/2050640619845987>

Apr 26, 2019 · Magnifying **Narrow Band Imaging (NBI)** during colonoscopy is a reliable method for differential and depth diagnoses of **colorectal lesions**. This study examined the **diagnostic** yield of magnifying **NBI** based on the **Japan NBI Expert Team (JNET)** classification in a clinical setting using a large-scale clinical practice database.

Cited by: 1**Author:** Shunsuke Kobayashi, Masayoshi Yama...**Publish Year:** 2019



国内版

国际版

Diagnostic efficacy of Japan NBI Expert Team and pit pattern classi



Sign in



ALL

IMAGES

VIDEOS



Get the new Microsoft Edge >

12,300 Results

Any time ▾

Diagnostic performance of Japan NBI Expert Team ...

<https://www.sciencedirect.com/science/article/pii/S0016510717301207>

Oct 01, 2017 · The **Japan NBI Expert Team** (JNET) classification is the first universal **narrow-band imaging** magnifying endoscopic classification of **colorectal** tumors. Considering each type in this classification, the **diagnostic** ability of Type 2B is the weakest. Generally, clinical behavior is believed to be different in each gross type of **colorectal** tumor.

Cited by: 16

Author: Kyoku Sumimoto, Shinji Tanaka, Kenjiro Shi...

Publish Year: 2017

Search Tools

Turn off Hover Translation (关闭取词)

Diagnostic yield of the Japan NBI Expert Team (JNET ...

<https://journals.sagepub.com/doi/full/10.1177/2050640619845987>

Apr 26, 2019 · Magnifying **Narrow Band Imaging** (NBI) during colonoscopy is a reliable method for differential and depth diagnoses of **colorectal lesions**. This study examined the **diagnostic** yield of magnifying **NBI** based on the **Japan NBI Expert Team** (JNET) classification in a clinical setting using a large-scale clinical practice database.

Cited by: 1

Author: Shunsuke Kobayashi, Masayoshi Yamada, ...

Publish Year: 2019

Diagnostic yield of the Japan NBI Expert Team (JNET ...

激活 Windows

转到“设置”以激活 Windows。



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

25,200 Results

Any time ▼

Diagnostic yield of the Japan NBI Expert Team (JNET ...

<https://journals.sagepub.com/doi/full/10.1177/2050640619845987>

Apr 26, 2019 · Magnifying **Narrow Band Imaging** (NBI) during colonoscopy is a reliable method for differential and depth diagnoses of **colorectal lesions**. This study examined the **diagnostic** yield of magnifying NBI based on the **Japan NBI Expert Team** (JNET) classification in a clinical setting using a large-scale clinical practice database.

Cited by: 2**Author:** Shunsuke Kobayashi, Masayoshi Yamad...**Publish Year:** 2019

Estimation of Invasion Depth: The First Key to Successful ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6453840>

Mar 27, 2019 · Strategy for the endoscopic treatment of **colorectal lesions** according to the **Japan Narrow-band Imaging** (NBI) **Expert Team** (JNET) classification. The dotted arrows suggest that magnifying chromoendoscopy for **pit pattern** diagnosis can be performed in selected cases of JNET 2A or 3 with low confidence.

Cited by: 3**Author:** Bo In Lee, Takahisa Matsuda**Publish Year:** 2019

Narrow-band imaging (NBI) magnifying endoscopic ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/den.12644>

Introduction. **Narrow-band imaging** (NBI) was initially developed by Sano et al . under the supervision of Dr Shigeaki Yoshida at the National Cancer Center Hospital East in 1999. 1, 2 A prototype short-wavelength narrow-band red/green/blue (RGB) filter was successfully created in 2001 (monochrome NBI), 1, 2 and the microvascular architecture of the gastrointestinal tract and tumor ...

Cited by: 137**Author:** Yasushi Sano, Shinji Tanaka, Shin-ei Kud...**Publish Year:** 2016

Narrow-band imaging (NBI) magnifying endoscopic ...

<https://onlinelibrary.wiley.com/doi/10.1111/den.12644>

Introduction. **Narrow-band imaging** (NBI) was initially developed by Sano et al. under the supervision of Dr Shigeaki Yoshida at the National Cancer Center Hospital East in 1999. 1, 2 A prototype short-wavelength narrow-band red/green/blue (RGB) filter was successfully created in 2001 (monochrome NBI), 1, 2 and the microvascular architecture of the gastrointestinal tract and tumor ...