

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 58462

Manuscript Type: CASE REPORT

**Multiple ectopic goiter in the retroperitoneum, abdominal wall, liver, and diaphragm
a case report and literature review**

Two-year follow-up of Multiple ectopic goiter

Lan-hui Qin, Feng-yi He, Jin-yuan Liao

Abstract

BACKGROUND

Ectopic thyroid is a rare developmental disorder, typically found in lingual areas, and most distantly in the abdomen. Thyroid ectopia in multiple regions is extremely rare. To date, there are no reports of ectopic goiter in four regions of the abdominal cavity in a single patient.

CASE SUMMARY

We present a case of thyroid ectopia in four areas of the abdomen, comprising normal orthotopic thyroid tissue. A 36-year-old woman, who underwent ovarian teratoma resection 10 years previously due to symptomatic ovarian teratoma, was referred to our hospital for treatment of an incidental hepatic mass. The magnetic resonance imaging

Match Overview

1 **Internet** 13 words
crawled on 13-Sep-2020
www.wjgnet.com

1%



ALL

IMAGES

VIDEOS

29,500 Results

Any time ▾

Two cases of ectopic liver attached to the gallbladder wall

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2020551>

Introduction. Ectopic liver is a rare developmental anomaly in which liver tissue is situated outside the liver and has no hepatic connection 1.Ectopic liver tissue can occur in several different organs such as the diaphragm, hepatic ligaments, omentum, stomach, retroperitoneum and thorax 2,3, but the gallbladder is the commonest site of origin.

Cited by: 19

Author: John Giniatsos, Ali A Riaz, Alberto M Isla

Publish Year: 2002

Abdominal Multiple Splenosis Mimicking Liver and Colon ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3283864>

Here we report a case of multiple splenosis in abdominal cavity including liver and colon sites that was not reported in published studies, and review related literatures. Case Report A 49-year-old female patient complained of subxiphoid pain with fatigue for half a month.

Cited by: 20

Author: Yahui Liu, Bai Ji, Guangyi Wang, Yingch...

Publish Year: 2012

Challenge in preoperative diagnosis of retroperitoneal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4694507>

Ectopic locations other than thorax were reported, such as diaphragm [5,6], esophagus [7,8], stomach , ileal mesentery , neck [11,12], abdominal wall , periscapular area , and so on. Most cysts measured <5 cm in diameter exhibit no symptoms, which are usually related to secondary infection or compression of adjacent structures.

Author: Zhang D, Zhang Y, Liu X, Zhu J, Feng ...

Publish Year: 2015

PEOPLE ALSO ASK

Can gallstones cause ectopic liver? ▾

Can ectopic liver tissue attach to gallbladder? ▾

What is ectopic liver? ▾

Multiple ectopic goiter in the retroperitoneum, abdominal wall, liver,



ALL

IMAGES

VIDEOS

56,200 Results

Any time ▾

Two cases of ectopic liver attached to the gallbladder wall

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2020551>

Introduction. **Ectopic liver** is a rare developmental anomaly in which **liver** tissue is situated outside the **liver** and has no hepatic connection 1. **Ectopic liver** tissue can occur in several different organs such as the **diaphragm**, hepatic ligaments, omentum, stomach, **retroperitoneum** and thorax 2,3, but the gallbladder is the commonest site of origin.

Cited by: 19

Author: John Griniatsos, Ali A Riaz, Alberto M Isla

Publish Year: 2002

Challenge in preoperative diagnosis of retroperitoneal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4694507>

Ectopic locations other than thorax were reported, such as **diaphragm** [5,6], esophagus [7,8], stomach , ileal mesentery , neck [11,12], **abdominal wall** , periscapular area , and so on. Most cysts measured <5 cm in diameter exhibit no symptoms, which are usually related to secondary infection or compression of adjacent structures.

Author: Zhang D, Zhang Y, Liu X, Zhu J, Feng C, ...

Publish Year: 2015

Hemoperitoneum in a Cirrhotic Patient Due to Rupture of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2673472>

Apr 23, 2009 · The **liver** was cirrhotic; an exuberant collateral circulation was seen on the **retroperitoneum** and on the **abdominal wall**, with canalization of the Umbilical vein, with an 8 mm diameter. The Retzius vein was identified in both paracolic gutters, with a lot of collaterals. A **liver** ...

Cited by: 20

Author: Igor Rafael Sincos, Grace Mulatti, Sheila M...

Publish Year: 2009



Multiple ectopic goiter in the retroperitoneum, abdominal wall, liver



ALL IMAGES VIDEOS MAPS NEWS SHOPPING

54,800 Results Any time ▾

Intrahepatic and intra-abdominal splenosis: A case report ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6960294>

Dec 27, 2019 · Liu Y, Ji B, Wang G, Wang Y. **Abdominal multiple splenosis mimicking liver and colon tumors: a case report and review of the literature.** Int J ...

Author: Kiruthika Ananthan, Gibran Timothy Y... **Publish Year:** 2019

Challenge in preoperative diagnosis of retroperitoneal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4694507>

Ectopic locations other than thorax were reported, such as **diaphragm** [5,6], esophagus [7,8], stomach , ileal mesentery , neck [11,12], **abdominal wall** , periscapular area , and so on. Most cysts measured <5 cm in diameter exhibit no symptoms, which are usually related to secondary infection or compression of adjacent structures.

Author: Zhang D, Zhang Y, Liu X, Zhu J, Feng... **Publish Year:** 2015

Two cases of ectopic liver attached to the gallbladder wall

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2020551>

Introduction. **Ectopic liver** is a rare developmental anomaly in which **liver** tissue is situated outside the **liver** and has no hepatic connection 1.**Ectopic liver** tissue can occur in several different organs such as the **diaphragm**, hepatic ligaments, omentum, stomach, **retroperitoneum** and thorax 2,3, but the gallbladder is the commonest site of origin.

Cited by: 19 **Author:** John Griniatsos, Ali A Riaz, Alberto M Isla

Publish Year: 2002

[PDF] Case Report Abdominal Multiple Splenosis Mimicking Liver ...

citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.271.6207&rep=rep1&type=pdf

Case Report Abdominal Multiple Splenosis Mimicking Liver and Colon Tumors: A Case Report and Review of the Literature Yahui Liu *, Bai Ji , Guangyi Wang, Yingchao Wang Department of Hepatobiliary and Pancreatic Surgery, the First Bethune Hospital, Jilin University, Jilin 130021, China. * Yahui Liu and Bai Ji contribute equally to this **case report**.

A case report of abdominal splenosis - a practical mini ...