



58557_Auto_Edited.docx

Quotes Excluded
Bibliography Excluded2%
SIMILAR**Name of Journal:** *World Journal of Gastrointestinal Oncology***Manuscript NO:** 58557**Manuscript Type:** ORIGINAL ARTICLE*Retrospective Study***Predictive factors for early distant metastasis after neoadjuvant chemoradiotherapy
in locally advanced rectal cancer**

Predictive factors for early distant metastasis

Hyojung Park

Match Overview

1	Internet 14 words crawled on 28-Dec-2020 atm.amegroups.com	<1%
2	Crossref 14 words Young Chul Kim, Jai Keun Kim, Myeong-Jin Kim, Jei Hee Lee, Young Bae Kim, Sung Jae Shin. "Feasibility of mesorectal	<1%
3	Crossref 13 words Francesco Sciafani, David Cunningham. "Neoadjuvant chemotherapy without radiotherapy for locally advanced rectal ca	<1%
4	Internet 12 words crawled on 25-Dec-2019 www.mdpi.com	<1%
5	Internet 12 words crawled on 10-Sep-2011 www.dccg.nl	<1%

激活 Windows

转到“设置”以激活 Windows

Text-Only Report

Predictive factors for early distant metastasis after neoadjuvant ch



ALL

IMAGES

VIDEOS

79,900 Results

Any time ▼

Pathological complete response may underestimate distant ...

<https://pubmed.ncbi.nlm.nih.gov/30879932>

Aim: To evaluate the pattern of tumor relapse of pathological complete response (pCR) patients with **locally advanced rectal cancer** (LARC) following **neoadjuvant chemoradiotherapy** (nCRT) and total mesorectal excision (TME), and to identify **predictive factors** of **distant metastasis** in pCR patients **after** nCRT. Method: This was a retrospective analysis of 118 LARC patients who ...

Predictors of Early Distant Relapse in Rectal Cancer ...

<https://www.ncbi.nlm.nih.gov/pubmed/32036373>

Feb 07, 2020 · BACKGROUND: **Neoadjuvant chemoradiotherapy** (CRT) is a standard treatment for **locally advanced rectal cancer**. CRT leads to a better local control; however, this does not translate into a survival benefit. Long-term survival is mostly affected by the development of **distant metastases after** surgery. This study aimed to evaluate **predictive** clinical ...

Cited by: 1

Author: Angelo Restivo, Paolo Delrio, Simona D...

Publish Year: 2020

Predictive Factors of Tumor Response After Neoadjuvant ...

<https://www.sciencedirect.com/science/article/pii/S0360301610002592>

Jun 01, 2011 · Purpose. **Neoadjuvant** chemoradiation followed by surgery is the standard of care for **locally advanced rectal cancer**. The aim of this study was to correlate tumor response to survival and to identify **predictive factors** for tumor response **after** chemoradiation.

Cited by: 71

Author: Laurence Moureau-Zabotto, Bertrand F...

Publish Year: 2011

Early recurrence after neoadjuvant chemoradiation therapy ...

<https://www.sciencedirect.com/science/article/pii/S1015958420302268>

Jul 25, 2020 · Some **locally advanced rectal cancer** (LARC) patients treated with **neoadjuvant chemoradiotherapy** (CRT) prior to total mesorectal excision (TME) show **early** recurrence with a short disease-free interval. This is unacceptable for patients and their families, necessitating re-

Predictive factors for early distant metastasis aft



ALL

IMAGES

VIDEOS



Add the Give with Bing extension >

76,300 Results

Any time ▾

Predictors of Early Distant Relapse in Rectal Cancer ...

<https://www.ncbi.nlm.nih.gov/pubmed/32036373>

Feb 07, 2020 · A higher rate of early distant relapse was associated with **CEA levels above 3 ng/dL (20% vs. 10%; p <0.001)**, **tumor lying under 5 cm from anal verge (20% vs. 9%; p <0.001)**, and **age under 63 years (17% vs. 11%; p = 0.036)**. Multivariate analysis confirmed these factors to be independently correlated with a higher risk of early metastasis.

Cited by: 1

Author: Angelo Restivo, Paolo Delrio, Simona Deidd...

Publish Year: 2020

Predictive Factors of Tumor Response After Neoadjuvant ...

<https://www.sciencedirect.com/science/article/pii/S0360301610002592>

Jun 01, 2011 · Purpose. **Neoadjuvant** chemoradiation followed by surgery is the standard of care for **locally advanced rectal cancer**. The aim of this study was to correlate tumor response to survival and to identify **predictive factors** for tumor response **after** chemoradiation.

Cited by: 72

Author: Laurence Moureau-Zabotto, Bertrand Farna...

Publish Year: 2011

Tumor mutation burden in blood predicts benefit from ...

Search Tools

[Turn on Hover Translation \(开启取词\)](#)

激活 Windows

转到“设置”以激活 Windows。

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

77,600 Results

Any time ▾

Open links in new tab



Pathological complete response may underestimate distant ...

<https://pubmed.ncbi.nlm.nih.gov/30879932>

Pathological complete response may underestimate **distant metastasis in locally advanced rectal cancer following neoadjuvant chemoradiotherapy** and radical surgery: Incidence, metastatic pattern, and risk factors. **Distant metastasis** still remained a major concern in pCR patients following nCRT and TME. **Tumor distance** from the anal verge, acellular mucin pools, and MUC1 expression were **associated with distant metastasis in patients ...**

Cited by: 6

Author: Yanwu Sun, Xuejing Wu, Yiyi Zhang, Huimin...

Publish Year: 2019

Predicting distant metastasis and chemotherapy benefit in ...

<https://www.nature.com/articles/s41467-020-18162-9>

Aug 27, 2020 · **Distant metastasis** (DM) is the main cause of treatment failure **in locally advanced rectal cancer**. Adjuvant chemotherapy is usually used for **distant** control. However, not all patients can benefit ...

Cited by: 2

Author: Zhenyu Liu, Xiaochun Meng, Hongmei Zha...

Publish Year: 2020

Predictors of Early Distant Relapse in Rectal Cancer ...

<https://www.ncbi.nlm.nih.gov/pubmed/32036373>

Feb 07, 2020 · A higher rate of **early distant relapse** was associated with CEA levels above 3 ng/dL (20% vs. 10%; $p < 0.001$), tumor lying under 5 cm from anal verge (20% vs. 9%; $p < 0.001$), and age under 63 years (17% vs. 11%; $p = 0.036$). Multivariate analysis confirmed these factors to be independently correlated with a higher risk of early metastasis.

Cited by: 2

Author: Angelo Restivo, Paolo Delrio, Simona Deidd...

Publish Year: 2020

PEOPLE ALSO ASK

Can rectal cancer cause metachronous metastasis? ▾

Can mesorectal excision help rectal cancer? ▾

What is neoadjuvant chemoradiotherapy? ▾