

Match Overview

1	Internet 34 words crawled on 27-Mar-2020 journals.sagepub.com	1%
2	Internet 34 words crawled on 29-May-2010 www.cmj.org	1%
3	Crossref 28 words Salih Serin, Fazıl Avcı, Onder Ercan, Bülent Köstü, Murat B akacak, Hakan Kiran. "Is neutrophil/lymphocyte ratio a usef	1%
4	Crossref 28 words Burak Yücel, Batuhan Ustun. "Neutrophil to lymphocyte rati o, platelet to lymphocyte ratio, mean platelet volume, red c	1%
5	Crossref 24 words Ana Rejec, Janos Butinar, Jerzy Gawor, Milan Petelin. "Ev aluation of Complete Blood Count Indices (NLR, PLR, MP	<1%
6	Internet 23 words crawled on 09-Aug-2020 www.thieme-connect.com	<1%
7	Internet 15 words crawled on 24-Dec-2019 www.mdpi.com	<1%
8	Internet 15 words crawled on 02-Dec-2020 f6publishing.blob.core.windows.net	<1%
9	Crossref 13 words "Chapter 5088 Halstead-Reitan Battery", Springer Science and Business Media LLC, 2014	<1%

8
Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 59372

Manuscript Type: ORIGINAL ARTICLE

Case Control Study

New indicators in evaluation of hemolysis, elevated liver enzymes, and low platelet syndrome: A case-control study

Kang SY *et al.* New indictors in evaluation of HELLP

Su-Ya Kang, Yun Wang, Li-Ping Zhou, Hong Zhang

ALL

IMAGES

VIDEOS

8,150 Results

Any time ▼

Neutrophil-to-Lymphocyte Ratio, Platelet-to-Lymphocyte ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6572204>

Background and objective: **Neutrophil-to-lymphocyte ratio** (NLR) and **platelet-to-lymphocyte ratio** (PLR) are **new** readily available inflammatory markers that have been analyzed in pregnancy-induced hypertensive disorders such as preeclampsia. Studies on the NLR/PLR **ratio** in hemolysis, elevated liver enzymes, low-**platelet** count (**HELLP**) **syndrome** are limited in the current literature.

Cited by: 2

Author: Giovanni Sisti, Andrea Faraci, Jessica Silva,...

Publish Year: 2019

Neutrophil-to-Lymphocyte Ratio, Platelet-to-Lymphocyte ...

<https://europepmc.org/article/MED/31072037> ▼

May 08, 2019 · **Neutrophil-to-Lymphocyte Ratio, Platelet-to-Lymphocyte Ratio**, and Routine Complete Blood Count Components in **HELLP Syndrome**: A Matched **Case Control Study**. Giovanni Sisti Lincoln Medical and Mental Health Center, Department of Obstetrics and ...

Cited by: 2

Author: Giovanni Sisti, Andrea Faraci, Jessica Silva,...

Publish Year: 2019

Neutrophil-to-Lymphocyte Ratio, Platelet-to-Lymphocyte ...

<https://www.mdpi.com/1010-660X/55/5/123/htm> ▼

Background and objective: **Neutrophil-to-lymphocyte ratio** (NLR) and **platelet-to-lymphocyte ratio** (PLR) are **new** readily available inflammatory markers that have been analyzed in pregnancy-induced hypertensive disorders such as preeclampsia. Studies on the NLR/PLR **ratio** in hemolysis, elevated liver enzymes, low-**platelet** count (**HELLP**) **syndrome** are limited in the current literature.

Cited by: 2

Author: Giovanni Sisti, Andrea Faraci, Jessica Silva,...

Publish Year: 2019

Neutrophil-to-lymphocyte ratio and platelet indices in pre ...

ALL

IMAGES

VIDEOS

69,800 Results

Any time ▼

[PDF] Hemolytic Anemia: Evaluation and Differential Diagnosis

<https://www.aafp.org/afp/2018/0915/afp20180915p354.pdf>

Sep 15, 2018 · **Evaluation** of suspected hemolysis. G6PD = glucose-6-phosphate dehydrogenase; HELLP = hemolysis, **elevated liver enzymes**, and **low platelet** count. Initial hemolysis workup (lactate

Risk Factors, Management, and Outcomes of Hemolysis ...

<https://journals.lww.com/greenjournal/Fulltext/...>

Hemolysis, **elevated liver enzymes**, and **low platelets** (HELLP) **syndrome** is a serious complication of pregnancy that usually occurs in women who have **signs** of preeclampsia. 1 In the absence of hemolysis, the condition has been called **elevated liver enzymes**, **low platelets** (ELLP) **syndrome**. 2,3 Incidence estimates vary from five to 76 per 10,000 deliveries 4,5 and between 8% and 24% of women with ...

PEOPLE ALSO ASK

Is liver enzyme test necessary?



What causes elevated transaminase in the liver?



How does hemolysis affect hemolytic anemia?



Is preeclampsia and hypertension related?



Feedback

Evaluation of Hemolysis as a Severe Feature of Preeclampsia.

<https://www.ncbi.nlm.nih.gov/pubmed/29941517>

We find that **elevated** LDH levels are associated with adverse maternal and neonatal outcomes in hypertension and preeclampsia, independent of hemolysis, **elevated liver enzymes**, and **low platelet**

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

69,300 Results

Any time ▼

[PDF] [Hemolytic Anemia: Evaluation and Differential Diagnosis](https://www.aafp.org/afp/2018/0915/afp20180915p354.pdf)

<https://www.aafp.org/afp/2018/0915/afp20180915p354.pdf>

Sep 15, 2018 · **Evaluation** of suspected hemolysis. G6PD = glucose-6-phosphate dehydrogenase; HELLP = hemolysis, **elevated liver enzymes**, and **low platelet** count. Initial hemolysis workup (lactate

HELLP syndrome (hemolysis, elevated liver enzymes, and low ...

<https://www.uptodate.com/contents/hellp-syndrome...> ▼

Aug 25, 2020 · HELLP is an acronym that refers to a **syndrome** in pregnant and postpartum women characterized by hemolysis with a microangiopathic blood smear, **elevated liver enzymes**, and a **low platelet** count . It probably represents a severe form of preeclampsia (table 1A-B), but the relationship between the two disorders remains controversial.

PEOPLE ALSO ASK

How does hemolysis affect hemolytic anemia? ▼

Is preeclampsia and hypertension related? ▼

What is HELLP syndrome eye morphology? ▼

What are the signs and symptoms of HLP syndrome? ▼

[Feedback](#)

Risk Factors, Management, and Outcomes of Hemolysis ...

<https://journals.lww.com/greenjournal/Fulltext/...>

Hemolysis, **elevated liver enzymes**, and **low** platelets (HELLP) **syndrome** is a serious complication of pregnancy that usually occurs in women who have **signs** of preeclampsia. 1 In the absence of hemolysis, the condition has been called **elevated liver enzymes**, **low** platelets (ELLP) **syndrome**. 2,3 Incidence estimates vary from five to 76 per 10 000 deliveries 4 5 and between 8% and 24% of women with