

Match Overview

1	Crossref 50 words Sen Chen, Yangqun Li, Ning Ma, Weixin Wang, Lisi Xu, Zhe Yang. "Staged Buccal Mucosa Graft Urethroplasty for	1%
2	Internet 40 words crawled on 17-Oct-2020 en.wikipedia.org	1%
3	Internet 38 words crawled on 16-Nov-2019 www.ncbi.nlm.nih.gov	1%
4	Crossref 29 words Xinhai Liu. "A New Use of Scrotal Septal Skin Flap in Repairing Hypospadias :", Annals of Plastic Surgery, 03/2011	1%
5	Internet 21 words crawled on 14-Oct-2020 www.researchsquare.com	1%
6	Crossref 19 words Richard E. Jones. "The Male Reproductive System", Elsevier BV, 1991	1%
7	Crossref Posted Content 19 words Xian Qin, Cai-Yuan Liu, Yi-Lin Xiong, Tao Bai, Lei Zhang, Xiao-Hua Hou, Jun Song. "The clinical features of chron...	1%

Name of Journal: World Journal of Clinical Cases

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Scrotal septal flap and two-stage operation for complex hypospadias: a retrospective study

Running title: Scrotal septal flap for complex hypospadias

Sen Chen, Zhe Yang, Ning Ma, Wei-xin Wang, Li-si Xu, Qi-yu Liu, Yang-qun Li

国内版

国际版

Scrotal septal flap and two-stage operation for complex hypospadias



ALL

IMAGES

VIDEOS

38,600 Results

Any time ▼

[Hypospadias repair: the two-stage alternative - BRACKA ...](#)

<https://bjui-journals.onlinelibrary.wiley.com/doi/...>

WILLIAM DeFOOR, JEFFREY WACKSMAN, Results of Single Staged **Hypospadias Surgery** to Repair Penoscrotal **Hypospadias** With Bifid **Scrotum** or Penoscrotal Transposition, Journal of Urology, 10.1097/01.ju.0000084141.51617.1b, 170, 4 Part 2, (1585-1588), (2003).

Cited by: 261

Author: A. Bracka

Publish Year: 1995

[Hypospadias - ncbi.nlm.nih.gov](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4869453>

Experience with **scrotal septal** Island skin **flap** in repair of penile **hypospadias** ... surgical procedure and its outcome in reconstruction of transposition of the **scrotum**. This is a **retrospective study** carried out at Dhahran Health Centre between 1985 and 1993. 11 patients with this description were seen. ... Recurrent **complex hypospadias** is ...

[Two-stage repair with buccal mucosa for severe and ...](#)

<https://www.onlinelibrary.wiley.com/doi/full/10.1111/j.1442-2042.2010.02684.x>

Objectives: To report the outcomes of **two-stage** repair for adult **hypospadias** using a buccal mucosa graft as a tube. Methods: A total of 57 adult patients who underwent **surgery** for severe urethral defect at our institution from January 2002 to March 2010 were included in this analysis. The urethroplasty procedure included penile urethra reconstruction with the buccal mucosa graft used as a ...

Cited by: 9

Author: Muxin Zhao, Yangqun Li, Yong Tang, We...

Publish Year: 2011

[Clinical efficacy of transverse preputial island flap ...](#)



ALL

IMAGES

VIDEOS

38,400 Results

Any time ▾

Open links in new tab



Hypospadias repair: the two-stage alternative - BRACKA ...

<https://bjui-journals.onlinelibrary.wiley.com/doi/...>

WILLIAM DeFOOR, JEFFREY WACKSMAN, Results of Single Staged **Hypospadias Surgery** to Repair Penoscrotal **Hypospadias** With Bifid **Scrotum** or Penoscrotal Transposition, Journal of Urology, 10.1097/01.ju.0000084141.51617.1b, 170, 4 Part 2, (1585-1588), (2003).

Cited by: 261

Author: A. Bracka

Publish Year: 1995

Search Tools

Turn off Hover Translation (关闭取词)

Hypospadias - ncbi.nlm.nih.gov

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4869453>

Experience with **scrotal septal** Island skin **flap** in repair of penile **hypospadias** ... surgical procedure and its outcome in reconstruction of transposition of the **scrotum**. This is a **retrospective study** carried out at Dhahran Health Centre between 1985 and 1993. 11 patients with this description were seen. ... Recurrent **complex hypospadias** is ...

Two-stage repair with buccal mucosa for severe and ...

<https://www.onlinelibrary.wiley.com/doi/full/10.1111/j.1442-2042.2010.02684.x>

Objectives: To report the outcomes of **two-stage** repair for adult **hypospadias** using a buccal mucosa graft as a tube. Methods: A total of 57 adult patients who underwent **surgery** for severe urethral defect at our institution from January 2002 to March 2010 were included in this analysis. The urethroplasty

ALL

IMAGES

VIDEOS

37,200 Results

Any time ▾

Open links in new tab



[Hypospadias repair: the two-stage alternative - BRACKA ...](#)

<https://bjui-journals.onlinelibrary.wiley.com/doi/...>

WILLIAM DeFOOR, JEFFREY WACKSMAN, Results of Single Staged **Hypospadias Surgery** to Repair Penoscrotal **Hypospadias** With Bifid Scrotum or Penoscrotal Transposition, Journal of Urology, 10.1097/01.ju.0000084141.51617.1b, 170, 4 Part 2, (1585-1588), (2003).

Cited by: 261

Author: A. Bracka

Publish Year: 1995

[Hypospadias - ncbi.nlm.nih.gov](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4869453>

Experience with **scrotal septal** Island skin **flap** in repair of penile **hypospadias** ... surgical procedure and its outcome in reconstruction of transposition of the **scrotum**. This is a **retrospective study** carried out at Dhahran Health Centre between 1985 and 1993. 11 patients with this description were seen. ... Recurrent **complex hypospadias** is ...

[Two-stage repair with buccal mucosa for severe and ...](#)

<https://www.onlinelibrary.wiley.com/doi/full/10.1111/j.1442-2042.2010.02684.x>

Objectives: To report the outcomes of **two-stage** repair for adult **hypospadias** using a buccal mucosa graft as a tube. Methods: A total of 57 adult patients who underwent **surgery** for severe urethral defect at our institution from January 2002 to March 2010 were included in this analysis. The urethroplasty procedure included penile urethra reconstruction with the buccal mucosa graft used as a ...

Cited by: 10

Author: Muxin Zhao, Yangqun Li, Yong Tang, We...

Publish Year: 2011

[The contemporary role of 1 vs. 2-stage repair for proximal ...](#)

<https://tau.amegroups.com/article/view/5287/6187> ▾

Two-stage repairs permit compartmentalization of the repair, providing the opportunity to reevaluate the situation along the way for **complex hypospadias** cases. The time interval between the first and second stage allows for growth of the penile structures and reveals complications that relate to recurrent VC or graft contracture.

Cited by: 6

Author: Shawn Dason, Nathan Wong, Luis H. Braga

Publish Year: 2014

[O63 Braga – Two stage repair for penoscrotal Hypospadias ...](#)

<https://www.hypospadias-society.org/media-lectures/...> ▾

O58 Jones – Family education seminar for **Hypospadias surgery**. O59 McGrath – Parental perception & attitudes towards disclosure of **Hypospadias** repair; O60 Andersson – Urological, cosmetic &

Search Tools

Turn off Hover Translation (关闭取词)