

Match Overview

1

Crossref 13 words

Dan-qing Liu, Hang Zhang, Le Xiao, Bing-yin Zhang, Wei-hui
Liu. "Single-operator cholangioscopy for the treatment of co ...

<1%

Name of Journal: *World Journal of Clinical Cases***Manuscript NO:** 59982**Manuscript Type:** CASE REPORT

**Cystic duct dilation through ERCP in the treatment of gallstones and
choledocholithiasis: a new treatment strategy? – A report of six cases**

A new clinical strategy for cholelithiasis treatment

Abstract**BACKGROUND**

Choledocholithiasis removal *via* endoscopic retrograde cholangiopancreatography (ERCP) then followed by Laparoscopic cholecystectomy (LC) has gradually become the principal method in the treatment of gallstones and choledocholithiasis. We use ERCP approached by cystic duct to treat gallstones combined with choledocholithiasis, which aims to preserve the normal function of the gallbladder simultaneously decrease the risk of biliary tract injury.

Cystic duct dilation through ERCP in the treatment of gallstones a



ALL

IMAGES

VIDEOS

17,800 Results

Any time ▾

Topical nitrate drip infusion using cystic duct tube for ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3692960>

Jun 27, 2013 · In 6 **cases** in which a retained stone was detected following cholecystectomy, topical nitrate drip infusion via **cystic duct** tube (C-tube) was carried out. Retained stones of 2–3 mm diameter and no dilated common bile **duct** in 3 patients were removed by drip infusion of 50 mg GTN or 10 mg ISDN, which was the regular dose of intravenous injection.

Author: Masatoshi Shoji, Hiroshi Sakuma, Yu... **Publish Year:** 2013

Dilation of the cystic duct confluence in laparoscopic ...

<https://bmcsurg.biomedcentral.com/articles/10.1186/s12893-020-00705-y> ▾

Mar 17, 2020 · Then, the **cystic duct** was clipped very near the gall bladder, and 3/4 of its cross section was cut open approximately 0.5 cm from the CBD. The choledochoscope (4.9 mm, Olympus, Tokyo, Japan) was inserted **through** the **cystic duct** and confluence for CBD exploration and stone extraction (success in 9 patients).

Author: Xiao-Bin Yang, An-Shu Xu, Jian-Gan... **Publish Year:** 2020

Laparoendoscopic Rendezvous for Concomitant ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5220453>

1. Introduction. Common bile **duct** (CBD) stones are concomitant with **gallstones** in 10–15% of the patients, with the percentage of association between 8% and 15% in patients under the age of 60 and between 15% and 60% in patients over the age of 60 [].The standard **treatment** for concomitant CBD stones and **gallstones** remains debatable, and there is no consensus on the optimal management **strategy**.

Cited by: 2 **Author:** Bader Hamza Shirah, Zaher Abdulaziz ...

Publish Year: 2016

PEOPLE ALSO ASK

Can cystic duct cause irritation?

Cystic duct dilation through ERCP in the treatment of gallstones and



ALL

IMAGES

VIDEOS

35,000 Results

Any time ▼

Topical nitrate drip infusion using cystic duct tube for ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3692960>

Jun 27, 2013 · In 6 **cases** in which a retained stone was detected following cholecystectomy, topical nitrate drip infusion via **cystic duct** tube (C-tube) was carried out. Retained stones of 2-3 mm diameter and no dilated common bile **duct** in 3 patients were removed by drip infusion of 50 mg GTN or 10 mg ISDN, which was the regular dose of intravenous injection.

Author: Masatoshi Shoji, Hiroshi Sakuma, Yutak... **Publish Year:** 2013

Dilation of the cystic duct confluence in laparoscopic ...

<https://bmcsurg.biomedcentral.com/articles/10.1186/s12893-020-00705-y> ▼

Mar 17, 2020 · Then, the **cystic duct** was clipped very near the gall bladder, and 3/4 of its cross section was cut open approximately 0.5 cm from the CBD. The choledochoscope (4.9 mm, Olympus, Tokyo, Japan) was inserted **through** the **cystic duct** and confluence for CBD exploration and stone extraction (success in 9 patients).

Author: Xiao-Bin Yang, An-Shu Xu, Jian-Gang Li,... **Publish Year:** 2020

Laparoendoscopic Rendezvous for Concomitant ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5220453>

1. Introduction. Common bile **duct** (CBD) stones are concomitant with **gallstones** in 10–15% of the patients, with the percentage of association between 8% and 15% in patients under the age of 60 and between 15% and 60% in patients over the age of 60 []. The standard **treatment** for concomitant CBD stones and **gallstones** remains debatable, and there is no consensus on the optimal management **strategy**.

Cited by: 2

Author: Bader Hamza Shirah, Zaher Abdulaziz Mik...

Publish Year: 2016

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

36,700 Results

Any time ▼

Dilation of the cystic duct confluence in laparoscopic ...

<https://bmcsurg.biomedcentral.com/articles/10.1186/s12893-020-00705-y> ▼

Mar 17, 2020 · Then, the **cystic duct** was clipped very near the gall bladder, and 3/4 of its cross section was cut open approximately 0.5 cm from the CBD. The choledochoscope (4.9 mm, Olympus, Tokyo,...

Author: Xiao-Bin Yang, An-Shu Xu, Jian-Gang Li,... **Publish Year:** 2020

Congenital pancreaticobiliary anomalies in an urban ...

<https://onlinelibrary.wiley.com/doi/10.1002/jgh3.12418>

Pancreaticobiliary anomalies were analyzed using **endoscopic retrograde cholangiopancreatography** (5522 patients) in an urban medical center in the United States. A total of 249 (4.5%) patients were fo...

Predicting cholecystocholedochal fistulas in patients with ...

<https://europepmc.org/article/MED/33177796> ▼

Endoscopic retrograde cholangiopancreatography could serve diagnostic and therapeutic purposes in patients with MS and also help detect cholecystocholedochal fistulas before operation. Our study...

Predictors of failure of endoscopic retrograde ...

[https://www.cell.com/heliyon/fulltext/S2405-8440\(20\)32358-6](https://www.cell.com/heliyon/fulltext/S2405-8440(20)32358-6) ▼

Evidence-based medicine; Internal medicine; Laboratory medicine; Surgery; Common bile **duct**, Lithiasis, ERCP Predictors of failure of **endoscopic retrograde** pancreatocholangiography during common bile **duc**...

Combined Endoscopic and Laparoscopic Management of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4800084>

1. Introduction. Mirizzi syndrome has been defined as bile **duct** obstruction from stone impaction in Hartman's pouch or **cystic duct** [].Mirizzi syndrome has been well-described in the **literature** in patients...

Cited by: 2 **Author:** Arpit Amin, Yuriy Zhurov, George Ibrahim, A...

Publish Year: 2016

Unusual Presentation of Recurrent Gallstone Ileus: A Case ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6794980>

2 **Case Report** A forty nine-year-old female presented to the emergency department due to episodes of