

World Journal of *Orthopedics*

World J Orthop 2021 May 18; 12(5): 254-345



REVIEW

- 254 Fibula fractures management
Canton G, Sborgia A, Maritan G, Fattori R, Roman F, Tomic M, Morandi MM, Murena L

MINIREVIEWS

- 270 Acute syndesmotic injuries in ankle fractures: From diagnosis to treatment and current concepts
Pogliacomini F, De Filippo M, Casalini D, Longhi A, Tacci F, Perotta R, Pagnini F, Tocco S, Ceccarelli F

ORIGINAL ARTICLE**Retrospective Study**

- 292 Reducing unnecessary crossmatching for hip fracture patients by accounting for preoperative hemoglobin concentration
Amin RM, Puvanesarajah V, Chaudhry YP, Best MJ, Rao SS, Frank SM, Hasenboehler EA

- 301 Usefulness of computed tomography based three-dimensional reconstructions to assess the critical shoulder angle
Mah D, Chamoli U, Smith GC

Observational Study

- 310 Potential contribution of pedicle screw design to loosening rate in patients with degenerative diseases of the lumbar spine: An observational study
Bokov A, Pavlova S, Bulkin A, Aleynik A, Mlyavykh S

SYSTEMATIC REVIEWS

- 320 Endoscopic fasciotomy for chronic exertional compartment syndrome of the forearm: Systematic review of outcomes and complications
Marwan Y, Addar A, Algarni N, Algarni N, Burman M, Martineau PA

CASE REPORT

- 329 Rare shear-type fracture of the talar head in a thirteen-year-old child - Is this a transitional fracture: A case report and review of the literature
Monestier L, Riva G, Faoro L, Surace MF
- 338 Three-dimensional printing technology for patient-matched instrument in treatment of cubitus varus deformity: A case report
Sri-utenchai N, Pengrung N, Srikong K, Puncreobutr C, Lohwongwatana B, Sa-ngasoongsong P

ABOUT COVER

Editorial Board Member of *World Journal of Orthopedics*, Mantu Jain, MD, Reader (Associate Professor), Department of Orthopedics, All India Institute of Medical Sciences, Bhubneswar 751019, Odisha, India. montu_jn@yahoo.com

AIMS AND SCOPE

The primary aim of *World Journal of Orthopedics (WJO, World J Orthop)* is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthritis, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

INDEXING/ABSTRACTING

The *WJO* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database. The *WJO*'s CiteScore for 2019 is 3.2 and Scopus CiteScore rank 2019: Orthopedics and Sports Medicine is 77/261.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yan-Xia Xing; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lai Wang.

NAME OF JOURNAL

World Journal of Orthopedics

ISSN

ISSN 2218-5836 (online)

LAUNCH DATE

November 18, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Massimiliano Leigheb

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/2218-5836/editorialboard.htm>

PUBLICATION DATE

May 18, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Fibula fractures management

Gianluca Canton, Andrea Sborgia, Guido Maritan, Roberto Fattori, Federico Roman, Marko Tomic, Massimo Max Morandi, Luigi Murena

ORCID number: Gianluca Canton 0000-0002-8094-0494; Andrea Sborgia 0000-0001-8133-9409; Guido Maritan 0000-0003-0305-6547; Roberto Fattori 0000-0002-5184-3740; Federico Roman 0000-0001-9949-7210; Marko Tomic 0000-0003-3287-9753; Massimo Max Morandi 0000-0002-5923-8553; Luigi Murena 0000-0001-9172-8627.

Author contributions: Canton G designed the structure of the overall manuscript, authored and drafted text in all sections and finally reviewed the manuscript; Sborgia A contributed to the conception and design of the manuscript and drafted and wrote text in all sections; Maritan G, Fattori R and Tomic M contributed to drafting and writing text in different sections; Roman F collected clinical cases and processed images and tables; Murena L and Morandi MM contributed to the supervision and final validation of the manuscript; All authors have read and approved the final manuscript.

Conflict-of-interest statement: Each of the authors declare no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative

Gianluca Canton, Andrea Sborgia, Guido Maritan, Roberto Fattori, Federico Roman, Marko Tomic, Luigi Murena, Department of Medical, Surgical and Life Sciences, Orthopaedics and Traumatology Unit, Trieste University, Trieste 34149, Italy

Massimo Max Morandi, Department of Orthopaedic Surgery, Louisiana State University Health Science Center, Shreveport, LA 71103, United States

Corresponding author: Gianluca Canton, MD, Associate Professor, Department of Medical, Surgical and Life Sciences, Orthopaedics and Traumatology Unit, Trieste University, 447 Strada di Fiume, Trieste 34149, Italy. gcanton@units.it

Abstract

Isolated distal fibula fractures represent the majority of ankle fractures. These fractures are often the result of a low-energy trauma with external rotation and supination mechanism. Diagnosis is based on clinical signs and radiographic exam. Stress X-rays have a role in detecting associated mortise instability. Management depends on fracture type, displacement and associated ankle instability. For simple, minimally displaced fractures without ankle instability, conservative treatment leads to excellent results. Conservative treatment must also be considered in overaged unhealthy patients, even in unstable fractures. Surgical treatment is indicated when fracture or ankle instability are present, with several techniques described. Outcome is excellent in most cases. Complications regarding wound healing are frequent, especially with plate fixation, whereas other complications are uncommon.

Key Words: Fibula fracture; Lateral malleolus; Distal fibula; Management; Treatment; Ankle

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Isolated fibula fractures are very common injuries. Diagnostic exams must rule out ankle instability. Surgical treatment must be considered in the case of associated ankle instability. Risk factors for wound related complications must be considered when choosing the surgical technique.

Citation: Canton G, Sborgia A, Maritan G, Fattori R, Roman F, Tomic M, Morandi MM,

Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Specialty type: Orthopedics

Country/Territory of origin: Italy

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): D
Grade E (Poor): 0

Received: October 31, 2020

Peer-review started: October 31, 2020

First decision: January 18, 2021

Revised: February 1, 2021

Accepted: April 5, 2021

Article in press: April 5, 2021

Published online: May 18, 2021

P-Reviewer: Amorniyotin S, Tsikopoulos K

S-Editor: Liu M

L-Editor: Filipodia

P-Editor: Yuan YY



Murena L. Fibula fractures management. *World J Orthop* 2021; 12(5): 254-269

URL: <https://www.wjgnet.com/2218-5836/full/v12/i5/254.htm>

DOI: <https://dx.doi.org/10.5312/wjo.v12.i5.254>

INTRODUCTION

Ankle fractures are frequent injuries[1], increasing in elderly patients as a consequence of osteoporosis[2]. In most literature reports, distal fibula fractures represent the majority of ankle fractures[3]. These fractures are often the result of a low energy trauma with an external rotation and supination mechanism.

Many authors recommend conservative treatment for isolated fibula fractures without signs of ankle instability as good clinical results are obtained in most cases[1-3]. However, the trend in recent years is headed towards surgical treatment, with advantages in terms of anatomic restoration and earlier recovery[1-3].

Depending on fracture type, displacement and degree of instability, several surgical treatment techniques have been described. These include lateral *vs* posterolateral plating, nonlocking *vs* locking plate fixation, isolated screws and intramedullary fixation[4].

The aim of the present paper is to review the most recent literature about the epidemiology, mechanism of injury, diagnosis, classification, management and complications of isolated distal fibula fractures treatment.

EPIDEMIOLOGY

Ankle fractures are frequent injuries, accounting for about 9% of all fractures[1]. Moreover, there has been a sharp increase in osteoporosis related ankle fracture incidence in recent years. Isolated distal fibula fractures represent the most frequent ankle fracture type[3,4]. Elsoe *et al*[2] recently reported the epidemiology of 9767 ankle fractures, identifying distal fibula fractures as the most common fracture type, accounting for 55% of cases. Furthermore, according to the Arbeitsgemeinschaft für Osteosynthesefragen/Orthopaedic Trauma Association (AO/OTA) classification, type B is generally reported to be the more common distal fibula fracture type. Court-Brown *et al*[5] reported the following distal fibula fracture type distribution among 1500 ankle fractures: 52% type B trans-syndesmotic fractures, 38% type A infra-syndesmotic fractures and 10% type C supra-syndesmotic fractures[5]. Age and gender-related differences in ankle fracture epidemiology have been reported. Distal fibula fractures are more frequent in young active male patients. Werner *et al*[6] reported an incidence of isolated distal fibula fractures reaching 83% of cases among a population of National Football League athletes reporting ankle fractures. Conversely, Hasselman *et al*[7] found isolated fibular fracture to cover 57.6% of cases in elderly (> 65 years) women reporting ankle fractures.

MECHANISM OF LATERAL MALLEOLUS FRACTURE

Stability of the ankle mortise is determined by bony components (fibula, tibia and talus) and ligamentous structures (syndesmosis complex and lateral and medial collateral ligaments). Dynamic musculotendinous stabilizers, which exact function is less understood, also play a relevant role. There is on-going research on interactions between these structures and mechanisms that cause fracture.

Ankle sprains/torsion injuries, accidental falls and sports related accidents are the most frequently reported causes of distal fibula fracture, with different rates according to the different AO/OTA types. In type A fractures the main cause is represented by torsion (32%) followed by falls (23%) and sports related trauma (22%). In type B fractures the trend is similar, with reported rates of 27% for torsion, 37% for falls and 13% for sports related trauma. For type C the trend is slightly different because torsion represents only 3.7% of cases while falls and sports related trauma represent 28% and 21% of cases, respectively.

The most frequently described traumatic mechanism is supination-external rotation (SER). In this type of trauma, the talus rotates pushing the tibia and fibula apart, rupturing the anterior-inferior tibiofibular ligament and causing a simple ankle sprain.

Further progression of deforming force causes a simple oblique fibula fracture at the level of the syndesmosis, equivalent to the AO/OTA type B fracture. In the second most frequent mechanism, supination-adduction, adduction of the hindfoot causes either talofibular ligament rupture (ankle sprain) or an avulsion fracture of the distal fibula, equivalent to the AO/OTA type A fracture. As reported by Lauge-Hansen[8], lateral structures are damaged after the medial side when traumatic forces act in pronation.

Nonetheless, a recent *in vivo* study by Kwon *et al*[9] analyzing injury videos posted on YouTube and matching them to their corresponding X-rays, found that the Lauge-Hansen system was only 58% overall accurate in predicting fracture patterns from deforming injury mechanism.

DIAGNOSIS

Clinical examination

Clinical signs of an isolated fracture of the distal fibula are not specific and may not be distinguishable from a severe ankle sprain. These signs include swelling, bruising, pain, ecchymosis, tenderness and reduced range of motion (ROM). Swelling is the most common reported sign and was found to be a constant feature of all ankle fractures[10]. Because most isolated distal fibula fractures are stable, weight bearing is usually possible[11], thus patients might be ambulating at clinical presentation. There are no specific clinical tests for this fracture. Nonetheless, it is essential to evaluate medial ankle structures stability to choose the correct management. In C type fractures, the syndesmosis complex integrity must also be investigated. However, clinical examination alone is not diagnostic in most cases because pain, edema and muscle contracture can hinder correct evaluation[12,13].

Radiographic evaluation

Standard ankle X-rays are the mainstay of instrumental diagnosis for all ankle fractures. However, ankle sprains that might possibly cause a fracture might not deserve radiographic examination in all cases. In fact, due to the already described unspecific clinical presentation, other criteria should be considered to reduce the number of unnecessary exams and length of hospital stay[14,15]. The Ottawa Criteria were introduced for this purpose, despite some studies questioning their clinical validity[16].

When cases amenable for radiographic evaluation are selected, three radiographic views should always be obtained according to the American College of Radiology guidelines: antero-posterior (AP), lateral and mortise view[17].

The AP view is performed along the long axis of the foot. In isolated fibula fractures, this view is particularly useful to evaluate signs of associated ankle and/or syndesmosis instability through the analysis of talus coronal inclination, tibio-fibular overlap, tibio-fibular clear space and medial clear space (MCS)[18].

In the lateral view, the talar dome must be centered and congruent with the tibial plafond. This view is useful in isolated fibula fractures to demonstrate AP displacement and external rotation type fractures[19].

The mortise view is taken by placing the foot on the table with about 15° of internal rotation. This visualization is useful in isolated fibula fractures to detect signs of associated syndesmosis instability and to obtain a clear view of the lateral malleolus without other overlapping structures. It is especially useful in undisplaced and incomplete fractures[18,20].

Theoretically, these views are sufficient to identify an isolated distal fibula fracture in almost all cases. As demonstrated in several clinical and cadaveric studies[21-23] in standard radiographic evaluation, a MCS increase in isolated distal fibula fractures is a typical sign of rupture of the deltoid ligament with consequent talar lateral shift. These factors suggest a possible mortise instability, which is fundamental to define treatment modality. However, reliable radiographic determination of deltoid ligament rupture in such uncertain cases is difficult in clinical practice[24]. To adequately evaluate mortise stability, different modalities to obtain stress radiographs have been described.

The manual stress view has been considered the method of choice for years. However, being an operator dependent exam, reproducibility and radiation exposure to the physician are a concern. Moreover, it is not clear in the published data which MCS values are to be considered as a cut off to identify a clinically relevant deltoid ligament rupture.

Michelson *et al*[25] described for the first time in 2001 the gravity stress view to investigate ankle joint instability. The patient is placed in the lateral decubitus position with the distal half of the leg off the end of the table. Then a standard AP and mortise view is taken. Gravity stress views proved to be just as effective as manual stress views to detect deltoid ligament injury in association with an isolated distal fibula fracture[26-28]. Many advantages, such as no radiation exposure to the physician and less pain to the patient are described. Moreover, the constant force of gravity makes the exam reproducible[27].

Nonetheless, manual and gravity stress radiographs can overestimate the need for surgical fixation by showing MCS widening in partial deltoid ligament lesions that might uneventfully heal with conservative treatment[29-31]. For this reason, in the literature many authors described the weight bearing radiographs as an alternative method to identify ankle instability in isolated distal fibula fractures. As proposed by Weber *et al*[32], these radiographs should be performed with the patient standing on both legs as pain allows. An AP, lateral and mortise view is then taken with the weight ideally distributed equally on both ankles[32]. The disadvantage of this technique is the possible variability in weight distribution between healthy and injured side that could hide the degree of ankle instability in some cases.

Finally, magnetic resonance imaging is not indicated in the acute diagnosis of isolated distal fibula fractures. Its use in these lesions is very limited because traditional and stress radiographs proved to be equally effective in identifying the severity of associated deltoid ligament injury[33]. Conversely, magnetic resonance imaging might be useful to detect associated chondral injuries or to diagnose a fracture when conventional radiographs are inconclusive, and clinical suspicion is very high[34].

CLASSIFICATION

Although the fibula carries only 10% of the body weight (compared to 90% carried by the tibia)[35], its role is crucial in the stability of the ankle mortise. Based on this statement, to define an ankle with isolated fibula fracture as stable or unstable is crucial to guide proper treatment. Nonetheless, there is still debate in the literature on which are the most suitable clinical and radiological criteria to obtain this goal. The optimal management is based on an accurate knowledge of the fracture. For a comprehensive assessment of the fracture, a reproducible classification method is essential.

Different classifications have been proposed through the years. Lauge-Hansen and Danis-Weber classifications are the most used. They are based on standard AP, lateral and mortise radiographic views of the ankle. Their aim is to describe the mechanism of injury, predict soft tissue conditions and finally guide treatment.

The Lauge-Hansen classification, developed in 1954, is based on the position of the foot at the time of injury (supination or pronation) and on the deforming forces acting on the foot (abduction, adduction or external rotation)[36]. This results in a combination of four categories and thirteen patterns of ankle fracture. Isolated distal fibula fractures usually occur in the supination-external rotation type II and supination-adduction Lauge-Hansen types. In the other classes (pronation-abduction and pronation-external rotation) an isolated fibula fracture might occur above the level of the syndesmosis in association with a medial ligament injury. The main limitation of the Lauge-Hansen classification is the poor interobserver agreement that is instead very high in the Danis-Weber classification due to its simplicity.

The Danis-Weber classification (Figure 1) was first described by Robert Danis in 1949 and later modified by Bernhard Georg Weber in 1966. It was then adopted by the AO/OTA Group. It evaluates the location of the main fibular fracture in relation to the syndesmosis. Type A fractures are generally stable injuries occurring below the level of the syndesmosis. Type B fractures occur at the level of the syndesmosis and might be unstable in some cases. Type C fractures are usually unstable injuries occurring above the level of the syndesmosis[37].

However, differentiating fracture types in relation to the syndesmosis might lead the medial side of the ankle to be overlooked. Moreover, injury extent in the tibiofibular syndesmosis is often not predictable. These findings are crucial to detect tibio-talar instability and consequently to decide between surgical and nonsurgical management, especially for type B fractures[38]. Probably, as suggested by Lampridis *et al*[39], a combination of the two main systems is the correct approach. Nonetheless, the limits of these classification systems are reported by several studies in the literature, especially their poor prognostic and therapeutically predictive capabili-



Figure 1 Danis-Weber classification of distal fibula isolated fractures. A: Type A; B: Type B; C: Type C.

ties[40-42].

MANAGEMENT

Clinical and biomechanical data indicate that maintenance of talar reduction is the most important factor for the prognosis of ankle fractures[43,44]. A residual dislocation leads to a series of complications including impaired healing, early osteoarthritis and residual instability.

Shortening and external rotation of the fibula can cause talar lateral shift, reducing the tibio-talar contact area and increasing peak pressure in the articular cartilage[45,46]. Many authors believe that the deltoid ligament is the most important structure to maintain the position of the talus when the fibula is fractured[47-49].

An isolated distal fibula fracture is considered stable when less than 2 mm of displacement occurs, and no deltoid ligament rupture is detected (MCS < 4 mm)[50,51]. Several clinical studies have shown that in isolated fractures without concomitant medial injury, conservative treatment leads to excellent long-term results[52,53].

Conversely, many clinical studies have shown significantly better results in ankle fractures with mortise instability and talus displacement when an accurate fracture reduction is achieved[54]. However, unlike cases associated with gross instability, proper management of isolated fibula fractures that demonstrate instability only after stress radiographs is still a matter of debate in the literature[55]. In many authors' opinions, stress radiographs can overestimate the need for surgical fixation[29-31]. Hoshino *et al*[31] analyzed 36 patients with isolated distal fibula fracture demonstrating MCS widening at external rotation stress radiographs. All patients were treated with nonsurgical treatment initially. After 7 d, only one (3%) patient had an unstable ankle mortise on weight bearing radiographs, thus requiring operative treatment. All patients demonstrated good to excellent functional scores at 1 year follow-up[31].

NONOPERATIVE TREATMENT

Conservative treatment is indicated for isolated distal fibula fractures with a stable ankle mortise. As far as fracture displacement is concerned, Lesic *et al*[56] set 2 mm as the threshold between conservative and surgical treatment. However, there is no strong evidence in the literature advocating surgery for fracture displacement more than 2 mm[57]. Other studies suggest that radiographic displacement might not be reliable as it is mostly a rotational displacement[58-60].

Nonetheless, minor radiographic displacement seems not to affect clinical outcome[61]. Two studies have shown a high percentage of good results even when the fibula is posteriorly displaced up to 5 mm[41,62]. Hence, any isolated distal fibula fracture with a stable ankle can be treated conservatively (Figure 2).



Figure 2 Lateral, mortise and antero-posterior radiographic views of a Danis-Weber type B left distal fibula fracture. Conservative treatment is the correct choice for this case because minimal displacement of the fracture and absence of associated ankle instability are demonstrated.

Weber type A fractures can be considered equivalent to a ligamentous ankle injury. Likely, satisfactory results with nonoperative treatment can be achieved as in ligament ruptures[63].

In the literature, randomized and nonrandomized studies show satisfactory outcomes for conservative treatment in minimally displaced or nondisplaced Weber B type fractures[41,61,62,64]. Dobbe *et al*[65] reported that 13% of 108 conservatively treated infra-syndesmotic fractures had difficulties with work- and life-related activities. However, no relationship was identified between outcomes and the degree of articular displacement or fragment width[65]. These results might suggest the need to adapt treatment according to age and activity level. Sanders *et al*[54] suggested operative intervention in younger individuals, with the aim to reduce the risk of malalignment and improve outcomes. Conversely, a different management can be reserved for older and less active patients, which can be safely treated with cast immobilization even in unstable fracture patterns[54].

While there are several studies that describe conservative treatment for Weber type B fractures, a recent review comparing different managements for Weber type C fractures found only one study included conservative treatment. This demonstrates the widespread preference for surgical management in these cases[66]. Donken *et al*[67] compared the results of nonoperative and operative treatment for Weber type C fractures, with conservative treatment reserved to cases that demonstrated joint congruity, no signs of deltoid ligament injury and no medial malleolus fractures. Clinical results were comparable with most patients reaching high-level functional results[67].

Nonoperative treatment modality is chosen based on patients' symptoms, bone and skin quality, time lapse from injury to clinical presentation and risk factors for impaired healing (Table 1). Different conservative treatment modalities are described in the literature, ranging from cast immobilization without weight bearing to immediate full weight bearing without cast or brace. Historically, treatment in a plaster cast for several weeks was recommended. This strategy arises from the evidence that 6 wk are needed for any fracture to tolerate weight bearing[41,61,62,68]. Although a high rate of fracture union was demonstrated, a prolonged immobilization can result in ankle stiffness and higher risk of deep vein thrombosis[69-73]. To overcome these complications, Kortekangas *et al*[74] recently showed that a 3-wk period of immobilization is noninferior to 6 wk in the treatment of an isolated stable Weber B type fracture.

Alternative methods of immobilization with immediate weight bearing have been proposed through the years. In 1979, Stover *et al*[75] proposed a bivalve pneumatic air stirrup in the management of ankle fractures. Stuart *et al*[71] in 1989 showed the brace to improve patient comfort, post fracture swelling, range of ankle motion at union and time to full rehabilitation. Similar findings were reported by Brink *et al*[76], who advocated the use of a hinged short-leg boot with good results in pain relief, increased ROM and earlier return to ambulation. It has been suggested that an ordinary elastic bandage is equally safe and beneficial[52], and no difference in the amount of pain experienced has been found between early mobilization or plaster cast[77]. Ryd *et al*[52] described 49 patients treated only with elastic bandage for isolated distal fibula fractures displaced less than 2 mm and without medial tenderness. All of them had excellent clinical outcomes and were back to normal activity in about 4 mo[52].

Table 1 Summary of main criteria for conservative indication in isolated distal fibula fractures treatment

Nonoperative treatment	
Displacement	< 2 mm
Medial stability	MCS < 4 mm in AP/mortise view and/or in dynamic radiographs view
Poor bone and skin quality	
Long time lapse from injury	
Advanced age, low functional demand	
High risk of local and general complications	

AP: Antero-posterior; MCS: Medial clear space.

The functional treatment of stable ankle fractures is also supported by van der Berg *et al*[78], who showed better Visual Analogue Scale score and total ROM with a brace rather than with a cast after 6 wk, while no significant difference was found at 1 year.

SURGICAL TREATMENT

Open reduction and internal fixation is the most common treatment for unstable ankle fractures (Table 2). There are several fixation methods described for distal fibula fractures fixation, including one-third tubular plate (Figure 3), dynamic compression plate and locking plate with or without an independent lag screw[79-81]. The most used plates are angular stable metaphyseal or anatomic distal fibula plates (Figure 4) [82]. They can serve as bridging plates, compression plates, tension band plates or neutralization plates. Most studies comparing locking plates and conventional one third tubular plates show no differences in clinical and radiographic outcomes as well as in wound complications incidence[79,83-85]. However, these fixation techniques have a complication rate of up to 30% of cases, with wound complications being the most common[86-88]. This is attributed to the surgical trauma occurring in an already injured area with limited soft tissue cover. This range increases in smokers, in elderly patients and in patients with comorbidities such as diabetes and peripheral vasculopathy[87-90]. Minimally invasive plate osteosynthesis is a suggested alternative to prevent periosteal damage and major soft tissues dissection. This technique allows plate insertion through a small incision and better respects fracture biology[79]. However, anatomic restoration is more technically demanding for the surgeon, and the technique is suitable for only a small amount of fracture patterns.

Distal fibula fractures in elderly patients are often comminuted and present with impaired soft tissues coverage. Consequently, the correct management of ankle fractures in these patients must account for bone quality and the risk of soft tissue complications[80,86,89]. Locking plates provide a biomechanical advantage in cases of poor bone quality and are therefore recommended over nonlocking plates in osteoporotic patients when surgical management is chosen[86,80].

Fibular nailing is considered a valid alternative method of fixation for distal fibula fractures. The use of intramedullary fibula fixation was first introduced in the mid-1980s to reduce complications of the traditional plating techniques[91]. However, early attempts in intramedullary fibula fixation, such as using rush rods, Inyo nail, K. wires, *etc.*, showed several complications and failures due to poor rotational and longitudinal stability, which led to loss of reduction, malunion and nail migration[92-94]. As a result, modern locking fibula nails have been developed to reduce such complications[93]. Modern fibula nails are designed with proximal and distal locking fixation systems. The use of proximal and distal locking screws as well as of intersyndesmotomic screws (Figure 5) if needed, allows optimal fracture stability with consequent advantages on early weight bearing[93,95]. The use of a modern fibula intramedullary nail should be considered especially in cases of suffering skin and/or patients with severe comorbidities[93-95]. Several authors compared the results of plating and nailing for internal fixation of the fibula in ankle fractures, and most of them show no differences in functional outcome[96,97]. Successful results with lower rates of wound breakdown and hardware prominence compared with standard open reduction and plating have been reported[93-96]. On the other hand, the main criticisms of this

Table 2 Summary of main criteria for surgical indication in isolated distal fibula fractures treatment

Surgical treatment	
Displacement	> 2-5 mm
Medial instability	MCS > 4 mm in AP/mortise view or in dynamic radiographs view
Type C	Any displacement

AP: Antero-posterior; MCS: Medial clear space.



Figure 3 Radiographic and clinical results at 3 mo from a one-third tubular plate fixation of a Danis-Weber type B left distal fibula fracture.

technique reside in the difficult management of rotational displacement, especially in type B fractures and in the absence of compression at the fracture site in simple oblique and spiral fracture patterns with a theoretical risk of delayed healing[93,95]. However, the minimally invasive approach that preserves fracture biology and periosteal blood supply might overcome this aspect[94,98].

COMPLICATIONS

Nonunion

Nonunion in ankle fractures is an extremely rare complication[99-103]. Distal fibula fractures usually heal uneventfully even with nonsurgical treatment. However, because most nonunions in this area are asymptomatic, the exact incidence of this complication is uncertain[99-103].

A higher incidence of fibular nonunion has been described when associated with medial instability. Sneppen *et al*[99] demonstrated a 0.7% incidence of fibular nonunion in ankle fractures involving the medial malleolus *vs* a 0.1% incidence in isolated fibular fractures. Complete nonunion was mostly seen in type A (Figure 6) and C fractures[100], whereas incomplete nonunion was described in Weber type B fractures[101].

Treatment is based on the type of nonunion, symptoms, initial fracture pattern, associated injuries and patient expectations. Asymptomatic nonunions are treated conservatively, with reports of spontaneous healing even several years after the initial injury[102]. For symptomatic nonunion, open reduction and internal fixation with or without bone grafting represents the best treatment choice resulting in successful outcomes in most cases[103].

Malunion

Angular malalignment occurs when the distal fibula heals in shortening or external rotation. This causes a lateral subluxation of the talus with ankle kinematics alteration leading to arthritis[44]. Most cases of angular malalignment occur after conservative treatment[104]. However, surgical treatment might be a cause of malunion if anatomic reduction is not achieved intraoperatively. Moreover, technical errors, suboptimal stability of fixation and unrecognized associated ligamentous instability might lead to loss of reduction and consequent malunion (Figure 7)[105]. Several radiographic parameters have been described to identify the correct length and rotation of the fibula



Figure 4 Clinical and radiographic results at 3 mo from surgical treatment of a Danis-Weber type B right distal fibula fracture with an anatomic angular stable plate.

in the ankle mortise: the Weber circle, the Shenton's line, the talocrural and bimalleolar angles[106]. These parameters can help the clinicians to more easily identify a malalignment that could cause continuous pain to the patient even after several months.

Surgical treatment is a demanding procedure, as anatomic reconstruction usually requires both fibula osteotomy and soft tissues release. Plate osteosynthesis is then required for fixation[107]. In several studies, the results of surgical treatment for malalignment were excellent. Yablon *et al*[108] reported good results over 7 years of follow up in 23 of 26 patients with fibular malalignment surgically treated 6 years after trauma. Ward *et al*[109] reported similar results in ankle fracture malunions treated with lengthening osteotomy of the fibula.

Wound dehiscence and infection

The distal fibula is subcutaneous and lacks a layer of overlying muscles. Thus, wound healing complications are the most common adverse events related to distal fibula fixation. Wound edge necrosis, wound dehiscence and superficial and deep infection have all been reported[110]. In the literature, the overall wound complication rate varies from 8.4% to 40.0% among studies[111-113]. The deep surgical infection rate is significantly lower, about 1.2% to 2.8% according to different reports[111,114,115].

The genesis is multifactorial as it depends on soft tissue compromise at admission, timing of surgery, fracture type and patient characteristics. Age, diabetes, steroid intake, smoking and peripheral vascular disease are associated with a greater risk of wound complications[88,112,116-118]. The type of implant also might have a role. Schepers *et al*[119] reported in a retrospective study of 165 patients a higher wound complication rate with locking plates (17.5%) than with thinner one third tubular plates (5.5%). Conversely, Tsukada *et al*[84] in two different trials did not find any difference in complication rates between locking and nonlocking plates[84,115].



Figure 5 Clinical case of a Danis-Weber Type B left distal fibula fracture treated with locked nailing. A: Intraoperative image demonstrating fibula nail insertion with guided instrumentation; B: Detail of the clinical result of minimally invasive approach for fibular nail insertion; C: Antero-posterior X-rays demonstrating fracture fixation with fibular nail completed with two guided intersyndesmodic screws.



Figure 6 Lateral and antero-posterior view X-rays taken 4 mo after conservative treatment of a Danis-Weber type A distal fibula fracture. Despite the radiographic evidence of nonunion the patient is completely asymptomatic, and no further treatment is indicated.

The incidence of this complication can be minimized by treating the fracture as soon as reasonably possible[110] or postponing the surgery until the edema is resolved. The latter strategy has gained more and more popularity with the advent of damage control techniques. Whatever the timing, limiting the use of the tourniquet and closing the wound without tension is also advisable.

ROM limitation

Most patients after isolated distal fibula fractures recover with a completely functional ROM. When stiffness occurs, dorsiflexion deficit is more common. Lin *et al*[120] reported that in a population of 306 patients a 19% rate of plantar flexion limitation (only 2% > 10 degrees) occurred, while 41% of cases had restriction in dorsiflexion. In a review of 31 randomized trials about rehabilitation of ankle fractures, the authors found a positive effect on ankle ROM from early mobilization, early weight bearing and the use of a removable immobilization device. However, there is limited evidence supporting this strategy as patient compliance seems to play a significant role[120].



Figure 7 Clinical case of a Danis-Weber type C left distal fibula fracture treated with a one-third tubular plate. At post-op X-rays, malreduction with residual medial displacement is demonstrated. A dedicated adjustable tibio-fibular suture button fixation was added to obtain anatomic reduction and correction of associated ankle instability.

Early arthritis

Trauma is the most common cause of ankle osteoarthritis[121]. Arthritis results from a combination of direct cartilage damage and biomechanical alterations that affect joint kinematics[122]. It becomes evident 2 to 3 years after trauma and is often symptomatic. It is more frequent in higher grade fractures according to the Lauge-Hansen classification. Lübbecke *et al*[105], evaluating risk factors for development of ankle osteoarthritis, found a Weber type C or a fibula fracture with an associated fracture of the medial malleolus as the most important risk factors for the development of symptomatic radiographically advanced osteoarthritis. Surgical treatment seems to reduce the incidence of osteoarthritis by means of anatomic reduction and ankle stability restoration.

CONCLUSION

Isolated fibula fractures are very common injuries. Diagnostic exams must rule out ankle instability. Conservative treatment yields good results in stable fractures with stable ankle mortise. Open reduction internal fixation is indicated in case of associated ankle instability. Risk factors for wound related complications must be considered when choosing a surgical technique.

REFERENCES

- 1 **Court-Brown CM**, Caesar B. Epidemiology of adult fractures: A review. *Injury* 2006; **37**: 691-697 [PMID: 16814787 DOI: 10.1016/j.injury.2006.04.130]
- 2 **Elsoe R**, Ostgaard SE, Larsen P. Population-based epidemiology of 9767 ankle fractures. *Foot Ankle Surg* 2018; **24**: 34-39 [PMID: 29413771 DOI: 10.1016/j.fas.2016.11.002]
- 3 **Jehlicka D**, Bartonicek J, Svatos F, Dobiás J. [Fracture-dislocations of the ankle joint in adults. Part I: epidemiologic evaluation of patients during a 1-year period]. *Acta Chir Orthop Traumatol Cech* 2002; **69**: 243-247 [PMID: 12362627]
- 4 **Aiyer AA**, Zachwieja EC, Lawrie CM, Kaplan JRM. Management of Isolated Lateral Malleolus Fractures. *J Am Acad Orthop Surg* 2019; **27**: 50-59 [PMID: 30278012 DOI: 10.5435/JAAOS-D-17-00417]
- 5 **Court-Brown CM**, McBirnie J, Wilson G. Adult ankle fractures--an increasing problem? *Acta Orthop Scand* 1998; **69**: 43-47 [PMID: 9524517 DOI: 10.3109/17453679809002355]
- 6 **Werner BC**, Mack C, Franke K, Barnes RP, Warren RF, Rodeo SA. Distal Fibula Fractures in National Football League Athletes. *Orthop J Sports Med* 2017; **5**: 2325967117726515 [PMID: 28959699 DOI: 10.1177/2325967117726515]
- 7 **Hasselmann CT**, Vogt MT, Stone KL, Cauley JA, Conti SF. Foot and ankle fractures in elderly white women. Incidence and risk factors. *J Bone Joint Surg Am* 2003; **85**: 820-824 [PMID: 12728031 DOI: 10.2106/00004623-200305000-00008]
- 8 **Lauge-Hansen N**. Fractures of the ankle. II. Combined experimental-surgical and experimental-roentgenologic investigations. *Arch Surg* 1950; **60**: 957-985 [PMID: 15411319 DOI: 10.1093/00004623-200305000-00008]

- 10.1001/archsurg.1950.01250010980011]
- 9 **Kwon JY**, Chacko AT, Kadzielski JJ, Appleton PT, Rodriguez EK. A novel methodology for the study of injury mechanism: ankle fracture analysis using injury videos posted on YouTube.com. *J Orthop Trauma* 2010; **24**: 477-482 [PMID: 20657256 DOI: 10.1097/BOT.0b013e3181e99264]
 - 10 **Montague AP**, McQuillan RF. Clinical assessment of apparently sprained ankle and detection of fracture. *Injury* 1985; **16**: 545-546 [PMID: 4066016 DOI: 10.1016/0020-1383(85)90082-8]
 - 11 **Lynch SA**. Assessment of the Injured Ankle in the Athlete. *J Athl Train* 2002; **37**: 406-412 [PMID: 12937562]
 - 12 **Gebing R**, Fiedler V. [X-ray diagnosis of ligament lesions of the superior ankle joint]. *Radiologe* 1991; **31**: 594-600 [PMID: 1780435]
 - 13 **Oloff LM**, Sullivan BT, Heard GS, Thornton MC. Magnetic resonance imaging of traumatized ligaments of the ankle. *J Am Podiatr Med Assoc* 1992; **82**: 25-32 [PMID: 1545367 DOI: 10.7547/87507315-82-1-25]
 - 14 **Stiell IG**, Greenberg GH, McKnight RD, Nair RC, McDowell I, Reardon M, Stewart JP, Maloney J. Decision rules for the use of radiography in acute ankle injuries. Refinement and prospective validation. *JAMA* 1993; **269**: 1127-1132 [PMID: 8433468 DOI: 10.1001/jama.269.9.1127]
 - 15 **Stiell I**, Wells G, Laupacis A, Brison R, Verbeek R, Vandemheen K, Naylor CD. Multicentre trial to introduce the Ottawa ankle rules for use of radiography in acute ankle injuries. Multicentre Ankle Rule Study Group. *BMJ* 1995; **311**: 594-597 [PMID: 7663253 DOI: 10.1136/bmj.311.7005.594]
 - 16 **Stiell IG**, Greenberg GH, McKnight RD, Wells GA. The "real" Ottawa ankle rules. *Ann Emerg Med* 1996; **27**: 103-104 [PMID: 8572439 DOI: 10.1016/s0196-0644(96)70309-8]
 - 17 **Dalinka MK**, Alazraki N, Berquist TH, Daffner RH, DeSmet AA, el-Khoury GY, Goergen TG, Keats TE, Manaster BJ, Newberg A, Pavlov H, Haralson RH 3rd, McCabe JB, Sartoris D. Imaging evaluation of suspected ankle fractures. American College of Radiology. ACR Appropriateness Criteria. *Radiology* 2000; **215** Suppl: 239-241 [PMID: 11037432]
 - 18 **Krähenbühl N**, Weinberg MW, Davidson NP, Mills MK, Hintermann B, Saltzman CL, Barg A. Imaging in syndesmotic injury: a systematic literature review. *Skeletal Radiol* 2018; **47**: 631-648 [PMID: 29188345 DOI: 10.1007/s00256-017-2823-2]
 - 19 **Lampignano JP**, Kendrick LE. Bontrager's Textbook of Radiographic Positioning and Related Anatomy. 9th ed. Mosby, 2017: 239-242
 - 20 **Patel P**, Russell TG. Ankle Radiographic Evaluation. [cited 14 September 2020]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls [PMID: 32491394]
 - 21 **Harper MC**. The deltoid ligament. An evaluation of need for surgical repair. *Clin Orthop Relat Res* 1988; **(226)**: 156-168 [PMID: 3121227 DOI: 10.1097/00003086-198801000-00022]
 - 22 **Harper MC**. Deltoid ligament: an anatomical evaluation of function. *Foot Ankle* 1987; **8**: 19-22 [PMID: 3623356 DOI: 10.1177/107110078700800104]
 - 23 **Baird RA**, Jackson ST. Fractures of the distal part of the fibula with associated disruption of the deltoid ligament. Treatment without repair of the deltoid ligament. *J Bone Joint Surg Am* 1987; **69**: 1346-1352 [PMID: 3440794 DOI: 10.2106/00004623-198769090-00007]
 - 24 **Schuberth JM**, Collman DR, Rush SM, Ford LA. Deltoid ligament integrity in lateral malleolar fractures: a comparative analysis of arthroscopic and radiographic assessments. *J Foot Ankle Surg* 2004; **43**: 20-29 [PMID: 14752760 DOI: 10.1053/j.jfas.2003.11.005]
 - 25 **Michelson JD**, Varner KE, Checcone M. Diagnosing deltoid injury in ankle fractures: the gravity stress view. *Clin Orthop Relat Res* 2001; **(387)**: 178-182 [PMID: 11400880 DOI: 10.1097/00003086-200106000-00024]
 - 26 **Gill JB**, Risko T, Raducan V, Grimes JS, Schutt RC Jr. Comparison of manual and gravity stress radiographs for the evaluation of supination-external rotation fibular fractures. *J Bone Joint Surg Am* 2007; **89**: 994-999 [PMID: 17473136 DOI: 10.2106/JBJS.F.01002]
 - 27 **Schock HJ**, Pinzur M, Manion L, Stover M. The use of gravity or manual-stress radiographs in the assessment of supination-external rotation fractures of the ankle. *J Bone Joint Surg Br* 2007; **89**: 1055-1059 [PMID: 17785745 DOI: 10.1302/0301-620X.89B8.19134]
 - 28 **LeBa TB**, Gugala Z, Morris RP, Panchbhavi VK. Gravity vs manual external rotation stress view in evaluating ankle stability: a prospective study. *Foot Ankle Spec* 2015; **8**: 175-179 [PMID: 25552278 DOI: 10.1177/1938640014565048]
 - 29 **Dawe EJ**, Shafafy R, Quayle J, Gougoulas N, Wee A, Sakellariou A. The effect of different methods of stability assessment on fixation rate and complications in supination external rotation (SER) 2/4 ankle fractures. *Foot Ankle Surg* 2015; **21**: 86-90 [PMID: 25937406 DOI: 10.1016/j.fas.2014.09.010]
 - 30 **Hastie GR**, Akhtar S, Butt U, Baumann A, Barrie JL. Weightbearing Radiographs Facilitate Functional Treatment of Ankle Fractures of Uncertain Stability. *J Foot Ankle Surg* 2015; **54**: 1042-1046 [PMID: 26190779 DOI: 10.1053/j.jfas.2015.04.025]
 - 31 **Hoshino CM**, Nomoto EK, Norheim EP, Harris TG. Correlation of weightbearing radiographs and stability of stress positive ankle fractures. *Foot Ankle Int* 2012; **33**: 92-98 [PMID: 22381339 DOI: 10.3113/FAI.2012.0092]
 - 32 **Weber M**, Burmeister H, Flueckiger G, Krause FG. The use of weightbearing radiographs to assess the stability of supination-external rotation fractures of the ankle. *Arch Orthop Trauma Surg* 2010; **130**: 693-698 [PMID: 20082083 DOI: 10.1007/s00402-010-1051-1]
 - 33 **Nortunen S**, Lepojärvi S, Savola O, Niinimäki J, Ohtonen P, Flinkkilä T, Lantto I, Kortekangas T, Pakarinen H. Stability assessment of the ankle mortise in supination-external rotation-type ankle

- fractures: lack of additional diagnostic value of MRI. *J Bone Joint Surg Am* 2014; **96**: 1855-1862 [PMID: 25410502 DOI: 10.2106/JBJS.M.01533]
- 34 **Sadineni RT**, Pasumarthy A, Bellapa NC, Velicheti S. Imaging Patterns in MRI in Recent Bone Injuries Following Negative or Inconclusive Plain Radiographs. *J Clin Diagn Res* 2015; **9**: TC10-TC13 [PMID: 26557590 DOI: 10.7860/JCDR/2015/15451.6685]
- 35 **Crist B**, Fischer SJ, Dunbar RP. Ankle Fractures (Broken Ankle). [cited 27 August 2020]. In: American Academy of Orthopedic Surgeons [Internet]. Last reviewed: March 2013. Available from: <https://orthoinfo.aaos.org/en/diseases--conditions/ankle-fractures-broken-ankle/>
- 36 **Lauge-Hansen N**. Fractures of the ankle. III. Genetic roentgenologic diagnosis of fractures of the ankle. *Am J Roentgenol Radium Ther Nucl Med* 1954; **71**: 456-471 [PMID: 13124631]
- 37 **Danis R**. Les fractures malleolaires. In: Danis R. Théorie et pratique de l'ostéosynthèse. Paris: Masson, 1949: 133-165
- 38 **Fonseca LLD**, Nunes IG, Nogueira RR, Martins GEV, Mesencio AC, Kobata SI. Reproducibility of the Lauge-Hansen, Danis-Weber, and AO classifications for ankle fractures. *Rev Bras Ortop* 2018; **53**: 101-106 [PMID: 29367914 DOI: 10.1016/j.rboe.2017.11.013]
- 39 **Lampridis V**, Gougoulis N, Sakellariou A. Stability in ankle fractures: Diagnosis and treatment. *EFORT Open Rev* 2018; **3**: 294-303 [PMID: 29951269 DOI: 10.1302/2058-5241.3.170057]
- 40 **Broos PL**, Bisschop AP. Operative treatment of ankle fractures in adults: correlation between types of fracture and final results. *Injury* 1991; **22**: 403-406 [PMID: 1806506 DOI: 10.1016/0020-1383(91)90106-o]
- 41 **Bauer M**, Jonsson K, Nilsson B. Thirty-year follow-up of ankle fractures. *Acta Orthop Scand* 1985; **56**: 103-106 [PMID: 3925709 DOI: 10.3109/17453678508994329]
- 42 **Egol KA**, Amirtharajah M, Tejwani NC, Capla EL, Koval KJ. Ankle stress test for predicting the need for surgical fixation of isolated fibular fractures. *J Bone Joint Surg Am* 2004; **86**: 2393-2398 [PMID: 15523008 DOI: 10.2106/00004623-200411000-00005]
- 43 **Michelsen JD**, Ahn UM, Helgemo SL. Motion of the ankle in a simulated supination-external rotation fracture model. *J Bone Joint Surg Am* 1996; **78**: 1024-1031 [PMID: 8698719 DOI: 10.2106/00004623-199607000-00006]
- 44 **Clarke HJ**, Michelson JD, Cox QG, Jinnah RH. Tibio-talar stability in bimalleolar ankle fractures: a dynamic *in vitro* contact area study. *Foot Ankle* 1991; **11**: 222-227 [PMID: 1855708 DOI: 10.1177/107110079101100407]
- 45 **Curtis MJ**, Michelson JD, Urquhart MW, Byank RP, Jinnah RH. Tibiotalar contact and fibular malunion in ankle fractures. A cadaver study. *Acta Orthop Scand* 1992; **63**: 326-329 [PMID: 1609601 DOI: 10.3109/17453679209154793]
- 46 **Kimizuka M**, Kurosawa H, Fukubayashi T. Load-bearing pattern of the ankle joint. Contact area and pressure distribution. *Arch Orthop Trauma Surg* 1980; **96**: 45-49 [PMID: 7377925 DOI: 10.1007/BF01246141]
- 47 **Moody ML**, Koeneman J, Hettinger E, Karpman RR. The effects of fibular and talar displacement on joint contact areas about the ankle. *Orthop Rev* 1992; **21**: 741-744 [PMID: 1614720]
- 48 **Michelson JD**, Clarke HJ, Jinnah RH. The effect of loading on tibiotalar alignment in cadaver ankles. *Foot Ankle* 1990; **10**: 280-284 [PMID: 2111269 DOI: 10.1177/107110079001000507]
- 49 **Tornetta P 3rd**. Competence of the deltoid ligament in bimalleolar ankle fractures after medial malleolar fixation. *J Bone Joint Surg Am* 2000; **82**: 843-848 [PMID: 10859104 DOI: 10.2106/00004623-200006000-00011]
- 50 **Ramsey PL**, Hamilton W. Changes in tibiotalar area of contact caused by lateral talar shift. *J Bone Joint Surg Am* 1976; **58**: 356-357 [PMID: 1262367 DOI: 10.2106/00004623-197658030-00010]
- 51 **Riede UN**, Schenk RK, Willenegger H. [Joint mechanical studies on post-traumatic arthrosis in the ankle joint. I. The intra-articular model fracture]. *Langenbecks Arch Chir* 1971; **328**: 258-271 [PMID: 4996716 DOI: 10.1007/BF01251346]
- 52 **Ryd L**, Bengtsson S. Isolated fracture of the lateral malleolus requires no treatment. 49 prospective cases of supination-eversion type II ankle fractures. *Acta Orthop Scand* 1992; **63**: 443-446 [PMID: 1529699 DOI: 10.3109/17453679209154764]
- 53 **Zeegers AV**, Van Raay JJ, van der Werken C. Ankle fractures treated with a stabilizing shoe. *Acta Orthop Scand* 1989; **60**: 597-599 [PMID: 2603663 DOI: 10.3109/17453678909150130]
- 54 **Sanders DW**, Tieszer C, Corbett B; Canadian Orthopedic Trauma Society. Operative vs nonoperative treatment of unstable lateral malleolar fractures: a randomized multicenter trial. *J Orthop Trauma* 2012; **26**: 129-134 [PMID: 22330975 DOI: 10.1097/BOT.0b013e3182460837]
- 55 **DeAngelis NA**, Eskander MS, French BG. Does medial tenderness predict deep deltoid ligament incompetence in supination-external rotation type ankle fractures? *J Orthop Trauma* 2007; **21**: 244-247 [PMID: 17414551 DOI: 10.1097/BOT.0b013e3180413835]
- 56 **Lesic A**, Bumbasirevic M. Ankle fractures. *Curr Orthop* 2004; **18**: 232-244 [DOI: 10.1016/j.cuor.2004.03.001]
- 57 **Smith G**. The isolated lateral malleolar fracture: where are we and how did we get here? *Surgeon* 2013; **11**: 6-9 [PMID: 22459666 DOI: 10.1016/j.surge.2012.02.008]
- 58 **Michelson JD**, Magid D, Ney DR, Fishman EK. Examination of the pathologic anatomy of ankle fractures. *J Trauma* 1992; **32**: 65-70 [PMID: 1732577 DOI: 10.1097/00005373-199201000-00014]
- 59 **Harper MC**. The short oblique fracture of the distal fibula without medial injury: an assessment of displacement. *Foot Ankle Int* 1995; **16**: 181-186 [PMID: 7787973 DOI: 10.1177/107110079501600402]

- 60 **van den Bekerom MP**, van Dijk CN. Is fibular fracture displacement consistent with tibiotalar displacement? *Clin Orthop Relat Res* 2010; **468**: 969-974 [PMID: [19582527](#) DOI: [10.1007/s11999-009-0959-7](#)]
- 61 **Yde J**, Kristensen KD. Ankle fractures. Supination-eversion fractures stage II. Primary and late results of operative and non-operative treatment. *Acta Orthop Scand* 1980; **51**: 695-702 [PMID: [6778066](#) DOI: [10.3109/17453678008990863](#)]
- 62 **Kristensen KD**, Hansen T. Closed treatment of ankle fractures. Stage II supination-eversion fractures followed for 20 years. *Acta Orthop Scand* 1985; **56**: 107-109 [PMID: [3925710](#) DOI: [10.3109/17453678508994330](#)]
- 63 **Haraguchi N**, Toga H, Shiba N, Kato F. Avulsion fracture of the lateral ankle ligament complex in severe inversion injury: incidence and clinical outcome. *Am J Sports Med* 2007; **35**: 1144-1152 [PMID: [17379919](#) DOI: [10.1177/0363546507299531](#)]
- 64 **Donken CC**, Al-Khateeb H, Verhofstad MH, van Laarhoven CJ. Surgical vs conservative interventions for treating ankle fractures in adults. *Cochrane Database Syst Rev* 2012; CD008470 [PMID: [22895975](#) DOI: [10.1002/14651858.CD008470.pub2](#)]
- 65 **Dobbe A**, Beaupre LA, Ali Almansoori K. Functional Outcomes of Isolated Infrasyndesmotom Fibula Fractures. *Foot Ankle Int* 2020; **5**: 1-9 [DOI: [10.1177/2473011419892227](#)]
- 66 **Yap RY**, Babel A, Phoon KM, Ward AE. Functional Outcomes Following Operative and Nonoperative Management of Weber C Ankle Fractures: A Systematic Review. *J Foot Ankle Surg* 2020; **59**: 105-111 [PMID: [31882132](#) DOI: [10.1053/j.jfas.2019.06.005](#)]
- 67 **Donken CC**, Verhofstad MH, Edwards MJ, van Laarhoven CJ. Twenty-two-year follow-up of pronation external rotation type III-IV (OTA type C) ankle fractures: a retrospective cohort study. *J Orthop Trauma* 2012; **26**: e115-e122 [PMID: [22198655](#) DOI: [10.1097/BOT.0b013e31823bc320](#)]
- 68 **McKibbin B**. The biology of fracture healing in long bones. *J Bone Joint Surg Br* 1978; **60-B**: 150-162 [PMID: [350882](#) DOI: [10.1302/0301-620X.60B2.350882](#)]
- 69 **Patil S**, Gandhi J, Curzon I, Hui AC. Incidence of deep-vein thrombosis in patients with fractures of the ankle treated in a plaster cast. *J Bone Joint Surg Br* 2007; **89**: 1340-1343 [PMID: [17957074](#) DOI: [10.1302/0301-620X.89B10.19241](#)]
- 70 **Drakos MC**, Behrens SB, Paller D, Murphy C, DiGiovanni CW. Biomechanical Comparison of an Open vs Arthroscopic Approach for Lateral Ankle Instability. *Foot Ankle Int* 2014; **35**: 809-815 [PMID: [24850160](#) DOI: [10.1177/1071100714535765](#)]
- 71 **Stuart PR**, Brumby C, Smith SR. Comparative study of functional bracing and plaster cast treatment of stable lateral malleolar fractures. *Injury* 1989; **20**: 323-326 [PMID: [2516838](#) DOI: [10.1016/0020-1383\(89\)90003-x](#)]
- 72 **Heijnders IL**, Lin CW. Treatment of acute ankle sprains: evidence on the use of an ankle brace is unclear. *Br J Sports Med* 2012; **46**: 852-853 [PMID: [22844037](#) DOI: [10.1136/bjsports-2012-091533](#)]
- 73 **Moseley AM**, Beckenkamp PR, Haas M, Herbert RD, Lin CW; EXACT Team. Rehabilitation After Immobilization for Ankle Fracture: The EXACT Randomized Clinical Trial. *JAMA* 2015; **314**: 1376-1385 [PMID: [26441182](#) DOI: [10.1001/jama.2015.12180](#)]
- 74 **Kortekangas T**, Haapasalo H, Flinkkilä T, Ohtonen P, Nortunen S, Laine HJ, Järvinen TL, Pakarinen H. Three week vs six week immobilisation for stable Weber B type ankle fractures: randomised, multicentre, non-inferiority clinical trial. *BMJ* 2019; **364**: k5432 [PMID: [30674451](#) DOI: [10.1136/bmj.k5432](#)]
- 75 **Stover CN**. A Functional Semirigid Support System for Ankle Injuries. *Phys Sportsmed* 1979; **7**: 71-78 [PMID: [27448078](#) DOI: [10.1080/00913847.1979.11710861](#)]
- 76 **Brink O**, Staunstrup H, Sommer J. Stable lateral malleolar fractures treated with aircast ankle brace and DonJoy R.O.M.-Walker brace: a prospective randomized study. *Foot Ankle Int* 1996; **17**: 679-684 [PMID: [8946182](#) DOI: [10.1177/107110079601701106](#)]
- 77 **Port AM**, McVie JL, Naylor G, Kreibich DN. Comparison of two conservative methods of treating an isolated fracture of the lateral malleolus. *J Bone Joint Surg Br* 1996; **78**: 568-572 [PMID: [8682822](#) DOI: [10.1302/0301-620X.78B4.0780568](#)]
- 78 **van den Berg C**, Haak T, Weil NL, Hoogendoorn JM. Functional bracing treatment for stable type B ankle fractures. *Injury* 2018; **49**: 1607-1611 [PMID: [29903578](#) DOI: [10.1016/j.injury.2018.06.009](#)]
- 79 **Huang Z**, Liu L, Tu C, Zhang H, Fang Y, Yang T, Pei F. Comparison of three plate system for lateral malleolar fixation. *BMC Musculoskelet Disord* 2014; **15**: 360 [PMID: [25358474](#) DOI: [10.1186/1471-2474-15-360](#)]
- 80 **Minihane KP**, Lee C, Ahn C, Zhang LQ, Merk BR. Comparison of lateral locking plate and antiglide plate for fixation of distal fibular fractures in osteoporotic bone: a biomechanical study. *J Orthop Trauma* 2006; **20**: 562-566 [PMID: [16990728](#) DOI: [10.1097/01.bot.0000245684.96775.82](#)]
- 81 **Knutsen AR**, Sangiorgio SN, Liu C, Zhou S, Warganich T, Fleming J, Harris TG, Ebramzadeh E. Distal fibula fracture fixation: Biomechanical evaluation of three different fixation implants. *Foot Ankle Surg* 2016; **22**: 278-285 [PMID: [27810029](#) DOI: [10.1016/j.fas.2016.08.007](#)]
- 82 **Anglen J**, Kyle RF, Marsh JL, Virkus WW, Watters WC 3rd, Keith MW, Turkelson CM, Wies JL, Boyer KM. Locking plates for extremity fractures. *J Am Acad Orthop Surg* 2009; **17**: 465-472 [PMID: [19571302](#) DOI: [10.5435/00124635-200907000-00007](#)]
- 83 **Bilgetekin YG**, Çatma MF, Öztürk A, Ünlü S, Ersan Ö. Comparison of different locking plate fixation methods in lateral malleolus fractures. *Foot Ankle Surg* 2019; **25**: 366-370 [PMID: [30321977](#) DOI: [10.1016/j.fas.2018.01.004](#)]

- 84 **Tsukada S**, Otsuji M, Shiozaki A, Yamamoto A, Komatsu S, Yoshimura H, Ikeda H, Hoshino A. Locking vs non-locking neutralization plates for treatment of lateral malleolar fractures: a randomized controlled trial. *Int Orthop* 2013; **37**: 2451-2456 [PMID: 24077867 DOI: 10.1007/s00264-013-2109-9]
- 85 **Eckel TT**, Glisson RR, Anand P, Parekh SG. Biomechanical comparison of 4 different lateral plate constructs for distal fibula fractures. *Foot Ankle Int* 2013; **34**: 1588-1595 [PMID: 23818460 DOI: 10.1177/1071100713496223]
- 86 **Herrera-Pérez M**, Gutiérrez-Morales MJ, Guerra-Ferraz A, Pais-Brito JL, Boluda-Mengod J, Garcés GL. Locking vs non-locking one-third tubular plates for treating osteoporotic distal fibula fractures: a comparative study. *Injury* 2017; **48** Suppl 6: S60-S65 [PMID: 29162244 DOI: 10.1016/S0020-1383(17)30796-9]
- 87 **Miller AG**, Margules A, Raikin SM. Risk factors for wound complications after ankle fracture surgery. *J Bone Joint Surg Am* 2012; **94**: 2047-2052 [PMID: 23172322 DOI: 10.2106/JBJS.K.01088]
- 88 **SooHoo NF**, Krenk L, Eagan MJ, Gurbani B, Ko CY, Zingmond DS. Complication rates following open reduction and internal fixation of ankle fractures. *J Bone Joint Surg Am* 2009; **91**: 1042-1049 [PMID: 19411451 DOI: 10.2106/JBJS.H.00653]
- 89 **Varenne Y**, Curado J, Asloum Y, Salle de Chou E, Colin F, Gouin F. Analysis of risk factors of the postoperative complications of surgical treatment of ankle fractures in the elderly: A series of 477 patients. *Orthop Traumatol Surg Res* 2016; **102**: S245-S248 [PMID: 27033839 DOI: 10.1016/j.otsr.2016.03.004]
- 90 **Lanzetti RM**, Lupariello D, Venditto T, Guzzini M, Ponzo A, De Carli A, Ferretti A. The role of diabetes mellitus and BMI in the surgical treatment of ankle fractures. *Diabetes Metab Res Rev* 2018; **34** [PMID: 29031012 DOI: 10.1002/dmrr.2954]
- 91 **McLennan JG**, Ungersma JA. A new approach to the treatment of ankle fractures. The Inyo nail. *Clin Orthop Relat Res* 1986; **(213)**: 125-136 [PMID: 3780081 DOI: 10.1097/00003086-198612000-00015]
- 92 **Jain S**, Haughton BA, Brew C. Intramedullary fixation of distal fibular fractures: a systematic review of clinical and functional outcomes. *J Orthop Traumatol* 2014; **15**: 245-254 [PMID: 25304004 DOI: 10.1007/s10195-014-0320-0]
- 93 **Coifman O**, Bariteau JT, Shazar N, Tenenbaum SA. Lateral malleolus closed reduction and internal fixation with intramedullary fibular rod using minimal invasive approach for the treatment of ankle fractures. *Foot Ankle Surg* 2019; **25**: 79-83 [PMID: 29409300 DOI: 10.1016/j.fas.2017.08.008]
- 94 **Rehman H**, Gardner WT, Rankin I, Johnstone AJ. The implants used for intramedullary fixation of distal fibula fractures: A review of literature. *Int J Surg* 2018; **56**: 294-300 [PMID: 29964180 DOI: 10.1016/j.ijsu.2018.06.019]
- 95 **Bugler KE**, Watson CD, Hardie AR, Appleton P, McQueen MM, Court-Brown CM, White TO. The treatment of unstable fractures of the ankle using the Acumed fibular nail: development of a technique. *J Bone Joint Surg Br* 2012; **94**: 1107-1112 [PMID: 22844054 DOI: 10.1302/0301-620X.94B8.28620]
- 96 **Asloum Y**, Bedin B, Roger T, Charissoux JL, Arnaud JP, Mabit C. Internal fixation of the fibula in ankle fractures: a prospective, randomized and comparative study: plating vs nailing. *Orthop Traumatol Surg Res* 2014; **100**: S255-S259 [PMID: 24709304 DOI: 10.1016/j.otsr.2014.03.005]
- 97 **White TO**, Bugler KE, Appleton P, Will E, McQueen MM, Court-Brown CM. A prospective randomised controlled trial of the fibular nail vs standard open reduction and internal fixation for fixation of ankle fractures in elderly patients. *Bone Joint J* 2016; **98-B**: 1248-1252 [PMID: 27587528 DOI: 10.1302/0301-620X.98B9.35837]
- 98 **Al-Obaidi B**, Wiik AV, Bhattacharyya R, Mushtaq N, Bhattacharya R. Fibular nails for open and closed ankle fractures: Results from a non-designer level I major trauma centre. *J Orthop Surg (Hong Kong)* 2019; **27**: 2309499019832420 [PMID: 30803314 DOI: 10.1177/2309499019832420]
- 99 **Sneppen O**. Pseudarthrosis of the lateral malleolus. *Acta Orthop Scand* 1971; **42**: 187-200 [PMID: 5000187 DOI: 10.3109/17453677108989039]
- 100 **Siliski J**, Blitzer C, Healy W, Baumgaertner M, Carr C. Non-union of the fibula after ankle fracture. *Orthop Trans* 1993; **18**: 22 [DOI: 10.1097/00005131-199304000-00100]
- 101 **Walsh EF**, DiGiovanni C. Fibular nonunion after closed rotational ankle fracture. *Foot Ankle Int* 2004; **25**: 488-495 [PMID: 15319107 DOI: 10.1177/107110070402500708]
- 102 **Böstman O**, Kyrö A. Delayed union of fibular fractures accompanying fractures of the tibial shaft. *J Trauma* 1991; **31**: 99-102 [PMID: 1986140 DOI: 10.1097/00005373-199101000-00018]
- 103 **König M**, Gotzen L. [Pseudarthroses of the fibula following fracture of the lower leg]. *Unfallchirurg* 1989; **92**: 191-194 [PMID: 2727714]
- 104 **Marti RK**, Raaymakers EL, Nolte PA. Malunited ankle fractures. The late results of reconstruction. *J Bone Joint Surg Br* 1990; **72**: 709-713 [PMID: 2116416 DOI: 10.1302/0301-620X.72B4.2116416]
- 105 **Lübbecke A**, Salvo D, Stern R, Hoffmeyer P, Holzer N, Assal M. Risk factors for post-traumatic osteoarthritis of the ankle: an eighteen year follow-up study. *Int Orthop* 2012; **36**: 1403-1410 [PMID: 22249843 DOI: 10.1007/s00264-011-1472-7]
- 106 **Ovaska M**. Complications in ankle fracture surgery. *Acta Orthop Suppl* 2015; **86**: 1-32 [PMID: 25586467 DOI: 10.3109/17453674.2014.100227]
- 107 **Weber D**, Borisch N, Weber M. Treatment of malunion in ankle fractures. *Eur J Trauma Emerg Surg* 2010; **36**: 521-524 [PMID: 26816306 DOI: 10.1007/s00068-010-0060-7]

- 108 **Yablon IG**, Leach RE. Reconstruction of malunited fractures of the lateral malleolus. *J Bone Joint Surg Am* 1989; **71**: 521-527 [PMID: 2703512 DOI: 10.2106/00004623-198971040-00007]
- 109 **Ward AJ**, Ackroyd CE, Baker AS. Late lengthening of the fibula for malaligned ankle fractures. *J Bone Joint Surg Br* 1990; **72**: 714-717 [PMID: 2116417 DOI: 10.1302/0301-620X.72B4.2116417]
- 110 **Schepers T**, De Vries MR, Van Lieshout EM, Van der Elst M. The timing of ankle fracture surgery and the effect on infectious complications; a case series and systematic review of the literature. *Int Orthop* 2013; **37**: 489-494 [PMID: 23288046 DOI: 10.1007/s00264-012-1753-9]
- 111 **Bäcker HC**, Greisberg JK, Vosseller JT. Fibular Plate Fixation and Correlated Short-term Complications. *Foot Ankle Spec* 2020; **13**: 378-382 [PMID: 31538819 DOI: 10.1177/1938640019873539]
- 112 **Anderson SA**, Li X, Franklin P, Wixted JJ. Ankle fractures in the elderly: initial and long-term outcomes. *Foot Ankle Int* 2008; **29**: 1184-1188 [PMID: 19138481 DOI: 10.3113/FAI.2008.1184]
- 113 **Zaghloul A**, Haddad B, Barksfield R, Davis B. Early complications of surgery in operative treatment of ankle fractures in those over 60: a review of 186 cases. *Injury* 2014; **45**: 780-783 [PMID: 24388418 DOI: 10.1016/j.injury.2013.11.008]
- 114 **Naumann MG**, Sigurdson U, Utvåg SE, Stavem K. Incidence and risk factors for removal of an internal fixation following surgery for ankle fracture: A retrospective cohort study of 997 patients. *Injury* 2016; **47**: 1783-1788 [PMID: 27262772 DOI: 10.1016/j.injury.2016.05.011]
- 115 **Lyle SA**, Malik C, Oddy MJ. Comparison of Locking Versus Nonlocking Plates for Distal Fibula Fractures. *J Foot Ankle Surg* 2018; **57**: 664-667 [PMID: 29681437 DOI: 10.1053/j.jfas.2017.11.035]
- 116 **Koval KJ**, Zhou W, Sparks MJ, Cantu RV, Hecht P, Lurie J. Complications after ankle fracture in elderly patients. *Foot Ankle Int* 2007; **28**: 1249-1255 [PMID: 18173987 DOI: 10.3113/FAI.2007.1249]
- 117 **Lynde MJ**, Sautter T, Hamilton GA, Schubert JM. Complications after open reduction and internal fixation of ankle fractures in the elderly. *Foot Ankle Surg* 2012; **18**: 103-107 [PMID: 22443995 DOI: 10.1016/j.fas.2011.03.010]
- 118 **Flynn JM**, Rodriguez-del Rio F, Pizá PA. Closed ankle fractures in the diabetic patient. *Foot Ankle Int* 2000; **21**: 311-319 [PMID: 10808971 DOI: 10.1177/107110070002100407]
- 119 **Schepers T**, Van Lieshout EM, De Vries MR, Van der Elst M. Increased rates of wound complications with locking plates in distal fibular fractures. *Injury* 2011; **42**: 1125-1129 [PMID: 21329921 DOI: 10.1016/j.injury.2011.01.009]
- 120 **Lin CW**, Donkers NA, Refshauge KM, Beckenkamp PR, Khera K, Moseley AM. Rehabilitation for ankle fractures in adults. *Cochrane Database Syst Rev* 2012; **11**: CD005595 [PMID: 23152232 DOI: 10.1002/14651858.CD005595.pub3]
- 121 **Saltzman CL**, Salamon ML, Blanchard GM, Huff T, Hayes A, Buckwalter JA, Amendola A. Epidemiology of ankle arthritis: report of a consecutive series of 639 patients from a tertiary orthopaedic center. *Iowa Orthop J* 2005; **25**: 44-46 [PMID: 16089071]
- 122 **Martin RL**, Stewart GW, Conti SF. Posttraumatic ankle arthritis: an update on conservative and surgical management. *J Orthop Sports Phys Ther* 2007; **37**: 253-259 [PMID: 17549954 DOI: 10.2519/jospt.2007.2404]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

