

Microsoft Bing

国内版

国际版

Progressive liver injury and increased mortality risk in COVID-19 pat

Chat with Bing

Sign in

ALLIMAGESVIDEOS

221,000 ResultsAny time

Fatty liver, cirrhosis double mortality risk in COVID-19

[https://www.healio.com/news/hepatology/20200828/...](https://www.healio.com/news/hepatology/20200828/)

Aug 28, 2020 · In this registry of more than 1,000 patients with liver disease, Berg showed that non-cirrhotic liver disease did not confer an additional risk with COVID-19 but once a patient ...

Liver injury in COVID-19: management and challenges - The ...

[https://www.thelancet.com/journals/langas/article/...](https://www.thelancet.com/journals/langas/article/)

The number of confirmed cases of 2019 coronavirus disease (COVID-19) has reached 87 137 worldwide as of March 1, 2020, according to WHO COVID-19 situation report 41; most of these patients are in Wuhan, China. Many cases of COVID-19 are acute and resolve quickly, but the disease can also be fatal, with a mortality rate of around 3%.

Cited by: 569Author: Chao Zhang, Lei Shi, Fu-Sheng Wang

Publish Year: 2020

Abnormal Liver Tests in COVID-19: A Retrospective ...

<https://aasldpubs.onlinelibrary.wiley.com/doi/10.1002/hep.31487>

Jul 29, 2020 · Furthermore, although studies from China have reported associations between abnormal liver tests and hospitalization length of stay, risk for progression to severe COVID-19, (16-18) and mortality, there are limited data about these relationships in U.S. cohorts.

Search Tools

Turn off Hover Translation (关闭翻译)

国内版 国际版

Progressive liver injury and increased mortality risk in COVID-19 pat



ALL IMAGES VIDEOS

227,000 Results

Any time ▼

Progressive liver injury and increased mortality risk in ...

<https://www.authorea.com/users/342559/articles/...> ▼

A document by Shuisheng Zhang, written on Authorea. Clinical characteristic and risk factors

Acute Liver Injury in COVID-19: Prevalence and Association ...

<https://aasldpubs.onlinelibrary.wiley.com/doi/abs/10.1002/hep.31404>

May 30, 2020 · We aim to identify the prevalence of and risk factors for development of COVID-19-associated ALI in a large cohort in the United States. Approach and Results. In this retrospective cohort study, all patients who underwent SARS-CoV-2 testing at three hospitals in the NewYork-Presbyterian network were assessed.

Cited by: 57

Author: Meaghan M. Phipps, Luis H. Barraza, Elijah ...

Publish Year: 2020

Clinical characteristics and risk factors of liver injury ...

<https://pubmed.ncbi.nlm.nih.gov/33026573>

Background: Coronavirus disease 2019 (COVID-19) has rapidly become a major international public health concern. This study was designed to evaluate the clinical characteristics and risk factors of COVID-19-associated liver injury. Methods: A fraction of 657 COVID-19 patients were retrospectively analyzed. Clinical and laboratory data were derived from electronic medical records and compared ...

Fatty liver, cirrhosis double mortality risk in COVID-19

<https://www.healio.com/news/hepatology/20200828/...> ▼

Aug 28, 2020 · In this registry of more than 1,000 patients with liver disease, Berg showed that non-cirrhotic liver disease did not confer an additional risk with COVID-19 but once a patient ...

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 61015

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Progressive liver injury and increased mortality risk in COVID-19 patients: A retrospective cohort study in China

Liver injury in COVID-19

Abstract

BACKGROUND

Liver injury is common and also can be fatal, particularly in severe or critical patients with coronavirus disease 2019 (COVID-19).

AIM

We aim to conduct an in-depth investigation into the risk factors of liver injury and into

Match Overview

1	Crossref 116 words Caiting Yang, Xiaoxia Ma, Jili Wu, Jie Han, Zhe Zheng, Huipeng Duan, Qun Liu, Changxin Wu, Yongkang Dong, Li Do	3%
2	Crossref Posted Content 66 words Weiping Jiang, Xuanqi Liu, Xiaodan Tang, Xiaoxiao Ge et al. I. "Cardiac Troponin T as the Underlying Risk Factor Assr ...	2%
3	Crossref 46 words Zhenyu Fan, Liping Chen, Jun Li, Xin Cheng, Jingmao Yang, Cheng Tian, Yajun Zhang, Shaoping Huang, Zhanju Li ...	1%
4	Crossref 37 words "Clinical characteristics and outcomes in coronavirus disease 2019 (COVID-19) patients with and without hypertens ...	1%
5	Internet 33 words crawled on 23-Oct-2020 onlinelibrary.wiley.com	1%
6	Internet 25 words crawled on 09-Nov-2020 www.researchsquare.com	1%
7	Internet 21 words crawled on 16-Oct-2019 www.nature.com	<1%
8	Internet 18 words crawled on 12-Aug-2020 www.researchgate.net	<1%

国内版 国际版

Progressive liver injury and increased mortality risk in COVID-19 pat



ALL IMAGES VIDEOS

237,000 Results Any time ▾

Acute Liver Injury in COVID-19: Prevalence and Association ...

<https://aasldpubs.onlinelibrary.wiley.com/doi/abs/10.1002/hep.31404>

May 30, 2020 · We aim to identify the prevalence of and risk factors for development of COVID-19-associated ALI in a large cohort in the United States. Approach and Results. In this retrospective cohort study, all patients who underwent SARS-CoV-2 testing at three hospitals in the NewYork-Presbyterian network were assessed.

Cited by: 57 Author: Meaghan M. Phipps, Luis H. Barraza, Elij...
Publish Year: 2020

Fatty liver, cirrhosis double mortality risk in COVID-19

<https://www.healio.com/news/hepatology/20200828/...> ▾

Aug 28, 2020 · In this registry of more than 1,000 patients with liver disease, Berg showed that non-cirrhotic liver disease did not confer an additional risk with COVID-19 but once a patient ...

Liver injury in COVID-19: management and challenges - The ...

[https://www.thelancet.com/journals/langas/article/PIIS2468-1253\(20\)30057-1/fulltext](https://www.thelancet.com/journals/langas/article/PIIS2468-1253(20)30057-1/fulltext)

The number of confirmed cases of 2019 coronavirus disease (COVID-19) has reached 87 137 worldwide as of March 1, 2020, according to WHO COVID-19 situation report 41; most of these patients are in Wuhan, China. Many cases of COVID-19 are acute and resolve quickly, but the disease can also be fatal, with a mortality rate of around 3%.

Cited by: 762 Author: Chao Zhang, Lei Shi, Fu-Sheng Wang
Publish Year: 2020

Liver injury in COVID-19: The current evidence - Saleh A ...

<https://journals.sagepub.com/doi/full/10.1177/2050640620924157>

May 26, 2020 · In SARS, the highest mortality rates were observed in the elderly and adults with underlying liver disease. 45 Therefore, it has to be expected that the patients with COVID-19 are also more vulnerable to hepatic injury. 9 In a case series from the Zhejiang province, a prevalence of 11% of underlying liver disease was reported.

Cited by: 25 Author: Saleh A Alqahtani, Jörn M Schattenberg
Publish Year: 2020

Abnormal Liver Tests in COVID-19: A Retrospective ...