

ALL IMAGES VIDEOS

35,200 Results Any time ▾

Fibrinogen/Albumin Ratio Index Is an Independent Prognosis ...

<https://europepmc.org/article/PMC/PMC7462001>

Moreover, multivariate analysis indicated that preoperative FARI is an independent predictor of poor prognosis in esophageal carcinoma patients. In another study, involving 154 gallbladder cancer (GBC) patients, the findings revealed that patients with a FARI of more than 0.09 have a worse OS compared with patients with a FARI ≤ 0.08 (26).

(PDF) Nutritional assessment and prognosis of oral cancer ...

<https://www.researchgate.net/publication/339436909>

Results: Patients with BMI < 18.5 kg/m2 (VS 18.5 kg/m2 ≤ BMI < 24 kg/m2) had a poor survival outcome (HR = 1.585; 95% CI: 1.207-2.082). ... was an independent predictor of a poor prognosis in ...

Implications of BMI for the Prognosis of Gastric Cancer ...

<https://www.researchgate.net/publication/283492845>

The patients were principally divided into 3 groups on the basis of BMI: low (<18.5 kg/m2), normal (≥18.5-25.0 kg/m2) and high (≥25.0 kg/m2). Results: The low-BMI patients had more advanced ...

Clinical characteristics and survival of 5283 esophageal ...

<https://academic.oup.com/ajco/advance-article-abstract/doi/10.1093/ajco/kp212/5283>

Search Tools

Turn off Hover Translation (关闭浮词)

ALL IMAGES VIDEOS

51,200 Results Any time Open links in new tab

Impact of body mass index on long-term survival outcome in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6160684>

Purpose: Being elevated **body mass index** (BMI) has been considered a **poor** prognostic factor in **patients with cancer**. However, studies about the impact of elevated BMI on the survival outcome after cancer surgery have conflicting results. This study aimed to evaluate the impact of BMI on long-term postoperative survival outcome in a large cohort of Asian population with solid ...

Cited by: 2 Author: Chia-Yen Hung, Cheng-Chou Lai, Ping-T...

Publish Year: 2018

Use of Body Mass Index to Predict the Prognosis of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5502225>

Gastric cancer is one of the most common malignancies and death from **gastric cancer** ranks second among all cancer deaths worldwide. 1 Remnant **gastric cancer** (RGC) is a less prevalent form of the disease, and the incidence of RGC has been reported as 1–7% of all **gastric cancer** cases. 2-4 The most frequently used definition of RGC is **cancer** in the remnant stomach after ...

Cited by: 2 Author: Tomoyuki Matsunaga, Hiroaki Saito, To...

Publish Year: 2017

Prognostic impact of body mass index stratified by smoking ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5502225>

Search Tools

Turn off Hover Translation (关闭取词)

激活 Windows
转到“设置”以激活 Windows。

Low BMI (<18.5 kg/m²) is an independent predictor of poor long-term prognosis among patients with resectable gastric cancer: a retrospective study at Chinese National Cancer Center

Shuai Ma, Hao Liu, Fu-Hai Ma, Yang Li, Peng Jin, Hai-Tao Hu, Wen-Zhe Kang, Wei-Kun Li, Jian-Ping Xiong, Yan-Tao Tian

| Match Overview | | |
|----------------|---|----|
| 1 | Internet 137 words crawled on 16-Feb-2020 #publishing blob core windows.net | 4% |
| 2 | Internet 68 words crawled on 17-Jul-2019 www.jcancer.org | 2% |
| 3 | Internet 64 words crawled on 21-Jul-2020 www.wjgnet.com | 2% |
| 4 | Internet 32 words crawled on 09-Jan-2018 ascipubs.org | 1% |
| 5 | Internet 32 words crawled on 07-Dec-2020 synapse.koreamed.org | 1% |
| 6 | Internet 26 words crawled on 12-May-2020 bmccancer.biomedcentral.com | 1% |

国内版 国际版

Low body mass index is an independent predictor of poor long-term



ALL IMAGES VIDEOS

50,200 Results Any time ▾

Use of Body Mass Index to Predict the Prognosis of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5502225>

Gastric cancer is one of the most common malignancies and death from gastric cancer ranks second among all cancer deaths worldwide. 1 Remnant gastric cancer (RGC) is a less prevalent form of the disease, and the incidence of RGC has been reported as 1–7% of all gastric cancer cases. 2–4 The most frequently used definition of RGC is cancer in the remnant stomach after surgery for benign ...

Cited by: 2 Author: Tomoyuki Matsunaga, Hiroaki Saito, Tom...

Publish Year: 2017

Prognostic impact of body mass index stratified by smoking ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5074733>

French oncologists found that a low BMI (<18 kg/m²) was associated with inferior survival in 87 ESCC cases with definitive chemoradiotherapy,³⁰ which was consistent with data from another French study reported by Clavier et al.³¹ The poor prognosis of low BMI was also identified in two other Chinese studies.^{28,29} Consistent with these results, we also confirmed, in our large retrospective study, the ...

Cited by: 4 Author: Peng Sun, Fei Zhang, Cui Chen, Chao Re...

Publish Year: 2016

A Novel Prognostic Score, Based on Preoperative ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5118680>

Oct 25, 2016 · Introduction. Gastric cancer (GC) is one of the most common malignancies worldwide, accounting for approximately one million new cases annually 1, 2. Despite great advancements in early detection, surgical techniques, and multidisciplinary treatment in recent years, the postoperative long-term survival of patients is still poor 3. Therefore, there is increasing interest in prognostic factors to ...

Cited by: 10 Author: Xuechao Liu, Haibo Qiu, Jianjun Liu, Sha...

Publish Year: 2016

A Novel Inflammation- and Nutrition-Based Prognostic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5033418>

Increased RDW is associated with poor prognosis in cancer patients. According to WHO, there are three weight categories: underweight (BMI < 18.5), normal weight (18.5 ≤ BMI < 25) and overweight (BMI ≥ 25). Low BMI values are associated with increased risk of cancer mortality in patients.

Cited by: 7 Author: Yan Fu, Yize Mao, Shiqi Chen, Ankui Ya...

Publish Year: 2016