

# World Journal of *Clinical Cases*

*World J Clin Cases* 2021 March 16; 9(8): 1761-2021



**REVIEW**

- 1761 Cardiac rehabilitation and its essential role in the secondary prevention of cardiovascular diseases  
*Winnige P, Vysoky R, Dosbaba F, Batalik L*

**ORIGINAL ARTICLE****Case Control Study**

- 1785 Association between homeobox protein transcript antisense intergenic ribonucleic acid genetic polymorphisms and cholangiocarcinoma  
*Lampropoulou DI, Laschos K, Aravantinos G, Georgiou K, Papiris K, Theodoropoulos G, Gazouli M, Filippou D*

**Retrospective Study**

- 1793 Risk factors for post-hepatectomy liver failure in 80 patients  
*Xing Y, Liu ZR, Yu W, Zhang HY, Song MM*

- 1803 Outcomes of laparoscopic bile duct exploration for choledocholithiasis with small common bile duct  
*Huang XX, Wu JY, Bai YN, Wu JY, Lv JH, Chen WZ, Huang LM, Huang RF, Yan ML*

**Observational Study**

- 1814 Three-dimensional finite element analysis with different internal fixation methods through the anterior approach  
*Xie XJ, Cao SL, Tong K, Zhong ZY, Wang G*
- 1827 Bedside cardiopulmonary ultrasonography evaluates lung water content in very low-weight preterm neonates with patent ductus arteriosus  
*Yu LF, Xu CK, Zhao M, Niu L, Huang XM, Zhang ZQ*

**CASE REPORT**

- 1835 Conservative endodontic management using a calcium silicate bioceramic sealer for delayed root fracture: A case report and review of the literature  
*Zheng P, Shen ZY, Fu BP*
- 1844 Brain magnetic resonance imaging findings and radiologic review of maple syrup urine disease: Report of three cases  
*Li Y, Liu X, Duan CF, Song XF, Zhuang XH*
- 1853 A three-year clinical investigation of a Chinese child with craniometaphyseal dysplasia caused by a mutated ANKH gene  
*Wu JL, Li XL, Chen SM, Lan XP, Chen JJ, Li XY, Wang W*
- 1863 Intradural osteomas: Report of two cases  
*Li L, Ying GY, Tang YJ, Wu H*

- 1871** Gastroesophageal varices in a patient presenting with essential thrombocythemia: A case report  
*Wang JB, Gao Y, Liu JW, Dai MG, Yang SW, Ye B*
- 1877** Chest pain showing precordial ST-segment elevation in a 96-year-old woman with right coronary artery occlusion: A case report  
*Wu HY, Cheng G, Cao YW*
- 1885** Subcutaneous panniculitis-like T-cell lymphoma invading central nervous system in long-term clinical remission with lenalidomide: A case report  
*Sun J, Ma XS, Qu LM, Song XS*
- 1893** Imaging findings of primary pulmonary synovial sarcoma with secondary distant metastases: A case report  
*Li R, Teng X, Han WH, Li Y, Liu QW*
- 1901** Severe community-acquired pneumonia caused by *Leptospira interrogans*: A case report and review of literature  
*Bao QH, Yu L, Ding JJ, Chen YJ, Wang JW, Pang JM, Jin Q*
- 1909** Bilateral common peroneal neuropathy due to rapid and marked weight loss after biliary surgery: A case report  
*Oh MW, Gu MS, Kong HH*
- 1916** Retroperitoneal laparoscopic partial resection of the renal pelvis for urothelial carcinoma: A case report  
*Wang YL, Zhang HL, Du H, Wang W, Gao HF, Yu GH, Ren Y*
- 1923** 17 $\alpha$ -hydroxylase/17,20 carbon chain lyase deficiency caused by p.Tyr329fs homozygous mutation: Three case reports  
*Zhang D, Sun JR, Xu J, Xing Y, Zheng M, Ye SD, Zhu J*
- 1931** Epithelioid angiomyolipoma of the pancreas: A case report and review of the literature  
*Zhu QQ, Niu ZF, Yu FD, Wu Y, Wang GB*
- 1940** Computed tomography imaging features for amyloid dacryolith in the nasolacrimal excretory system: A case report  
*Che ZG, Ni T, Wang ZC, Wang DW*
- 1946** Epidural analgesia followed by epidural hydroxyethyl starch prevented post-dural puncture headache: Twenty case reports and a review of the literature  
*Song LL, Zhou Y, Geng ZY*
- 1953** Extracorporeal membrane oxygenation for coronavirus disease 2019-associated acute respiratory distress syndrome: Report of two cases and review of the literature  
*Wen JL, Sun QZ, Cheng Z, Liao XZ, Wang LQ, Yuan Y, Li JW, Hou LS, Gao WJ, Wang WJ, Soh WY, Li BF, Ma DQ*
- 1968** Human parvovirus B19-associated early postoperative acquired pure red cell aplasia in simultaneous pancreas-kidney transplantation: A case report  
*Wang H, Fu YX, Song WL, Wang Z, Feng G, Zhao J, Nian YQ, Cao Y*

- 1976** Diabetes insipidus with impaired vision caused by germinoma and perioptic meningeal seeding: A case report  
*Yang N, Zhu HJ, Yao Y, He LY, Li YX, You H, Zhang HB*
- 1983** Madelung disease: A case report  
*Chen KK, Ni LS, Yu WH*
- 1989** Laryngopharyngeal reflux disease management for recurrent laryngeal contact granuloma: A case report  
*Li K, Chen WY, Li YY, Wang TL, Tan MJ, Chen Z, Chen H*
- 1996** *Mycobacterium abscessus* infection after facial injection of argireline: A case report  
*Chen CF, Liu J, Wang SS, Yao YF, Yu B, Hu XP*
- 2001** Inadvertent globe penetration during retrobulbar anesthesia: A case report  
*Dai Y, Sun T, Gong JF*
- 2008** Systemic lupus erythematosus combined with primary hyperfibrinolysis and protein C and protein S deficiency: A case report  
*Liao YX, Guo YF, Wang YX, Liu AH, Zhang CL*
- 2015** Interstitial lung disease induced by the roots of *Achyranthes japonica* Nakai: Three case reports  
*Moon DS, Yoon SH, Lee SI, Park SG, Na YS*

**ABOUT COVER**

Gokul Sridharan, MD, PhD, Associate Professor, Oral Pathology and Microbiology, YMT Dental College and Hospital, Navi Mumbai, Mumbai 400018, Maharashtra, India. drgokuls@gmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for WJCC as 1.013; IF without journal self cites: 0.991; Ranking: 120 among 165 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2019 is 0.3 and Scopus CiteScore rank 2019: General Medicine is 394/529.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Jia-Hui Li; Production Department Director: Yu-Jie Ma; Editorial Office Director: Jin-Li Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

March 16, 2021

**COPYRIGHT**

© 2021 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## *Mycobacterium abscessus* infection after facial injection of argireline: A case report

Chao-Feng Chen, Jie Liu, Shuang-Shuang Wang, Yu-Fang Yao, Bo Yu, Xiao-Ping Hu

**ORCID number:** Chao-Feng Chen 0000-0001-6706-412X; Jie Liu 0000-0001-6420-5177; Shuang-Shuang Wang 0000-0001-8241-9048; Yu-Fang Yao 0000-0003-1114-3034; Bo Yu 0000-0002-5579-6806; Xiao-Ping Hu 0000-0002-5532-979X.

**Author contributions:** These authors contributed equally to this work and should be considered as joint first authors. Hu XP reviewed the data and consulted the authors; Chen CF and Liu J wrote the article and extracted the data; Wang SS, Yao YF, and Yu B revised and edited the article.

**Supported by** The Scientific Research Project of Peking University Shenzhen Hospital, No. JCYJ2018011 and the San-ming Project of Medicine in Shenzhen, No. SZSM201812059.

**Informed consent statement:** Informed written consent was obtained from the patient and her parents for the publication of this report and any accompanying images.

**Conflict-of-interest statement:** No conflict of interest is declared.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE

Chao-Feng Chen, Jie Liu, Shuang-Shuang Wang, Yu-Fang Yao, Bo Yu, Xiao-Ping Hu, Department of Dermatology & Venereology, Peking University Shenzhen Hospital, Shenzhen 518036, Guangdong Province, China

**Corresponding author:** Xiao-Ping Hu, PhD, Associate Professor, Department of Dermatology & Venereology, Peking University Shenzhen Hospital, No. 1120 Lianhua Road, Futian District, Shenzhen 518036, Guangdong Province, China. [xiaoping7752@sohu.com](mailto:xiaoping7752@sohu.com)

### Abstract

#### BACKGROUND

The incidence of infection with *Mycobacterium abscessus* (*M. abscessus*) has increased in recent years. This increase is partly associated with invasive cosmetic procedures.

#### CASE SUMMARY

The purpose of this case summary is to increase clinicians' awareness of *M. abscessus* infection and reduce mycobacterial infection caused by cosmetic procedures. We report the case of a 45-year-old woman who received acetyl hexapeptide-8 (argireline) injections in the forehead and temples, and erythema, nodules, and abscesses appeared at the injection sites after one week. The pus specimens were examined by microbiological culture and confirmed to be positive for *M. abscessus*. Clarithromycin 500 mg twice daily and moxifloxacin 400 mg once daily were administered for 5 mo and the lesions gradually subsided.

#### CONCLUSION

We report here for the first time a case of infection with *M. abscessus* after argireline injection. This condition is easily misdiagnosed as a common bacterial infection. Microbiological examinations are helpful for diagnosis and standardized cosmetic procedures can prevent infection with *M. abscessus*.

**Key Words:** Argireline; *Mycobacterium abscessus*; Nontuberculous *Mycobacterium*; Infection; Injection; Antibiotic therapy; Case report

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Unsolicited manuscript

**Specialty type:** Dermatology

**Country/Territory of origin:** China

**Peer-review report's scientific quality classification**

Grade A (Excellent): A

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

**Received:** November 23, 2020

**Peer-review started:** November 23, 2020

**First decision:** December 13, 2020

**Revised:** December 14, 2020

**Accepted:** January 15, 2021

**Article in press:** January 15, 2021

**Published online:** March 16, 2021

**P-Reviewer:** Fateh A, Munerato MC

**S-Editor:** Zhang H

**L-Editor:** Wang TQ

**P-Editor:** Li JH



**Core Tip:** *Mycobacterium abscessus* (*M. abscessus*) is a rapidly growing nontuberculous mycobacterium that can lead to infections of the lung, lymph node, skin, and soft tissue. The incidence of infection with *M. abscessus* has increased in recent years. This increase is partly associated with invasive cosmetic procedures. We report here for the first time a case of infection with *M. abscessus* after argireline injection. The purpose of this case summary is to increase clinicians' awareness of *M. abscessus* infection and reduce mycobacterial infection caused by cosmetic procedures.

**Citation:** Chen CF, Liu J, Wang SS, Yao YF, Yu B, Hu XP. *Mycobacterium abscessus* infection after facial injection of argireline: A case report. *World J Clin Cases* 2021; 9(8): 1996-2000

**URL:** <https://www.wjgnet.com/2307-8960/full/v9/i8/1996.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v9.i8.1996>

## INTRODUCTION

*Mycobacterium abscessus* (*M. abscessus*) is a rapidly growing nontuberculous mycobacterium (NTM) that can lead to infections of the lung, lymph node, skin, and soft tissue<sup>[1]</sup>. Compared with *Mycobacterium tuberculosis*, NTM has weaker virulence and pathogenicity. In humans, *M. abscessus* is usually an opportunistic pathogen, with poor pathogenicity, and it is more likely to infect patients with chronic diseases or immunocompromised patients. With the recent popularization of cosmetic procedures, the number of healthy patients infected with NTM is further increasing<sup>[2]</sup>. We report here for the first time a case of *M. abscessus* infection in a middle-aged woman 1 wk following argireline injection in the forehead and temples.

## CASE PRESENTATION

### Chief complaints

Erythema, nodules, and abscesses appeared at the face injection sites for 9 wk.

### History of present illness

Ten weeks prior to presentation, to improve the appearance of wrinkles in the forehead and temples, a 45-year-old female patient went to a private cosmetic clinic and received an injection of a bottle of argireline (concentration unknown). Injection sites should not come into contact with water for 2 d. Erythema, nodules, and abscesses began to appear at the injection sites after 1 wk, with mild pain but no fever. Six weeks prior to presentation, the patient went to a private clinic and received injections of cephalosporin antibiotics for 10 d. The skin lesions slightly improved but then relapsed after treatment was stopped. Four weeks prior to presentation, she went to the district hospital for an abscess incision and drainage. The pus specimens were cultured and found to be positive for mycobacteria. Three weeks prior to presentation, she came to the outpatient clinic of our hospital.

### History of past illness


The patient was healthy.

### Physical examination


Physical examination showed that there were eight red nodules and abscesses on the forehead and temples. The diameter was approximately 0.5-1.0 cm (Figure 1).

### Laboratory examinations

Biopsy was performed. Hematoxylin-eosin staining revealed a cyst in the dermis that was surrounded by granulomatous structures formed by histiocytes, scattered multinucleate giant cells, and lymphocyte, plasma cell, and neutrophil infiltration (Figure 2). Periodic acid-Schiff staining was negative. Pus was taken for mycobacterial culture. Colonies appeared after 4 d of mycobacterial culture. *M. abscessus* was identified with a Vitek MS MALDI-TOF mass spectrometry system (bioMérieux,



**Figure 1 Abscesses and nodules over the argireline injection sites.** A: An abscess with a diameter of approximately 1.0 cm, following an incision, can be seen on the left temple; B: Three red nodules and abscesses with a diameter of approximately 0.5 cm can be seen on the forehead; C: Four red nodules and abscesses with a diameter of approximately 0.5 cm can be seen on the right temple.



**Figure 2 Pathological biopsy of the lesion was performed.** A: A cystic cavity can be seen in the dermis, surrounded by granulomatous structures formed by histiocytes [Hematoxylin-eosin staining (HE)  $\times$  40]; B: Scattered around the cystic cavity are multinucleate giant cells, accompanied by lymphocyte, plasma cell, and neutrophil infiltration (HE  $\times$  200).

France).

---

## FINAL DIAGNOSIS

---

Facial *M. abscessus* infection.

---

## TREATMENT

---

The patient was given oral clarithromycin 500 mg twice daily combined with oral minocycline 100 mg twice daily. After a week of treatment, the patient developed significant gastrointestinal side effects. Then, we adjusted the oral clarithromycin treatment to 500 mg twice daily and moxifloxacin 400 mg once daily for 5 mo.

---

## OUTCOME AND FOLLOW-UP

---

The lesions resolved, with scars and pigmentation (Figure 3) still observed at follow-up.




Figure 3 There was clinical resolution after 5 mo of treatment, but scars and pigmentation remained.

## DISCUSSION

Most mycobacteria are found in the native environment, for example, in soil, water, or aerosols, and are usually not pathogenic<sup>[3]</sup>. NTM can cause infections in the lungs, lymph nodes, and skin and subcutaneous soft tissues. Being dependent on the growth rate, NTM can be divided into rapidly growing mycobacteria (RGM) or slowly growing mycobacteria (SGM)<sup>[4]</sup>. Common RGM include *M. abscessus*, *Mycobacterium fortuitum*, and *Mycobacterium chelonae*. Lesions can be characterized by erythema, nodules, abscesses, and mild pain<sup>[5]</sup>. Biopsy and microbiological examination should be carried out in patients in whom this disease is highly suspected. In this case, the microbiological examination provided evidence for the diagnosis. However, the low positive rate of bacterial culture is the main factor that makes the disease difficult to diagnose. In this patient, microbial culture of a pus specimen was performed, mycobacteria were observed 4 d later, and *M. abscessus* was identified with a Vitek MS MALDI-TOF mass spectrometry system (bioMérieux, France). The Vitek MS MALDI-TOF mass spectrometry system was approved by the United States Food and Drug Administration on July 31, 2017 for the identification of mycobacteria. *M. abscessus* is naturally resistant to first-line anti-tuberculosis drugs. The results of the *in vitro* susceptibility test for *M. abscessus* are inconsistent. The relatively consistent results show that *M. abscessus* is sensitive (greater than 90%) to clarithromycin, amikacin, and cefoxitin; moderately sensitive (40%-60%) to linezolid, imipenem, moxifloxacin, and tetracycline antibiotics; and poorly sensitive (approximately 10%) to ciprofloxacin and the sulfamethoxazole tobramycin<sup>[6]</sup>. There is no standardized treatment for cutaneous *M. abscessus* infection, and many NTM strains show drug resistance with limited drug selection. Clarithromycin is the cornerstone of all therapeutic drugs. The treatment effect of a combination of clarithromycin as the preferred drug with other antibacterial agents lasts for at least 4 mo; however, surgical treatment is sometimes required<sup>[7]</sup>. In this case, we chose clarithromycin combined with minocycline for oral treatment, but after 1 wk of taking the drug, the patient developed a significant gastrointestinal reaction, so we adjusted the treatment plan to clarithromycin combined with moxifloxacin. The lesions improved, although with hyperpigmentation and scarring.

There are two major factors contributing to cutaneous NTM infection. The first is trauma, surgery, or direct exposure to the environment, and the other is systemic infection with cutaneous or subcutaneous soft tissue complications<sup>[5]</sup>. In recent years, cases of *M. abscessus* infection have been reported, mainly related to mesodermal therapy or cosmetic fillings<sup>[7]</sup>. It has been declared in the literature that *M. abscessus* infection has occurred after tattooing, cosmetic injection, acupuncture, and even hair transplantation<sup>[8-11]</sup>. Infection with *M. abscessus* after facial injection of argireline has not previously been reported.

In this case, infection occurred at all injection sites, and we suspect that the drug, drug solvent, or syringe was contaminated with the bacteria. Erythema, mound pustules, nodules, or abscesses appear within 1 wk to several weeks at the injection site. We should suspect infection with nontuberculous mycobacteria or fungi. Cosmetic procedures have become popular worldwide. Many clinics perform tattooing and injection operations. The environment must be strictly disinfected when we are performing treatment, and the surgical instruments should be sterilized. In

addition, water and solvent may also cause infection with *M. abscessus* during surgery, and sterile injection water should be utilized to dissolve the drug. When choosing locations to undergo cosmetic procedures, cosmetic patients must not mistakenly believe in false advertisements and must choose a formal institution.

Argireline is a novel anti-ageing product that is mostly used in cosmetics and is formulated into ointments and creams to prevent skin wrinkles. The anti-ageing mechanism of argireline is similar to that of botulinum toxin, which inhibits the release of neurotransmitters at neuromuscular synapses. Argireline is rarely used for subcutaneous or intramuscular injection<sup>[12]</sup>. We report here for the first time a case of *M. abscessus* infection following facial injection with argireline. When undergoing cosmetic injection, attention should be paid to the side effects and risks of treatment. If there is a nodule or abscess at the treatment site, the possibility of NTM infection should be investigated.

## CONCLUSION

We report here for the first time a case of infection with *M. abscessus* after argireline injection. This condition is easily misdiagnosed as a common bacterial infection. Microbiological examinations are helpful for diagnosis and standardized cosmetic procedures can prevent infection with *M. abscessus*.

## REFERENCES

- 1 **Griffith DE**, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, Holland SM, Horsburgh R, Huitt G, Iademarco MF, Iseman M, Olivier K, Ruoss S, von Reyn CF, Wallace RJ Jr, Winthrop K; ATS Mycobacterial Diseases Subcommittee; American Thoracic Society; Infectious Disease Society of America. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med* 2007; **175**: 367-416 [PMID: 17277290 DOI: 10.1164/rccm.200604-571ST]
- 2 **Lee WJ**, Kang SM, Sung H, Won CH, Chang SE, Lee MW, Kim MN, Choi JH, Moon KC. Nontuberculous mycobacterial infections of the skin: a retrospective study of 29 cases. *J Dermatol* 2010; **37**: 965-972 [PMID: 21039785 DOI: 10.1111/j.1346-8138.2010.00960.x]
- 3 **Bottai D**, Stinear TP, Supply P, Brosch R. Mycobacterial Pathogenomics and Evolution. *Microbiol Spectr* 2014; **2**: MGM2-0025 [PMID: 26082120 DOI: 10.1128/microbiolspec.MGM2-0025-2013]
- 4 **Brown-Elliott BA**, Wallace RJ Jr. Clinical and taxonomic status of pathogenic nonpigmented or late-pigmenting rapidly growing mycobacteria. *Clin Microbiol Rev* 2002; **15**: 716-746 [PMID: 12364376 DOI: 10.1128/cmr.15.4.716-746.2002]
- 5 **Kothavade RJ**, Dhurat RS, Mishra SN, Kothavade UR. Clinical and laboratory aspects of the diagnosis and management of cutaneous and subcutaneous infections caused by rapidly growing mycobacteria. *Eur J Clin Microbiol Infect Dis* 2013; **32**: 161-188 [PMID: 23139042 DOI: 10.1007/s10096-012-1766-8]
- 6 **Huang YC**, Liu MF, Shen GH, Lin CF, Kao CC, Liu PY, Shi ZY. Clinical outcome of *Mycobacterium abscessus* infection and antimicrobial susceptibility testing. *J Microbiol Immunol Infect* 2010; **43**: 401-406 [PMID: 21075707 DOI: 10.1016/S1684-1182(10)60063-1]
- 7 **Lee MR**, Sheng WH, Hung CC, Yu CJ, Lee LN, Hsueh PR. *Mycobacterium abscessus* Complex Infections in Humans. *Emerg Infect Dis* 2015; **21**: 1638-1646 [PMID: 26295364 DOI: 10.3201/2109.141634]
- 8 **Bechara C**, Macheras E, Heym B, Pages A, Auffret N. *Mycobacterium abscessus* skin infection after tattooing: first case report and review of the literature. *Dermatology* 2010; **221**: 1-4 [PMID: 20558973 DOI: 10.1159/000313974]
- 9 **Fang RY**, Sun QN. *Mycobacterium abscessus* infections following injection of botulinum toxin. *J Cosmet Dermatol* 2020; **19**: 817-819 [PMID: 31402555 DOI: 10.1111/jocd.13094]
- 10 **Choi WS**, Kim MJ, Park DW, Son SW, Yoon YK, Song T, Bae SM, Sohn JW, Cheong HJ, Kim MJ. Clarithromycin and amikacin vs. clarithromycin and moxifloxacin for the treatment of post-acupuncture cutaneous infections due to *Mycobacterium abscessus*: a prospective observational study. *Clin Microbiol Infect* 2011; **17**: 1084-1090 [PMID: 20946409 DOI: 10.1111/j.1469-0691.2010.03395.x]
- 11 **Eustace K**, Jolliffe V, Sahota A, Gholam K. Cutaneous *Mycobacterium abscessus* infection following hair transplant. *Clin Exp Dermatol* 2016; **41**: 768-770 [PMID: 27663154 DOI: 10.1111/ced.12900]
- 12 **Grosicki M**, Latacz G, Szopa A, Cukier A, Kieć-Kononowicz K. The study of cellular cytotoxicity of argireline - an anti-ageing peptide. *Acta Biochim Pol* 2014; **61**: 29-32 [PMID: 24644551]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

