

Match Overview

1	Crossref 32 words Mohamed E. Ahmed, Vidhu B. Joshi, Mohamed Badawy, Lance C. Pagliaro et al. "Radium-223 in the third-line setti	1%
2	Internet 24 words crawled on 11-Jan-2020 link.springer.com	1%
3	Crossref 10 words Kosmas Daskalakis, Olov Norlén, Andreas Karakatsanis, Per Hellman, Rolf Larsson, Peter Nygren, Peter Stålberg.	<1%
4	Crossref 9 words "Abstracts", <i>Journal of Thoracic Oncology</i> , 2009	<1%
5	Crossref 9 words "EANM'15", <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2015	<1%
6	Crossref 9 words Bellmunt, J.. "Sunitinib in Advanced Renal Cell Carcinoma: Clinical Evidence", <i>European Urology Supplements</i> , 20	<1%
7	Internet 8 words crawled on 06-May-2016 jnm.snmjournals.org	<1%
8	Crossref 8 words Patard, J.J.. "Tyrosine Kinase Inhibitors in Clinical Practice: Patient Selection", <i>European Urology Supplements</i> , 20	<1%

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 61095

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Advanced small-bowel well-differentiated neuroendocrine tumours: An international survey of practice on 3rd line treatment

Lamarca A *et al.* Advanced SBNETs: Survey of practice

Angela Lamarca, Mauro Cives, Louis de Mestier, Joakim Crona, Francesca Spada, Kjell Öberg, Marianne Pavel, Teresa Alonso-Gordoa

Abstract

BACKGROUND

Somatostatin analogues are an established first line therapy for well differentiated small bowel neuroendocrine tumours (Wd-SBNETs), while and **peptide receptor radionuclide**

Advanced small-bowel well-differentiated neuroendocrine tumours:



Sign in



ALL

IMAGES

VIDEOS



Add the Give with Bing extension >

51,100 Results

Any time ▾

Management of Small Bowel Neuroendocrine Tumors | JCO ...

<https://ascopubs.org/doi/full/10.1200/JOP.18.00135>

Aug 10, 2018 · Small bowel **neuroendocrine** tumors (NETs) are increasing in incidence and are now the most common primary malignancies of the small intestine. Despite this increase, the vague presentation and slow growth of these tumors lead to long delays in ...

Cited by: 9

Author: Aaron T Scott, James R Howe

Publish Year: 2018

Neuroendocrine tumours: what gastroenterologists need to ...

<https://fg.bmj.com/content/early/2020/09/22/flgastro-2020-101431> ▾

Sep 22, 2020 · Gastroenterologists are intermittently involved in diagnosing and managing patients who have **neuroendocrine tumours** (NETs). However, few UK gastroenterologists have received extensive training about this topic. This article aims to provide a brief introduction to NETs; it is aimed at a general gastroenterologist audience. NETs present in diverse ways and many symptomatic patients ...

Author: Mohid S Khan, D Mark Pritchard

Publish Year: 2020

Management of Small Bowel Neuroendocrine Tumors

<https://www.mdpi.com/2072-6604/11/9/1305/html>

Search Tools

Turn on Hover Translation (开启取词)

激活 Windows

转到“设置”以激活 Windows

Advanced small-bowel well-differentiated neuroendocrine tumours:



Sign in



Add the Give with Bing extension >

ALL

IMAGES

VIDEOS

53,000 Results

Any time ▾

Management of Small Bowel Neuroendocrine Tumors | JCO ...

<https://ascopubs.org/doi/10.1200/JOP.18.00135>

Aug 10, 2018 · Small bowel **neuroendocrine** tumors (NETs) are increasing in incidence and are now the most common primary malignancies of the small intestine. Despite this increase, the vague presentation and slow growth of these tumors lead to long delays in ...

Cited by: 10

Author: Aaron T Scott, James R Howe

Publish Year: 2018

Progress in the Evaluation and Treatment of Small Bowel ...

<https://ascopubs.org/doi/full/10.1200/JOP.18.00440>

The incidence of well-differentiated **neuroendocrine** tumors (NETs) has been climbing since at least the 1970s. 1 This feature, coupled with a typically indolent disease course, has translated into a relatively high prevalence of NETs. 1 With a median survival of 8.5 years in the setting of metastatic disease for small bowel NETs, 1 a thoughtful approach to care is necessary, balancing the ...

Author: Claire K. Mulvey, Emily K. Bergsland

Publish Year: 2018

Management of Small Bowel Neuroendocrine Tumors

<https://www.mdpi.com/2072-6694/11/9/1395/htm> ▾

This is due to the fact that small bowel **neuroendocrine** tumors have a significant metastatic potential, even at a size less than 2 cm. Rorstad et al compiled data in a systematic review showing that small bowel **neuroendocrine** tumors (NETs) had nodal and distant metastases at diagnosis, respectively, in 12% and 5% for tumors less than 1 cm, 70% ...

Cited by: 4

Author: Vincent Larouche, Vincent Larouche, Amit ...

Publish Year: 2019

The Surgical Management of Small Bowel Neuroendocrine ...

<https://journals.lww.com/pancreasjournal/Fulltext/...>

Small bowel **neuroendocrine** tumors (SBNETs) have been increasing in frequency over the past decades, and are now the most common type of small bowel **tumor**. Consequently, general surgeons and surgical oncologists are seeing more patients with SBNETs in their practices than ever before.

Neuroendocrine tumours: what gastroenterologists need to ...

<https://fg.bmj.com/content/early/2020/09/22/flgastro-2020-101431> ▾

Sep 22, 2020 · Gastroenterologists are intermittently involved in diagnosing and managing patients who have **neuroendocrine tumours** (NETs). However, few UK gastroenterologists have received extensive training about this topic. This article aims to provide a brief introduction to NETs; it is aimed at a general gastroenterologist audience. NETs present in diverse ways and many symptomatic patients ...

Author: Mohid S Khan, D Mark Pritchard

Publish Year: 2020

Some results are removed in response to a notice of local law requirement. For more information, please see [here](#).

1

2

3

4

5

>

Search Tools

Turn off Hover Translation (关闭取词)

36,800 Results Any time ▾

Management of Small Bowel Neuroendocrine Tumors | JCO ...
<https://ascopubs.org/doi/full/10.1200/JOP.18.00135>
 Aug 10, 2018 · Small bowel **neuroendocrine** tumors (NETs) are increasing in incidence and are now the most common primary malignancies of the small intestine. Despite this increase, the vague presentation and slow growth of these tumors lead to long delays in diagnosis, and many patients present with metastases.
Cited by: 11 **Author:** Aaron T Scott, James R Howe
Publish Year: 2018

PEOPLE ALSO ASK

- What is a small bowel neuroendocrine tumor? ▾
- Is metastatic disease a functional disorder? ▾
- What are the most common neuroendocrine tumours? ▾
- What are non pancreatic neuroendocrine tumors? ▾

Feedback

Management of Small Bowel Neuroendocrine Tumors
<https://www.mdpi.com/2072-6694/11/9/1395/htm>
 This is due to the fact that small bowel **neuroendocrine** tumors have a significant metastatic potential, even at a size less than 2 cm. Rorstad et al compiled data in a systematic review showing that small bowel **neuroendocrine** tumors (NETs) had nodal and distant metastases at diagnosis, respectively, in 12% and 5% for tumors less than 1 cm, 70% ...
Cited by: 5 **Author:** Vincent Larouche, Vincent Larouche, A...
Publish Year: 2019

The Surgical Management of Small Bowel Neuroendocrine ...
<https://journals.lww.com/pancreasjournal/Fulltext/...>
 Small bowel **neuroendocrine** tumors (SBNETs) have been increasing in frequency over the past
 BPG胶片核对等8个会话