

Hepatocellular carcinoma with neuroendocrine ...

<https://www.sciencedirect.com/science/article/pii/S0014480017304446>

Oct 01, 2017 · Hepatocellular carcinoma with neuroendocrine differentiation, where tumor cells stain for both hepatocellular and neuroendocrine markers, is extremely rare. We report a case of a 65-year-old man who presented with a 14-cm rapidly growing mass in the right lobe of his liver with local extension into the gallbladder and portal vein.

Cited by: 5 Author: Jiajie G. Lu, M. Aabid Farukhi, Donna Ma...

Publish Year: 2017

(PDF) Synchronous hepatocellular carcinoma and gallbladder ...

https://www.researchgate.net/publication/346326459_Synchronous_hepatocellular...

Synchronous hepatocellular carcinoma and gallbladder adenocarcinoma with neuroendocrine differentiation: a case report and literature review October 2020 BMC Surgery 20(1)

Synchronous hepatocellular carcinoma and gallbladder ...

<https://bmcsurg.biomedcentral.com/articles/10.1186/s12893-020-00905-6> ▾

Oct 20, 2020 · Double primary cancers have a low incidence rate, and synchronous hepatocellular carcinoma and gallbladder adenocarcinoma are rarely reported. Here, we report such a case—the 12th case of synchronous double primary cancers featuring HCC and GC, but the first case of neuroendocrine differentiation in the gallbladder. A 77-year-old female was admitted to the ...

Author: Yan Xu, Quan-Ning Chen, Hui Wang, ... Publish Year: 2020

Collision tumor of hepatocellular carcinoma and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5107604>

Nov 07, 2016 · Core tip: Only 2 cases of collision tumor of hepatocellular carcinoma and primary hepatic neuroendocrine carcinoma involving the liver have been reported in the literature. This case shows different clinical characteristics from the previous cases. And we analyzed total 8 previous cases reported as concurrent occurrence of hepatocellular carcinoma and neuroendocrine carcinoma.

Cited by: 6 Author: Gyu Ho Choi, Sun Young Ann, Soon Il Le...

Publish Year: 2016

[PDF] CASE REPORT Open Access Synchronous hepatocellular ...

https://www.researchgate.net/journal/1471-2482_BMC...

Here, we report such a case the 12th case of synchronous double primary cancers featuring HCC and GC, but the first case of neuroendocrine differentiation in the gallbladder.

[PDF] Synchronous hepatocellular carcinoma and gallbladder ...

ALL IMAGES VIDEOS

79,100 Results Any time Open links in new tab

Hepatocellular carcinoma with neuroendocrine ...

https://www.sciencedirect.com/science/article/pii/S0014480017304446

Oct 01, 2017 · Hepatocellular carcinoma with neuroendocrine differentiation, where tumor cells stain for both hepatocellular and neuroendocrine markers, is extremely rare. We report a case of a 65-year-old man who presented with a 14-cm rapidly growing mass in the right lobe of his liver with local extension into the gallbladder and portal vein.

Cited by: 5 Author: Jiajie G. Lu, M. Aabid Farukhi, Donna M... Publish Year: 2017

(PDF) Synchronous hepatocellular carcinoma and gallbladder ...

https://www.researchgate.net/publication/346326459...

Synchronous hepatocellular carcinoma and gallbladder adenocarcinoma with neuroendocrine differentiation: a case report and literature review October 2020 BMC Surgery 20(1)

Intrahepatic bile duct mixed adenoneuroendocrine carcinoma ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6265897/30

Background. Malignancies affecting the extrahepatic bile duct are rare, accounting for around 0.1–0.2% of all cancer diagnoses []. Of these cholangiocarcinoma is the commonest cause, with adenocarcinoma making up around 80% of cases and other subtypes occurring much less

Hepatocellular Carcinoma

Medical Condition

The most common form of liver cancer often seen in people with chronic liver diseases like cirrhosis.

Rare (Fewer than 200,000 cases per year in US)

May be preventable by vaccine

Often requires lab test or imaging

Treatment from medical professional advised

Can last several months or years

Caused mainly due to the chronic infection of the liver or DNA mutation of the liver cells. Characterized by unexplained weight loss, upper abdominal pain and yellowing of the skin. Treatment options include hepatectomy, target drug delivery and the use of radiation to kill the cancerous cells.

Symptoms

The early stages of the disease do not present any symptoms but in the later stages the symptoms which are noted are as follows:

激活 Windows

Windows 激活工具

Name of Journal: World Journal of Clinical Oncology
 Manuscript Type: CASE REPORT

Hepatocellular carcinoma with Biliary and Neuroendocrine Differentiation: A Case Report and Review of the Literature

Dimopoulos YP, Winslow ER, He AR and Ozdemirli M

Abstract

BACKGROUND

Liver tumors with dual differentiations (combined hepatocellular carcinoma and

Match Overview

1 **Crossref** 17 words
 U. Syversen "Clinical significance of elevated serum chro...
 ogramin A levels", Scandinavian Journal of Gastroenterology. <1%

2 **Internet** 14 words
 crawled on 07-Sep-2019
[tesisena.com/publications.com](https://www.tesisena.com/publications.com) <1%

激活 Windows
转到设置以激活 Windows。

Hepatocellular carcinoma with biliary and neuroendocrine differenti



ALL IMAGES VIDEOS

145,000 Results Any time ▾

Hepatocellular carcinoma with neuroendocrine ...

<https://www.sciencedirect.com/science/article/pii/S0014480017304446>

Oct 01, 2017 - Hepatocellular carcinoma with neuroendocrine differentiation, where **tumor cells stain** for both hepatocellular and neuroendocrine markers, is extremely rare. We report a case of a 65-year-old man who presented with a 14-cm rapidly growing mass in the right lobe of his liver with local extension into the gallbladder and portal vein.

Cited by: 5 Author: Jiajie G. Lu, M. Aabid Farukhi, Donna Ma...
Publish Year: 2017

Synchronous hepatocellular carcinoma and gallbladder ...

<https://pubmed.ncbi.nlm.nih.gov/33081789>

Here, we report such a case- the 12th case of synchronous double primary cancers featuring HCC and GC, but the first case of neuroendocrine differentiation in the gallbladder. Case presentation: A 77-year-old female was admitted to the hospital complaining of weakness and inappetence for six months. Contrast-enhanced computed tomography (CT) of the abdomen indicated an 11 cm space-occupying ...

PEOPLE ALSO ASK

What is hepatocellular carcinoma? ▾

What is the prevalence of hepatic cancer? ▾

Is CD56 a neuroendocrine marker? ▾

Feedback

Mucin-producing hepatocellular carcinoma without ...

<https://www.ncbi.nlm.nih.gov/pubmed/30200114>

RATIONALE: Hepatocellular carcinoma has been believed not to produce mucin unless it has biliary differentiation. However, some cases of hepatocellular carcinoma with extracellular myxoid change have been reported recently, raising the possibility that, in rare cases, hepatocellular carcinoma cells ...

Cited by: 1 Author: Masayuki Ueno, Hiroyuki Takabatake, Ta...
Publish Year: 2018