

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 61652

Manuscript Type: CASE REPORT

"Antifreeze glycoprotein" bundles for an extremely preterm infant who underwent difficult removal of a peripherally inserted central catheter: A case report

Qiong Chen, Yan-Ling Hu, Shao-Yu Su, Xi Huang, Ying-Xin Li

Match Overview

1	Internet 116 words crawled on 16-Nov-2019 journals.sagepub.com	4%
2	Crossref 41 words Lawrence S. Miall. "Peripherally Inserted Central Cathete ... in Children With Cystic Fibrosis", <i>Journal of Infusion Nursin</i>	1%
3	Crossref 35 words Renfeng Li, Xia Cao, Tian Shi, Lei Xiong. "Application of pe ripherally inserted central catheters in critically ill newbor ...	1%
4	Internet 22 words crawled on 18-Jul-2020 www.readbag.com	1%
5	Internet 17 words crawled on 13-Feb-2014 www.health.qld.gov.au	1%
6	Crossref 13 words H Ekelund, U Hedner. "Prothrombin and vitamin K deficie ...	<1%

"AFGP" bundle for an extremely preterm infant who underwent level



ALL

IMAGES

VIDEOS

10,700 Results

Any time ▼

The development of a risk score for unplanned removal of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4547071>

to develop a risk score for unplanned **removal** of **peripherally inserted central catheter** in newborns. prospective cohort study conducted in a neonatal intensive care unit with newborn babies **who underwent** 524 **catheter** insertions. The clinical characteristics ...

Cited by: 2**Author:** Priscila Costa, Amélia Fumiko Kimura, Debr...**Publish Year:** 2015

Search Tools

[Turn off Hover Translation \(关闭取词\)](#)

Central line-associated bloodstream infections in the NICU ...

<https://www.sciencedirect.com/science/article/pii/S0146000517300198>

Apr 01, 2017 · A **bundle** is a small group of evidence-based interventions that, when implemented together, result in a better outcome than when executed individually. 22 **Central** line insertion and maintenance **bundle** components are often derived from recommendations by national organizations such as the Centers for Disease Control and Prevention (CDC) and the Healthcare Infection Control ...

Cited by: 18**Author:** Renée E. Mobley, Matthew J. Bizzarro**Publish Year:** 2017

2020 Southern Regional Meeting | Journal of Investigative ...

<https://jim.bmj.com/content/68/2/435.1> ▼

"Antifreeze glycoprotein" bundles for an extremely preterm infant w



ALL

IMAGES

VIDEOS

39,000 Results

Any time ▾

Open links in new tab



[Central line-associated bloodstream infections in the NICU ...](#)

<https://www.sciencedirect.com/science/article/pii/S0146000517300198>

Apr 01, 2017 · Introduction. **Preterm** infants in the neonatal intensive care unit (NICU) are highly susceptible to healthcare-acquired infections (HAI) given their immature immune systems and need for invasive procedures, support devices, and prolonged hospitalization. 1 **Preterm infants** with HAI are significantly more likely to die and, in survivors, to experience neurodevelopmental and growth ...

Cited by: 18

Author: Renée E. Mobley, Matthew J. Bizzarro

Publish Year: 2017

[\(PDF\) Variation in infection prevention practices for ...](#)

<https://www.researchgate.net/publication/328682356...>

Nov 01, 2018 · After implementation of **central-line bundles** the CLABSI incidence ranged from 0 to 19·5 per 1000 **catheter-days** (median 2·6, IQR 1·2-4·4) in all types of ICUs.

[Health Care–Associated Infections in the Neonatal ...](#)

<https://www.sciencedirect.com/science/article/pii/S0146000512000742>

Dec 01, 2012 · **Central** lines are defined as those whose tip terminates in one of the great vessel, such

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

16,300 Results

Any time ▾

The development of a risk score for unplanned removal of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4547071>

to develop a risk score for unplanned **removal** of **peripherally inserted central catheter** in newborns. prospective cohort study conducted in a neonatal intensive care unit with newborn babies **who underwent** 524 **catheter** insertions. The clinical characteristics ...

Cited by: 2**Author:** Priscila Costa, Amélia Fumiko Kimura, ...**Publish Year:** 2015

Antimicrobial-impregnated central venous catheters for ...

<https://www.researchgate.net/publication/345968053...>

Nov 17, 2020 · Rifampicin resistance in positive **peripherally inserted central venous catheter** tip cultures was higher in the antimicrobial-impregnated **peripherally inserted central venous catheter** group ...

Central line-associated bloodstream infections in the NICU ...

<https://www.sciencedirect.com/science/article/pii/S0146000517300198>

Apr 01, 2017 · Introduction. **Preterm** infants in the neonatal intensive care unit (NICU) are highly susceptible to healthcare-acquired infections (HAI) given their immature immune systems and need for invasive procedures, support devices, and prolonged hospitalization. 1 **Preterm** infants with HAI are significantly more likely to die and, in survivors, to experience neurodevelopmental ...

Cited by: 19**Author:** Renée E. Mobley, Matthew J. Bizzarro**Publish Year:** 2017

(PDF) Variation in infection prevention practices for ...

<https://www.researchgate.net/publication/328682356...>

Nov 01, 2018 · After implementation of **central-line bundles** the CLABSI incidence ranged from 0 to 19.5 per 1000 **catheter**-days (median 2.6, IQR 1.2-4.4) in all types of ICUs.

The development of a risk score for unplanned removal of ...

www.scielo.br/scielo.php?script=sci_arttext&pid=S0104-11692015000300475 ▾

METHOD: prospective cohort study conducted in a neonatal intensive care unit with newborn babies **who underwent** 524 **catheter** insertions. The clinical characteristics of the newborn