

4 Name of Journal: *World Journal of Gastrointestinal Surgery*

Manuscript NO: 61677

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Therapeutic effects of the TST36 stapler on rectocele combined with internal rectal prolapse

Meng J *et al.* Therapeutic effects of the TST36 stapler

Jin Meng, Zhi-Tao Yin, Ying-Yi Zhang, Yong Zhang, Xiu Zhao, Qing Zhai, De-Yu Chen, Wei-Gang Yu, Lei Wang, Zhi-Gang Wang

Match Overview

1	Internet 43 words crawled on 01-Dec-2020 www.spandidos-publications.com	1%
2	Internet 28 words crawled on 11-Aug-2020 pesquisa.bvsalud.org	1%
3	Internet 15 words crawled on 23-Apr-2015 surgwiki.com	<1%
4	Internet 13 words crawled on 05-Dec-2019 f6publishing.blob.core.windows.net	<1%
5	Internet 12 words crawled on 15-Jul-2020 opendentistryjournal.com	<1%
6	Internet 12 words crawled on 01-Feb-2019	<1%

ALL

IMAGES

VIDEOS

9,660 Results

Any time ▾

TST36 stapling for rectocele and hemorrhoidal prolapse ...

<https://www.researchgate.net/publication/312330521...>

Altomare ODS score and CCIS improved significantly after surgery. Conclusions: Despite a notable complication rate during surgery and the postoperative period, the TST36 can be considered as an...

A transanal procedure using TST STARR Plus for the ...

<https://www.sciencedirect.com/science/article/pii/S1743919116301996>

Aug 01, 2016 - The optimal technique for ODS caused by rectocele and internal rectal prolapse is still controversial. Nevertheless, after 18 month mid-term follow up, our study suggests that the TST STARR Plus procedure is a simple and safe method which effectively improves defecation in carefully selected patients with rectocele and/or internal rectal prolapse.

Cited by: 7

Author: X.H. Ren, S.M. Yaseen, Y.L. Cao, W.C. L...

Publish Year: 2016

PEOPLE ALSO ASK

Is perineal stapled prolapse resection effective? ▾

What is the procedure for rectal prolapse? ▾

What is the treatment for internal prolapse? ▾

What kind of Doctor treats rectal prolapse? ▾

Feedback

Khubchandani's procedure combined with stapled posterior ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6429338>

Mar 21, 2019 - Our department developed a new procedure that combined Khubchandani's procedure with stapled posterior rectal wall resection (KSPRWR) in 42 female patients with ODS caused by rectocele combined with or without rectal mucosal prolapse to evaluate and compare its safety and effectiveness with the STARR procedure.

Cited by: 1

Author: Yi Shao, Yong-Xing Fu, Qing-Fa Wang, Z...

Publish Year: 2019



ALL

IMAGES

VIDEOS

9,930 Results

Any time ▾

Open links in new tab



[A transanal procedure using TST STARR Plus for the ...](#)

<https://www.sciencedirect.com/science/article/pii/S1743919116301996>

Aug 01, 2016 · The optimal technique for ODS caused by **rectocele** and **internal rectal prolapse** is still controversial. Nevertheless, after 18 month mid-term follow up, our study suggests that the TST STARR Plus procedure is a simple and safe method which effectively improves defecation in carefully selected patients with **rectocele** and/or **internal rectal prolapse**.

Cited by: 7

Author: X.H. Ren, S.M. Yaseen, Y.L. Cao, W.C....

Publish Year: 2016

[Khubchandani's procedure combined with stapled posterior ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6429338>

Mar 21, 2019 · Our department developed a new procedure that **combined** Khubchandani's procedure with stapled posterior **rectal** wall resection (KSPRWR) in 42 female patients with ODS caused by **rectocele** **combined** with or without **rectal** mucosal **prolapse** to evaluate and compare its safety and effectiveness with the STARR procedure.

Cited by: 1

Author: Yi Shao, Yong-Xing Fu, Qing-Fa Wang,...

Publish Year: 2019

PEOPLE ALSO ASK

What is solitary rectal prolapse?



Can a prolapse cause a rectal ulcer?



What is the treatment for rectal prolapse?



Can chronic straining cause rectal ulcers?



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

TOOLS ▾

[Combined Perineal and Endorectal Repair of Rectocele by ...](https://www.researchgate.net/publication/11035671...)

<https://www.researchgate.net/publication/11035671...>

Combined perineal and endorectal stapler repair of rectocele may be a useful new surgical tool for **correcting distention rectocele** associated with mucosal prolapse or hemorrhoids and perineal...

[Khubchandani's procedure combined with stapled posterior ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6429338)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6429338>

Mar 21, 2019 · Our department developed a new procedure that **combined** Khubchandani's procedure with stapled posterior **rectal** wall resection (KSPRWR) in 42 female patients with ODS caused by **rectocele combined** with or without **rectal** mucosal **prolapse** to evaluate and compare its safety and effectiveness with the STARR procedure.

Cited by: 2

Author: Yi Shao, Yong-Xing Fu, Qing-Fa Wang, Z...

Publish Year: 2019

[GMS | GMS German Medical Science - GMS | Homepage](http://www.egms.de/static/de/journals/gms/2016-14/000241.shtml)

www.egms.de/static/de/journals/gms/2016-14/000241.shtml ▾

Conclusions: Despite a notable complication rate during surgery and the postoperative period, the **TST36** can be considered as an effective tool for low **rectal** stapling for anorectal **prolapse** causing hemorrhoids or obstructed defecation.

[A transanal procedure using TST STARR Plus for the ...](https://www.sciencedirect.com/science/article/pii/S1743919116301996)

<https://www.sciencedirect.com/science/article/pii/S1743919116301996>

Aug 01, 2016 · **Rectocele** and **internal rectal prolapse** are the most common causes of ODS. Based on the stapled haemorrhoidopexy procedure, an alternative minimally invasive transanal operation technique called STARR was developed by Longo to treat ODS patients with **rectocele** and **internal rectal prolapse**. The main purpose of this procedure was to eliminate **rectocele** and **internal prolapse** by strengthening the anterior **rectal** ...

Cited by: 7

Author: X.H. Ren, S.M. Yaseen, Y.L. Cao, W.C. Li...

Publish Year: 2016

[\[Transtar™ operation for rectocele and obstructed ...](https://pubmed.ncbi.nlm.nih.gov/27742925)

<https://pubmed.ncbi.nlm.nih.gov/27742925>

The Contour® Transtar™ operation represents a further methodological development of conventional transanal stapled **rectal** resection (STARR) for the treatment of obstructed defecation syndrome (ODS) and/or full thickness **rectal prolapse**. In contrast to the conventional STARR technique a specially des ...