



Supplementary Figure 1 Distribution of patients in the pre-transplant period, immediate postoperative period and over 10 years follow-up of the orthotopic liver transplantation.

Supplementary Table 1 METAVIR histology scores

Histological activity	Inflammatory infiltrate	Fibrosis	Fibrosis
A0	No activity	F0	Without fibrosis
A1	Minimum activity: ¹ Fragmented necrosis-grade: 1; ² Lobular necrosis-grade: 0 a 1	F1	Portal fibrosis without septa
A2	Moderate activity: ¹ Fragmented necrosis-grade: 2; ² Lobular necrosis-grade: 0 a 1	F2	Portal fibrosis with few septa
A3	Severe activity: ¹ Fragmented necrosis-grade: 3; ² Lobular necrosis-grade: 0 a 2	F3	Numerous septa without cirrhosis
		F4	Cirrhosis

¹Fragmented necrosis (0: Absent; 1: Focal change in the periportal plate in some portal tracts; 2: Diffuse change in the periportal plate in some portal tracts or focal lesions in all portal tracts; 3: Diffuse change in the periportal plate in all portals); ²Lobular necrosis (0: Less than one necroinflammatory focus per lobule; 1: At least one necroinflammatory focus per lobule; 2: Several necroinflammatory focus per lobule or confluent or bridging necrosis).