

国内版

国际版

Should we use full analgesic dose of opioids for organ procurement



ALL

IMAGES

VIDEOS

50,000 Results

Any time ▾

[PDF] Opioid Analgesics - Stanford Medicine

med.stanford.edu/.../documents/opioid-analgesics.pdf

Opioid Analgesics Ardin S. Berger, D.O. Department of Anesthesiology & Pain Management Ver revised 10/12/19 ... respiratory drive center brainstem (6-18 hours after administration) ... –Doses analgesic dose –Coadministration with mu agonists may result in partial reversal

Pharmacists' guide to the management of organ donors ...

<https://academic.oup.com/ajhp/article-abstract/73/22/1829/5102132> ▾

Nov 15, 2016 · Solid organ transplantation remains a vital area of healthcare, with over 24,000 transplants performed in 2014 alone. 1 As of August 2016, approximately 120,000 patients were in of a lifesaving organ. Every 10 minutes, 1 person is added to the list of those needing a transplant. 21 people die each day waiting for a transplant. 1 The number of patients on this list continues to

Cited by: 4

Author: Catherine Korte, Jennifer L. Garber, Jillian...

Publish Year: 2016

Intraoperative management of brain-dead organ donors by ...

<https://bmcanaesthesiol.biomedcentral.com/articles/10.1186/s12871-019-0766-y>

Jun 15, 2019 · Brain-dead donors (BDD) currently remain the primary source of grafts for solid organ transplantation across the world [1, 2]. In this context, appropriate management of organ donors from diagnosis of brain death to the end of the organ procurement (OP) procedure is of paramount importance to optimize the function of potential grafts.

Cited by: 1

Author: Benoit Champigneulle, Arthur Neuschwan...

Publish Year: 2019

Opioids and alpha-2-agonists for analgesia and sedation in ...

<https://systematicreviewsjournal.biomedcentral.com/...> ▾

Aug 20, 2020 · The analgesic and sedative properties of the alpha-2-agonists without respiratory or gastrointestinal side effects, along with the potential beneficial effects to the immature brain, have promoted their use to enable reduced opioid dosage or even to replace opioid ...

Author: Mari Kinoshita, Mari Kinoshita, Katarz... Publish Year: 2020

Anaesthesia for organ donation in the brainstem dead - Why ...

<https://www.researchgate.net/publication/12662430...>

[\[PDF\] Opioid Analgesics - Stanford Medicine](#)med.stanford.edu/.../documents/opioid-analgesics.pdf

Opioid Analgesics Ardin S. Berger, D.O. Department of Anesthesiology & Pain Management Version 1, revised 10/12/19 ... respiratory drive center brainstem (6-18 hours after administration) ... -Doses ≤ analgesic dose -Coadministration with mu agonists may result in partial reversal

[Pharmacists' guide to the management of organ donors ...](#)<https://academic.oup.com/ajhp/article-abstract/73/22/1829/5102132> ▾

Nov 15, 2016 · Solid organ transplantation remains a vital area of healthcare, with over 24,000 transplants performed in 2014 alone. 1 As of August 2016, approximately 120,000 patients were in need of a lifesaving organ. Every 10 minutes, 1 person is added to the list of those needing a transplant, while 21 people die each day waiting for a transplant. 1 The number of patients on this list continues to grow ...

Cited by: 4

Author: Catherine Korte, Jennifer L. Garber, Jillian...

Publish Year: 2016

PEOPLE ALSO ASK

How long does it take to release opioids? ▾

What is the breakthrough dose of morphine? ▾

What is immediate release morphine? ▾

Feedback

[Opioids and alpha-2-agonists for analgesia and sedation in ...](#)<https://systematicreviewsjournal.biomedcentral.com/...> ▾

Aug 20, 2020 · The analgesic and sedative properties of the alpha-2-agonists without respiratory and gastrointestinal side effects, along with the potential beneficial effects to the immature brain, have promoted their use to enable reduced opioid dosage or even to replace opioid ...

Author: Mari Kinoshita, Mari Kinoshita, Katarz... Publish Year: 2020

[Intraoperative management of brain-dead organ donors by ...](#)<https://bmcanesthesiol.biomedcentral.com/articles/10.1186/s12871-019-0766-y> ▾

14 Feb 2021 04:32 PM 62050 words + 0 matches + 0 sources 62050_Auto_Edited.docx Quotes Included Bibliography Included 0%

iThenticate

Name of Journal: *World Journal of Meta-Analysis*
Manuscript NO: 62050
Manuscript Type: MINIREVIEWS

Should we use full analgesic dose of opioids for organ procurement in brainstem dead?

Charlier P *et al.* Dose of opioids for organ procurement in brainstem dead

Philippe Charlier, John Rebibo, Nadia Benmoussa

Abstract
Families facing the growing demand for organ removal from their loved ones are

Match Overview

There are no matching sources for this paper.

PAGE 1 OF 3 Test Only Report

国内版 国际版

Should we use full analgesic dose of opioids for organ procurement



ALL IMAGES VIDEOS

49,600 Results Any time ▾

[PDF] Opioid Analgesics - Stanford Medicine

med.stanford.edu/.../documents/opioid-analgesics.pdf

Opioid Analgesics Ardin S. Berger, D.O. Department of Anesthesiology & Pain Management Version 1, revised 10/12/19 ... respiratory drive center brainstem (6-18 hours after administration) ... -Doses ≤ analgesic dose -Coadministration with mu agonists may result in partial reversal

Pharmacists' guide to the management of organ donors ...

<https://academic.oup.com/ajhp/article-abstract/73/22/1829/5102132> ▾

Nov 15, 2016 · Solid organ transplantation remains a vital area of healthcare, with over 24,000 transplants performed in 2014 alone. 1 As of August 2016, approximately 120,000 patients were in need of a lifesaving organ. Every 10 minutes, 1 person is added to the list of those needing a transplant, while 21 people die each day waiting for a transplant. 1 The number of patients on this list continues to grow ...

Cited by: 4 Author: Catherine Korte, Jennifer L. Garber, Jillian...

Publish Year: 2016

PEOPLE ALSO ASK

How long does it take to release opioids? ▾

How are intrathecal opioids determined? ▾

What is the fate of opioids after injection into CSF? ▾

How does fentanyl work in the spinal cord? ▾

Feedback

Intraoperative management of brain-dead organ donors by ...

<https://bmcanesthesiol.biomedcentral.com/articles/10.1186/s12871-019-0766-y> ▾

Jun 15, 2019 · Brain-dead donors (BDD) currently remain the primary source of grafts for solid organ transplantation across the world [1, 2]. In this context, appropriate management of organ donors from the diagnosis of brain death to the end of the organ procurement (OP) procedure is of paramount importance to optimize the function of potential grafts.

Cited by: 1 Author: Benoit Champigneulle, Arthur Neuschwan...

Publish Year: 2019