

32,200 Results Any time

### Hepatic Abscess: A Practical Approach

<https://www.researchgate.net/publication/328859366...>

We reported a rare case of abscess of the caudate lobe of the liver in a 60-year old man. We first tried computed tomography (CT) guided percutaneous drainage of the abscess but failed to ...

Estimated Reading Time: 10 mins

### Bronchobiliary Fistula after Radiofrequency Thermal ...

[https://www.researchgate.net/publication/7976570\\_Bronchobiliary\\_Fistula\\_after...](https://www.researchgate.net/publication/7976570_Bronchobiliary_Fistula_after...)

Bronchobiliary fistula is a rare disorder usually caused by liver abscess [10 ... and caudate lobe in one. The tumor size was 2.0 to 6.4 cm (mean, 3.9 cm). ... which might be secondary to a ...

### Inflammatory Pseudotumor: The Great Mimicker | Request PDF

<https://www.researchgate.net/publication/221854640...>

Hepatic inflammatory pseudotumor secondary to a foreign body at right lobe of liver. ... the left liver



[Alveolar Echinococcosis: Spectrum of Findings at Cross...](https://pubs.rsna.org/doi/full/10.1148/rg.327125708)  
<https://pubs.rsna.org/doi/full/10.1148/rg.327125708>

[Introduction](#) [Epidemiologic and Pathophys...](#) [Clinical Features](#) [Diagnosis](#)

**Alveolar echinococcosis** is caused by the metacystode of the fox tapeworm *Echinococcus multilocularis*. In humans, metacystode cells of *E. multilocularis* proliferate in the liver, inducing slowly progressive, life-threatening tumortike growths (1-4). The parasitic lesions may subsequently disseminate to neighboring organs and tissues or metastasize to distant locations. The prognosis is generally poor, and liver transplantation may be required in patients with inoperable lesions, chronic liver failure, or bot...

See more on [pubs.rsna.org](https://pubs.rsna.org)

Cited by: 122 Author: Mecit Kantarci, Ummugusum Bayraktuta...  
Publish Year: 2012

[Successful diagnosis and treatment of ingested wooden ...](#)

<https://journals.lww.com/md-journal/Fulltext/2018/...>

Nov 22, 2017 - Foreign body ingestion is a common phenomenon and foreign bodies are mostly excreted in stool. Once sharp bodies are ingested without being realized, perforation of intestine is...

[Laparoscopic management of enterohepatic migrated fish ...](https://europepmc.org/article/MED/30882633)

<https://europepmc.org/article/MED/30882633>

Mar 01, 2019 - When there is a strong suspicion of bowel perforation by a foreign body or if a foreign body was detected by imaging with liver impairment, surgery is the mainstay of diagnosis and...

Search Tools

Turn off Hover Translation (关闭翻译)

**Name of Journal:** *World Journal of Clinical Cases*

**Manuscript NO:** 62062

**Manuscript Type:** CASE REPORT

**Hepatic abscess caused by esophageal foreign body misdiagnosed as cystadenocarcinoma by magnetic resonance imaging: a Case report**

Pan W *et al.* Hepatic abscess misdiagnosed as cystadenocarcinoma

Wei Pan, Li-Jing Lin, Ze-Wu Meng, Xin-Ran Cai, Yan-Ling Chen

Match Overview

Match #	Source	Words	Similarity
1	Internet crawled on 08-May-2021 <a href="http://www.eajnet.com">www.eajnet.com</a>	100	4%
2	Internet crawled on 03-Nov-2020 <a href="http://jmedicalcasereports.biomedcentral.com">jmedicalcasereports.biomedcentral.com</a>	17	1%
3	Crossref Wensu Yu, Hailin Yu, Jun Ling, Jiang Du, Zhijiang Yin, Chengen Li, Min Zhou. "Hepatic Abscess Secondary to Stomach Perforation".	14	1%
4	Crossref "Abstracts from the 2016 Society of General Internal Medicine Annual Meeting", Journal of General Internal Medicine, 2016	12	1%

国内版 国际版

Hepatic abscess caused by esophageal foreign body misdiagnosed



ALL IMAGES VIDEOS

73,700 Results Any time

### Alveolar Echinococcosis: Spectrum of Findings at Cross ...

<https://pubs.rsna.org/doi/full/10.1148/rg.327125708>

< **Introduction** Epidemiologic and Pathophy... Clinical Features Dia >

**Alveolar echinococcosis** is caused by the metacestode of the fox tapeworm *Echinococcus multilocularis*. In humans, metacestode cells of *E. multilocularis* proliferate in the liver, inducing slowly progressive, life-threatening tumorlike growths (1–4). The parasitic lesions may subsequently disseminate to neighboring organs and tissues or metastasize to distant locations. The prognosis is generally poor, and liver transplantation may be required in patients with inoperable lesions, chronic liver failure, or bot...

See more on [pubs.rsna.org](https://pubs.rsna.org)

Cited by: 122 Author: Mecit Kantarci, Ummugulsum Bayraktuta...

Publish Year: 2012

### Dysphagia Revisited: Common and Unusual Causes | ...

<https://pubs.rsna.org/doi/full/10.1148/rg.351130150>

Jan 15, 2015 · Introduction. Dysphagia can be caused by functional or structural abnormalities of the oral cavity, pharynx, esophagus, and/or gastric cardia. It is important to distinguish between oropharyngeal and substernal esophageal dysphagia, since potential causes, radiologic evaluation, and treatment may differ (). Dysphagia is defined by medical dictionaries as difficulty in swallowing.

Cited by: 22 Author: Laura R Carucci, Mary Ann Turner

Publish Year: 2015

#### PEOPLE ALSO ASK

- Is it possible to misdiagnose foreign body ingestion? ▾
- What kind of infection can cause hepatic abscess in liver? ▾
- When to use CT for swallowed foreign mass? ▾
- How are fluoroscopic studies used to evaluate the esophagus? ▾