

**Name of Journal:** *World Journal of Clinical Cases*

**Manuscript NO:** 62081

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Study*

**Effects on newborns of applying bupivacaine combined with different doses of fentanyl for cesarean section**

Yun Wang, Wen-Xun Liu, Xiao-Hong Zhou, Min Yang, Xin Liu, Yuan Zhang, Ke-Rong Hai, Qing-Shan Ye

**Abstract**

BACKGROUND

### Match Overview

1	<b>Internet</b> 15 words crawled on 07-Jun-2020 <a href="http://www.dovepress.com">www.dovepress.com</a>	1%
2	<b>Internet</b> 13 words crawled on 31-Oct-2020 <a href="http://www.researchsquare.com">www.researchsquare.com</a>	1%
3	<b>Internet</b> 11 words crawled on 03-Dec-2020 <a href="http://www.spandidos-publications.com">www.spandidos-publications.com</a>	1%
4	<b>Internet</b> 10 words crawled on 31-Oct-2020 <a href="http://bmcnephrol.biomedcentral.com">bmcnephrol.biomedcentral.com</a>	1%



ALL

IMAGES

VIDEOS

Add the Give with Bing extension &gt;

22,400 Results

Any time ▾

## A Randomised Controlled Trial to Evaluate the ...

<https://www.ncbi.nlm.nih.gov/pubmed/25504002>

PURPOSE: The purpose of this study was to evaluate comparatively, in women undergoing caesarean section under spinal anesthesia, the effectiveness of hyperbaric **bupivacaine combined** with 3 different adjuvants (**fentanyl**, clonidine, and dexmedtomidine) on quality of ...

Cited by: 21

Author: Z. Li, M. Tian, C.-Y. Zhang, A.-Z. Li, A.-J. ...

Publish Year: 2014



See more

## Comparison of Maternal and Neonatal Effects of Combined ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4894102>

The reason of finding higher ephedrine doses was that the intrathecal local anaesthetic dose, 10 mg hyperbaric **bupivacaine** and 20 µg **fentanyl**, the determined ED50 and ED95 dose for caesarean section, was higher than the dose used in the study of Coppejans and colleagues . Thus, the epidural catheter was not used in any of the patients for ...

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

22,500 Results

Any time ▾

### [A Randomised Controlled Trial to Evaluate the ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/25504002>

PURPOSE: The purpose of this study was to evaluate comparatively, in women undergoing caesarean **section** under spinal anesthesia, the effectiveness of hyperbaric **bupivacaine combined** with 3 **different** adjuvants (**fentanyl**, clonidine, and dexmedetomidine) on quality of ...

Cited by: 21

Author: Z. Li, M. Tian, C.-Y. Zhang, A.-Z. Li, A.-J. Hua...

Publish Year: 2014

### [Comparison of Maternal and Neonatal Effects of Combined ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4894102>

The reason of finding higher ephedrine **doses** was that the intrathecal local anaesthetic **dose**, 10 mg hyperbaric **bupivacaine** and 20 µg **fentanyl**, the determined ED50 and ED95 **dose** for caesarean **section**, was higher than the **dose** used in the study of Coppejans and colleagues . Thus, the epidural catheter was not used in any of the patients for ...

Cited by: 1

Author: Ece Dumanlar Tan, Berrin Gunaydin

Publish Year: 2014

#### PEOPLE ALSO ASK

How much bupivacaine for breastfeeding? ▾

What is the dosing algorithm for bupivacaine in spinal anaesthesia? ▾

How much Bupivacaine is injected into the subarachnoid space? ▾

What is the difference between fentanyl and epidural? ▾

Feedback

### [A double-blind randomized control trial to compare the ...](#)

<https://www.joacp.org/article.asp?issn=0970-9185;...>

Furthermore, Goel et al. found that the patients receiving 7.5 µg of **fentanyl in combination with low-dose bupivacaine** had a significantly higher number of failed blocks (almost 27%) than those receiving 10 or 12.5 µg of **fentanyl**. Therefore, most of the authors prefer to **use doses** of intrathecal **fentanyl** ...