

国内版国际版

Microsoft Bing

Stevens Johnson Syndrome And Concurrent Hand Foot Syndrome I

Sign in

ALLIMAGESVIDEOS

93,900 ResultsAny time

Stevens-Johnson Syndrome after Treatment with ...

<https://www.researchgate.net/publication/5689709...>

Request PDF | On Apr 1, 2008, M.A.N. Sendur and others published Stevens-Johnson Syndrome after Treatment with Capecitabine | Find, read and cite all the research you need on ResearchGate

Capecitabine (Professional Patient Advice) - Drugs.com

<https://www.drugs.com/ppa/capecitabine.html>

Jun 20, 2020 - Persistent hand-and-foot syndrome (grade 2 and higher) could eventually lead to fingerprint loss. If grade 2 or 3 hand-and-foot syndrome occurs, interrupt administration of capecitabine until the event resolves or decreases in intensity to grade 1. Following grade 3 hand-and-foot syndrome, decrease subsequent doses of therapy.

PEOPLE ALSO ASK

Search Tools

Turn off Hover Translation (关闭取词)

ALL IMAGES VIDEOS TOOLS ▾

Stevens-Johnson Syndrome after Treatment with Capecitabine ...<https://www.researchgate.net/publication/5689709...>

The most common dose-limiting adverse effects associated with capecitabine monotherapy are hyperbilirubinemia, diarrhea, and hand-foot syndrome. Myelosuppression, fatigue ...

Stevens-Johnson syndrome and toxic epidermal necrolysis ...<https://www.uptodate.com/contents/stevens-johnson...>

Mar 12, 2019 - A Case Report-Stevens-Johnson Syndrome as an Adverse Effect of Capecitabine. J Gastrointest Cancer 2018; 49:349. Doesch J, Debus D, Meyer C, et al. Afatinib-associated Stevens-Johnson syndrome in an EGFR-mutated lung cancer patient.


PEOPLE ALSO ASK

What is Stevens Johnson syndrome?	▾
Is there a Phase 2 trial for capecitabine?	▾
When to use capecitabine instead of 5 - FU / LV?	▾
Can you take phenytoin and capecitabine at the same time?	▾

Feedback

[Yeloda \(capecitabine\) dose, indications, adverse effects](#)**Stevens-Johnson Syndrome**

Medical Condition



A rare and serious disorder that affects skin, mucous membranes, genitals and eyes.

📊 Very rare (Fewer than 20,000 cases per year in US)

👤 Often requires lab test or imaging

🩺 Treatment from medical professional advised

国内版 国际版

Stevens Johnson syndrome and concurrent hand foot syndrome dui



ALL IMAGES VIDEOS

45,100 Results Any time ▾

Stevens-Johnson Syndrome after Treatment with ...

<https://www.researchgate.net/publication/5689709...>

The most common dose-limiting adverse effects associated with capecitabine monotherapy are hyperbilirubinemia, diarrhea, and hand-foot syndrome. Myelosuppression, fatigue and ...

Stevens-Johnson syndrome and toxic epidermal necrolysis ...

<https://www.uptodate.com/contents/stevens-johnson...>

Mar 12, 2019 · A Case Report-Stevens-Johnson Syndrome as an Adverse Effect of Capecitabine. J Gastrointest Cancer 2018; 49:349. Doesch J, Debus D, Meyer C, et al. Afatinib-associated Stevens-Johnson syndrome in an EGFR-mutated lung cancer patient.

PEOPLE ALSO ASK

What is Stevens Johnson syndrome? ▾

When to use capecitabine in patients with renal failure? ▾

Is there a Phase 2 trial for capecitabine? ▾

Does capecitabine cause coagulopathy? ▾

Feedback

Capecitabine Side Effects: Common, Severe, Long Term ...

<https://www.drugs.com/sfx/capecitabine-side-effects.html> ▾

in Summary

For The Consumer

Further Information

Commonly reported side effects of capecitabine include: anemia, diarrhea, hyperbilirubinemia, increased serum alkaline phosphatase, increased serum transaminases, nausea, neutropenia, palmar-plantar erythrodysesthesia, paresthesia, stomatitis, thrombocytopenia, and vomiting. Other side effects include: erythema of skin. See below for a comprehensive list of adverse effects.

[See more on drugs.com](#)