

**3**  
**Name of Journal:** *World Journal of Gastrointestinal Endoscopy*  
**Manuscript NO:** 62202  
**Manuscript Type:** ORIGINAL ARTICLE

### *Observational Study*

**Comparison of endoscopic gastritis based on Kyoto classification between diffuse and intestinal gastric cancer**

Toyoshima O *et al.* Endoscopic gastritis for diffuse and intestinal gastric cancer

Osamu Toyoshima, Toshihiro Nishizawa, Shuntaro Yoshida, Tomonori Aoki, Fumiko Nagura, Kosuke Sakitani, Yosuke Tsuji, Hayato Nakagawa, Hidekazu Suzuki, Kazuhiko Koike

## Match Overview

1	<b>Internet</b> 44 words crawled on 22-Sep-2020 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	1%
2	<b>Crossref</b> 28 words Satoki Shichijo, Yoshihiro Hirata, Ryota Niikura, Yoku Haya kawa, Atsuo Yamada, Kazuhiko Koike. "Association betw ...	1%
3	<b>Internet</b> 15 words crawled on 19-Feb-2020 <a href="http://f6publishing.blob.core.windows.net">f6publishing.blob.core.windows.net</a>	<1%
4	<b>Internet</b> 14 words <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	<1%
5	<b>Internet</b> 14 words crawled on 17-Feb-2021 <a href="http://www.spandidos-publications.com">www.spandidos-publications.com</a>	<1%
6	<b>Internet</b> 13 words crawled on 22-Jul-2020 <a href="http://bmjopen.bmj.com">bmjopen.bmj.com</a>	<1%

Comparison of endoscopic gastritis based on Kyoto classification be



Sign in

ALL

IMAGES

VIDEOS

56,500 Results

Any time ▼

### [Endoscopy-based Kyoto classification score of gastritis ...](#)

<https://pubmed.ncbi.nlm.nih.gov/32982115>

Background: **Endoscopy-based Kyoto classification** for **gastritis** and pathological topographic distribution of neutrophil infiltration are correlated with **gastric cancer** risk. Aim: To investigate t...

**Author:** Osamu Toyoshima, Toshihiro Nishi... **Publish Year:** 2020

See more

### [Efficacy of the Kyoto Classification of Gastritis in ...](#)

<https://pubmed.ncbi.nlm.nih.gov/28321054>

**Kyoto gastritis** scores of atrophy and intestinal metaplasia in the H. pylori-positive **cancer** group were significantly higher than in subjects with **gastritis** alone (all  $p < 0.001$ ). No significant...

**Cited by:** 56 **Author:** Mitsushige Sugimoto, Hiromitsu Ban, ...

**Publish Year:** 2017



Comparison of endoscopic gastritis based on Kyoto classification be



ALL

IMAGES

VIDEOS

60,400 Results

Any time ▾

## [Endoscopy-based Kyoto classification score of gastritis ...](#)

<https://pubmed.ncbi.nlm.nih.gov/32982115>

Background: **Endoscopy-based Kyoto classification** for **gastritis** and pathological topographic distribution of neutrophil infiltration are correlated with **gastric cancer** risk. Aim: To investigate the association **between Kyoto classification** and the topographic distribution of neutrophil activity. Methods: **Kyoto classification** score, ranging from 0 to 8, consisted of atrophy, **intestinal metaplasia** ...

**Author:** Osamu Toyoshima, Toshihiro Nishiza... **Publish Year:** 2020

## [Efficacy of the Kyoto Classification of Gastritis in ...](#)

<https://pubmed.ncbi.nlm.nih.gov/28321054>

**Kyoto gastritis** scores of atrophy and **intestinal metaplasia** in the H. pylori-positive **cancer** group were significantly higher than in subjects with **gastritis** alone (all  $p < 0.001$ ). No significant differences were noted in the rates of **gastric** fold hypertrophy or **diffuse** redness **between** the two groups.

**Cited by:** 57 **Author:** Mitsushige Sugimoto, Hiromitsu Ban, Hit...

**Publish Year:** 2017

## [\(PDF\) Kyoto Classification of Gastritis and Gastric Cancer](#)

<https://www.researchgate.net/publication/333751162...>

Jun 10, 2019 · The **Kyoto Classification** of **Gastritis** is useful for diagnosis of H. pylori infection and risk assessment of **gastric cancer** in **endoscopic** practice, and is also expected to be useful for student ...

**Author:** Dong Hyun Lee

## [Association between gastric cancer and the Kyoto ...](#)

<https://onlinelibrary.wiley.com/doi/abs/10.1111/jgh.13764>

From August 2013 to September 2014, esophagogastroduodenoscopy was performed and the **gastric** findings evaluated according to the **Kyoto Classification** of **Gastritis** in a total of 4062



Endoscopy-based Kyoto classification score of gastritis ...

https://pubmed.ncbi.nlm.nih.gov/32982115

Kyoto scores were significantly higher in the order of inactive stomach, antrum-predominant gastritis, pangastritis, and corpus-predominant gastritis (3.05, 4.57, 5.21, and 5.96, respectively). Each individual score of endoscopic findings ( i.e ., atrophy, intestinal metaplasia, enlarged folds, nodularity, and diffuse redness) was correlated with the topographic distribution of neutrophil activity.

Author: Osamu Toyoshima, Toshihiro Nishizaw... Publish Year: 2020

Efficacy of the Kyoto Classification of Gastritis in ...

https://pubmed.ncbi.nlm.nih.gov/28321054

Kyoto gastritis scores of atrophy and intestinal metaplasia in the H. pylori-positive cancer group were significantly higher than in subjects with gastritis alone (all p<0.001). No significant differences were noted in the rates of gastric fold hypertrophy or diffuse redness between the two groups.

Cited by: 58 Author: Mitsushige Sugimoto, Hiromitsu Ban, Hito... Publish Year: 2017

PEOPLE ALSO ASK

- How to identify patients with high risk of gastric cancer?
- How is gastric cancer related to H pylori infection?
- What is Kyoto classification?
- How is B redness related to gastric cancer?

Feedback

(PDF) Endoscopic Findings of Kyoto Classification of Gastritis

https://www.researchgate.net/publication/333750947...

Jun 10, 2019 · The Kyoto Classification of Gastritis is useful for diagnosis of H. pylori infection and risk assessment of gastric cancer in endoscopic practice, and is also expected to be useful for student ...

Author: Gwang Ha Kim

(PDF) Kyoto Classification of Gastritis and Gastric Cancer

https://www.researchgate.net/publication/333751162

Gastritis

Medical Condition

A group of conditions that cause inflammation of the stomach lining.

- Common (More than 200,000 cases per year in US)
- Transmitted through fecal-oral route
- No vaccine available
- Rarely requires lab test or imaging
- Treatment from medical professional advised
- Can last several days or weeks

Gastritis can be caused by underlying conditions, such as infections, medications, excessive alcohol use, Crohn's disease and parasitic infections. Nausea, vomiting and abdominal pain are common symptoms. Treatment includes medication and diet modifications.

Symptoms

The common symptoms include:

- Nausea
- Abdominal bloating
- Abdominal pain
- Vomiting
- Indigestion
- Burning sensation in the stomach
- Hiccups
- Loss of appetite
- Black and tarry stools

Treatments

Treatment is based on the diagnosis. Most of the treatments are aimed at reducing stomach acid and symptoms.

Medication

- Antibiotics: Used to treat bacterial infection. Clarithromycin · Amoxicillin · Metronidazole
- Supplements: Supplements are taken to correct deficiencies of