

国内版国际版

Microsoft Bing

Progressive disfiguring facial mass with pupillary axis obstruction fre

ALL

IMAGES

VIDEOS

15,100 ResultsAny time

Chronic eyelid lymphedema and acne rosacea - Report of ...
<https://www.researchgate.net/publication/12229543...>
Morbihan's disease is classically defined as primarily a Caucasian disease of chronic, recurrent erythema and nonpitting edema of the middle and upper thirds of the face. The disease is often ...

Search Tools
Turn off Hover Translation (关闭取词)

Orbitotemporal Neurofibromatosis: Classification and ...
<https://www.researchgate.net/publication/5751088...>
In orbitofacial NF1 (OFNF), which occurs in 1-22% of patients, NFs may cause progressive, disfiguring tumours of the orbital, facial, and temporal areas. 3, 4 In this study, we review the ...

Extramedullary plasmacytoma in the maxillofacial region: A ...
<https://www.sciencedirect.com/science/article/pii/S221454191830018X>
Dec 01, 2018 The patient reported a 2-month history of a progressive mass in the upper maxillary ridge, with an aggressive pattern of growth. He had no history of nasal obstruction, orbital symptoms, ...
Author: Francisco Merino, Isabel Portillo Hern... Publish Year: 2018

Orbital Decompression in Thyroid Eye Disease - American ...
<https://www.aao.org/oculoplastics-center/orbital...>
Silver RD, Harrison AR, Goding GS. Combined endoscopic medial and external lateral orbital decompression for progressive thyroid eye disease. Otolaryngology-Head and Neck Surgery...

(PDF) Case Files Pathology 2nd Edition [PDF] | David ...

ALL

IMAGES

VIDEOS

3,080 Results

Any time ▾

The Silent Sinus Syndrome: A Case Series and Literature ...

<https://www.researchgate.net/publication/229527813...>

Silent sinus **syndrome** (SSS) is known as a relatively new pathology with clinical characterizations, represented by the **facial** asymmetry determined by **progressive** enophthalmos and hypoglobus and ...

Orbitotemporal Neurofibromatosis: Classification and ...

<https://www.researchgate.net/publication/5751088...>

In OFNF, which occurs in 1-22% of patients, orbitofacial NFs may cause **progressive, disfiguring** tumors of the lid, brow, temple, **face** and orbit [11, 33,48]. Large NFs involving the orbit tend to ...

[DOC] C&P Service Clinician's Guide - Veterans Affairs

www.benefits.va.gov/warms/docs/admin21/guide/cliniciansguide.doc · [Web view](#)

c. Allergic or vasomotor rhinitis: If diagnosis has been established, **report** whether there are nasal polyps and the percent of **obstruction** of nasal passage on each side. d. Chronic laryngitis: **Report** any hoarseness, and describe laryngeal findings, such as inflammation, nodules, polyps of cords, or any other cord abnormalities.

Author: GPalmer

Created Date: 6/13/2002 6:17:00 PM

Last modified by: John

Title: C&P Service Clinician's Guide

Glaucoma and Globe Enlargement Associated with ...

<https://www.researchgate.net/publication/26788319...>

Sep 01, 2009 · **Case Report:** The patient was a seven-year-old boy with unilateral glaucoma secondary to CEH and **facial asymmetry**, mild unilateral ptosis, and proptosis in the left eye.

iThenticate®

62296_Auto_Edited.docx

Quotes Excluded
Bibliography Excluded

2%
0.0000

Match Overview

1

Crossref 15 words
Tsung-Tai Lin, Hsian-He Hsu, Shih-Chun Lee, Yi-Jen Peng, Kai-Hsiung Ko. "Dynamic magnetic resonance imaging feature...

1%

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 62296

Manuscript Type: CASE REPORT

Progressive disfiguring facial mass with pupillary axis obstruction from Morbihan syndrome: A case report

Morbihan syndrome

Abstract

BACKGROUND

Morbihan syndrome is a rare illness, which presents with recurrent pitting edema on the upper two thirds of the face causing facial contour deformities, and Morbihan syndrome obstructing pupillary axis with ptosis was seldomly reported.

CASE SUMMARY

PAGE: 1 OF 6

Text-Only Report

国内版 国际版

Progressive disfiguring facial mass with pupillary axis obstruction fr



ALL IMAGES VIDEOS

6,260 Results

Any time ▾

[Melkersson-Rosenthal syndrome | MedLink Neurology](#)

<https://www.medlink.com/articles/melkersson-rosenthal-syndrome> ▾

Apr 20, 2004 · In a patient with upper facial swelling, an absence of orolabial swelling, and lack of facial neuropathy, a diagnosis of **Morbihan** disease is preferred over Melkersson-Rosenthal **syndrome**, especially if the patient has a history of rosacea (40). **Morbihan** disease is a rare condition characterized by chronic and persistent erythematous solid edema ...

[The Silent Sinus Syndrome: A Case Series and Literature ...](#)

<https://www.researchgate.net/publication/229527813...>

Silent sinus **syndrome** (SSS) is known as a relatively new pathology with clinical characterizations, represented by the facial asymmetry determined by progressive enophthalmos and hypoglobus and ...

[Orbitotemporal Neurofibromatosis: Classification and ...](#)

<https://www.researchgate.net/publication/5751088...>

In OFNF, which occurs in 1-22% of patients, orbitofacial NFs may cause progressive, disfiguring tumors of the lid, brow, temple, face and orbit [11, 33,48]. Large NFs involving the orbit tend to ...

[Management of Congenital Vascular Lesions of the Head ...](#)

<https://www.cancernetwork.com/view/management...> ▾



Hemangiomas

Diagnosis

Treatment

Vascular Malformations



Most hemangiomas first appear soon after birth, but **as many as 30% may be apparent at birth** . They characteristically proliferate during the first year of life and then involute sometime during childhood. Proliferative Phases Although the rate and timing of proliferation within this period is extremely variable, two periods of rapid growth are frequently seen. The first and most common of these proliferative phases takes place during the neonatal period and early infancy and the second, between 4 and 6 months of a...

[See more on cancernetwork.com](#)

[Morbihan's Disease: Treatment with CO 2 Laser ...](#)

https://www.researchgate.net/publication/8545296_Morbihan

Morbihan's disease is a rare condition characterized by chronic persistent erythema and edema of the