

ALL IMAGES VIDEOS

57,100 Results Anytime ▾

[Surgical manifestations of gastrointestinal ...](#)

<https://www.sciencedirect.com/science/article/pii/S0022346815003966>

Nov 01, 2015 · Invasive CMV gastrointestinal disease in our children was predominantly HIV-associated, or followed a major lower gastrointestinal inflammatory insult in infants younger than 6 months. Successful therapy requires a high index of suspicion of active CMV disease to allow early implementation of CMV viral load control and aggressive treatment of the underlying immune impairment.

Cited by: 7 Author: M. Arnold, R. Itzikowitz, B. Young, S.M. ...

Publish Year: 2015

[Cytomegalovirus \(CMV\)-associated portal vein thrombosis in ...](#)

<https://casereports.bmj.com/content/13/12/e238645>

Cytomegalovirus-related thrombosis is a complication of acute cytomegalovirus (CMV) infection. Its presentation was originally associated with an immunocompromised state, but a rising number of case reports in recent years have noted the increasing prevalence of ...

[Cytomegalovirus Enterocolitis Mimicking Necrotizing ...](#)

<https://academic.oup.com/jpids/article/2/1/71/1084746>**Abstract** Case Report Review of The Literature Acknowledgments

Although cytomegalovirus (CMV) colitis is a well-known entity in immune-compromised individuals, such as those infected with the human immunodeficiency virus (HIV), transplant recipients, and patients receiving immunosuppressive therapy [1], it is uncommon in immunocompetent individuals. CMV infection in infancy is generally asymptomatic, but approximately 10% of congenitally infected infants present with rash, jaundice, and microcephaly at birth [2]. Most term infants who acquire CMV through breast milk re...

See more on academic.oup.com

Cited by: 7 Author: Maskit Bar-Meir, Kathryn N. Farrow, Hect...

Publish Year: 2013 Published: Jun 22, 2012

[Gastrointestinal manifestations of postnatal ...](#)

<https://www.researchgate.net/publication/8496911...>

The incidence, clinical characteristics, symptoms, sequelae, and long-term outcome at corrected age [(CA): 2 years of CMV infection] were analyzed. While no infant developed CMV infection with ...

Jejunal diaphragm disease associated with cytomegalovirus ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6447397>

Cytomegalovirus (CMV) infection, usually associated with an immunosuppressed host, can cause symptomatic **gastrointestinal disease** in an immunocompetent patient. Although CMV can affect the entire **gastrointestinal tract**, the colon is most commonly involved, followed by the small bowel 1. Clinical manifestations may range from mild anorexia to obvious **gastrointestinal bleeding** and perforation.

Author: Sumika Kagebayashi, Shuji Yamamoto, ... Publish Year: 2019

Complications of measles: a case series | BMJ Case Reports

<https://casereports.bmj.com/content/13/2/e232408> ▾

Feb 01, 2020 · **Gastrointestinal** complications include measles-associated diarrhoea with **secondary** dehydration and cases of **appendicitis** have also been reported. Characteristic giant cells positive for measles can be seen on immunologic staining of gastric or appendiceal biopsies, which may be present before the emergence of the pathognomonic rash. 9

Cited by: 1 Author: Thelma Xerri, Nicola Darmanin, Maria Ales...

Publish Year: 2020

Neonatal gastrointestinal involvement and congenital ...

<https://www.ncbi.nlm.nih.gov/pubmed/28009139>

Nov 28, 2016 · Very low birth weight and extremely low birth weight newborns are at higher risk of symptomatic **CMV infection**, most commonly **secondary** and acquired through breast milk.

Gastrointestinal involvement is rare in acquired **CMV infections**, but it could be an important manifestation of postnatal infection in preterm infants admitted to neonatal intensive care units.

Cited by: 2 Author: Alessandro Porta, Adriana Avanzini, Marta ...

Publish Year: 2016

Cytomegalovirus induced colonic stricture presenting as

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 62643

Manuscript Type: CASE REPORT

Gastrointestinal cytomegalovirus disease secondary to measles in an immunocompetent infant: A case report

Yang QH *et al.* Infant prolonged fever and bloody diarrhea

Abstract

BACKGROUND

Gastrointestinal (GI) cytomegalovirus (CMV) disease occurs commonly in immunocompromised/immunodeficient patients with advanced HIV infection, neoplasm, solid organ transplantation, hematopoietic stem cell transplantation, or treatment with immunosuppressants, but is rarely reported in association with measles infection.

Match Overview

Match #	Source	Match %
1	Internet 62 words crawled on 03-Dec-2016 www.ncbi.nlm.nih.gov	2%
2	Crossref 33 words William J Moss. "Measles", <i>The Lancet</i> , 2017	1%
3	Crossref 20 words George D. Liatsos, Maria Pirounaki, Angelina Lazareva, Georgia Kikezeou, Spyridon P. Dourakis. "Cytomegalovirus infect..."	1%
4	Crossref 17 words Dongang Dai, Feixiu Wen, Shaoming Zhou, Zhe Su, Guosheng Liu, Mingbang Wang, Jiantai Zhou, Fusheng He. "Association	1%
5	Internet 17 words crawled on 24-Feb-2020 www.wjgnet.com	1%
6	Crossref 17 words Diane E Griffin. "Measles immunity and immunosuppression", <i>Current Opinion in Virology</i> , 2021	1%

国内版 国际版

Gastrointestinal cytomegalovirus disease secondary to measles in ar



ALL IMAGES VIDEOS

54,600 Results Any time ▾

Jejunal diaphragm disease associated with cytomegalovirus ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6447397>

Cytomegalovirus (CMV) infection, usually associated with an immunosuppressed host, can cause symptomatic **gastrointestinal disease** in an immunocompetent patient. Although CMV can affect the entire **gastrointestinal tract**, the colon is most commonly involved, followed by the small bowel 1. Clinical manifestations may range from mild anorexia to obvious **gastrointestinal bleeding** and perforation.

Author: Sumika Kagebayashi, Shuji Yamamot... Publish Year: 2019

Neonatal gastrointestinal involvement and congenital ...

<https://www.ncbi.nlm.nih.gov/pubmed/28009139>

Nov 28, 2016 · Very low birth weight and extremely low birth weight newborns are at higher risk of symptomatic **CMV infection**, most commonly **secondary** and acquired through breast milk. **Gastrointestinal involvement** is rare in acquired **CMV infections**, but it could be an important manifestation of postnatal infection in preterm **infants** admitted to neonatal intensive care units.

Cited by: 2 Author: Alessandro Porta, Adriana Avanzini, Mart...
Publish Year: 2016

Complications of measles: a case series | BMJ Case Reports

<https://casereports.bmj.com/content/13/2/e232408> ▾

Feb 01, 2020 · **Gastrointestinal complications** include measles-associated diarrhoea with **secondary dehydration** and cases of **appendicitis** have also been reported. Characteristic giant cells positive for **measles** can be seen on immunologic staining of gastric or appendiceal biopsies, which may be present before the emergence of the pathognomonic rash. 9

Cited by: 1 Author: Thelma Xerri, Nicola Darmanin, Maria Ale...
Publish Year: 2020

Cytomegalovirus-induced colonic stricture presenting as ...

<https://europepmc.org/article/PMC/PMC3794276> ▾

Sep 10, 2013 · **Gastrointestinal cytomegalovirus (CMV) infection** can have serious morbidity and mortality and it occurs as an opportunistic **infection** in **immunocompromised patients**. Fortunately, it is very rare in **immunocompetent patient** where it is usually mild and self-limiting. But it rarely induces serious complications in an **immunocompetent adult**.

Cited by: 6 Author: B V Dinesh, Karthikeyan Selvaraju, Samp...