

国内版

国际版

Celiac disease in Asia beyond Middle East and Indian subcontinent:



ALL

IMAGES

VIDEOS

57,500 Results

Any time ▾

Genetic predisposition to celiac disease in Kazakhstan ...

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0226546> ▾

Jan 02, 2020 - Background Celiac disease (CD) is a systemic immune-mediated disorder developing in HLA genetically predisposed individuals carrying HLA-DQ2 and/or HLA-DQ8 molecules. Recent evidences supported a predominant importance of HLA-DQB1 locus and, in particular, HLA-DQB1*02 alleles. This diagnosis is poorly considered in Kazakhstan, because of the assumption that CD is not ...

Cited by: 3

Author: Dimitri Poddighe, Aida Turganbekova, Da...

Publish Year: 2020

Prevalence of Celiac Disease in Shiraz, Southern Iran

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2702920>

Celiac disease (CD), also called gluten-sensitive enteropathy and nontropical sprue, is a common autoimmune disorder characterized by an immune response to ingested gluten.[] Until a few years ago, this disorder was considered to be almost exclusively affecting the people of European origin[]; however, recent reports from other regions of the world revealed similar rates of disease when ...

Cited by: 21

Author: Mehdi Saberi-Firouzi, Gholamhossein R ...

Publish Year: 2008

PEOPLE ALSO ASK

What is the incidence of cancer in Asia? ▾

Is there a cancer burden in Asia? ▾

Is asia a diverse continent? ▾

What diseases are caused by Aedes aegypti? ▾

Feedback

Demographic and Epidemiologic Drivers of Global ...

<https://www.nejm.org/doi/full/10.1056/NEJMoa1406656>

According to the Global Burden of Disease Study, between 1990 and 2013, global cardiovascular mortality increased by 55% as a result of aging and by 25% as a ...



ALL IMAGES VIDEOS

62,600 Results Any time ▾

Genetic predisposition to celiac disease in Kazakhstan ...

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0226546> ▾

Jan 02, 2020 · Celiac disease (CD) is a systemic immune-mediated disorder developing in HLA genetically predisposed individuals carrying HLA-DQ2 and/or HLA-DQ8 molecules. Recent evidences supported a predominant importance of HLA-DQB1 locus and, in particular, HLA-DQB1*02 alleles. This diagnosis is poorly considered in Kazakhstan, because of the assumption that CD is not prevalent in ...

Cited by: 4 Author: Dimitri Poddighe, Aida Turganbekova, Dana...
Publish Year: 2020

Prevalence of Celiac Disease in Shiraz, Southern Iran

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2702920>

Celiac disease (CD), also called gluten-sensitive enteropathy and nontropical sprue, is a common autoimmune disorder characterized by an immune response to ingested gluten.[] Until a few years ago, this disorder was considered to be almost exclusively affecting the people of European origin[]; however, recent reports from other regions of the world revealed similar rates of disease when ...

Cited by: 21 Author: Mehdi Saberi-Firouzi, Gholamhossein R O...
Publish Year: 2008

PEOPLE ALSO ASK

- What is the incidence of cancer in Asia? ▾
- Is there a cancer burden in Asia? ▾
- What is the global burden of Disease Study? ▾

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 62726

Manuscript Type: EDITORIAL

Celiac disease in Asia beyond Middle East and Indian subcontinent: Epidemiological burden and diagnostic barriers

Celiac disease in Asia

Abstract

Celiac Disease had been considered uncommon in Asia for a long time. However, several studies suggested that, in the Indian subcontinent and Middle East countries, Celiac Disease is present and as prevalent as in Western countries. Outside these Asian regions, the information about the epidemiology of Celiac Disease is still lacking or largely incomplete, because of different and variable reasons. Here, we discuss the epidemiological aspects and the diagnostic barriers in the following Asian regions: China, Japan, South East Asia and Russia/Central Asia. In some of these region (and in

Match Overview

1 **Internet** 33 words
crawled on 16-Jan-2020
www.mdpi.com 1%

2 **Crossref** 13 words
Chu-Nan Chen, Jing-Nan Li. "Insufficient awareness of celiac disease in China", *Chinese Medical Journal*, 2019 1%

国内版 国际版

Celiac disease in Asia beyond Middle East and Indian subcontinent:



ALL IMAGES VIDEOS

59,500 Results

Any time ▾

Genetic predisposition to celiac disease in Kazakhstan ...

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0226546> ▾

Jan 02, 2020 · Celiac disease (CD) is a systemic immune-mediated disorder developing in HLA genetically predisposed individuals carrying HLA-DQ2 and/or HLA-DQ8 molecules. Recent evidences supported a predominant importance of HLA-DQB1 locus and, in particular, HLA-DQB1*02 alleles. This diagnosis is poorly considered in Kazakhstan, because of the assumption that CD is not prevalent in ...

Cited by: 4

Author: Dimitri Poddighe, Aida Turganbekova, Da...

Publish Year: 2020

Prevalence of Celiac Disease in Shiraz, Southern Iran

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2702920>

Celiac disease (CD), also called gluten-sensitive enteropathy and nontropical sprue, is a common autoimmune disorder characterized by an immune response to ingested gluten.[] Until a few years ago, this disorder was considered to be almost exclusively affecting the people of European origin[]; however, recent reports from other regions of the world revealed similar rates of disease when ...

Cited by: 21

Author: Mehdi Saberi-Firouzi, Gholamhossein R ...

Publish Year: 2008

Primary Prevention of Celiac Disease: Environmental ...

<https://www.nestlenutrition-institute.org/...> ▾

Celiac disease affects around 1% of the general population in Europe, greatly affecting patients' quality of life and incurring a significant cost burden on society. There has always been a great deal of inconsistency in the literature regarding the impact of the timing of gluten introduction on the risk of celiac disease.

Leishmaniasis - The Lancet

[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(18\)31204-2/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(18)31204-2/fulltext)

Leishmaniasis is a poverty-related disease with two main clinical forms: visceral leishmaniasis and cutaneous leishmaniasis. An estimated 0.7–1 million new cases of leishmaniasis per year are reported from nearly 100 endemic countries. The number of reported visceral leishmaniasis cases has decreased substantially in the past decade as a result of better access to diagnosis and treatment ...

Ethnicity and cardiovascular disease prevention in the ...

<https://www.nature.com/articles/11002126>