

4,760 Results

Any time ▾

[PDF] Changing TACTICS in intermediate HCC: TACE plus ...<https://gut.bmj.com/content/gutjnl/69/8/1374.full.pdf>

intermediate-stage HCC (Barcelona Clinic Liver Cancer (BCLC) stage B), despite the fact that this stage comprises patients with a wide range of liver function, variable tumour number and size. 1–4 In clinical practice, only 50%–60% of patients with BCLC stage B benefit from TACE, so TACE is usually repeated to achieve maximum tumour recession.

Author: Pompilia Radu, Jean-François Dufour Publish Year: 2020

The Barcelona approach: Diagnosis, staging, and treatment ...<https://aasldpubs.onlinelibrary.wiley.com/doi/full/10.1002/lt.20034>

Jan 30, 2004 · Barcelona-Clinic Liver Cancer (BCLC) staging classification and treatment schedule.

Stage 0: Patients with very early HCC are optimal candidates for resection. Stage A: Patients with early HCC are candidates for radical therapies (resection and ablation, liver ...

Cited by: 771

Author: Josep M. Llovet, Josep Fuster, Jordi Bruix

Publish Year: 2004

(PDF) Intermediate stage hepatocellular carcinoma: a ...<https://www.researchgate.net/publication/334389958...>

Intermediate stage or Barcelona Clinic Liver Cancer (BCLC) stage B HCC includes multiple tumorous lesions confined to the liver without vascular invasion in a patient with preserved liver ...

Ascites and alpha-fetoprotein improve prognostic ...<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4427691>

May 14, 2015 · Core tip: Although the Barcelona Clinic Liver Cancer (BCLC) system contains Child-Turcotte-Pugh classification as a main variable, a patient within BCLC stage A or B may have ascites, which can interfere with the recommended treatment. An elevated alpha-fetoprotein level, which is an integral part of other staging systems, may also impact prognosis and influence treatment decision.

Cited by: 3

Author: Asmaa I Gomaa, Alzhraa Al-Khatib, Wael...

Publish Year: 2015

Prognosis of patients with intermediate-stage ...https://www.researchgate.net/publication/265609669_Prognosis_of_patients_with...

Intermediate stage HCC, as defined by the Barcelona Clinic Liver Cancer (BCLC) system stage B,

国内版

国际版

Rethinking the Barcelona Clinic liver cancer guidelines: Intermediate



ALL

IMAGES

VIDEOS

4,640 Results

Any time ▾

[Treatment algorithm for intermediate and advanced stage ...](#)

<https://pubmed.ncbi.nlm.nih.gov/22212948>

The practice **guideline** for hepatocellular carcinoma (HCC) in Korea was revised in 2009. It was based on clinical evidence. The treatment algorithm was divided into curative and noncurative treatments.

According to **Barcelona Clinic Liver Cancer** (BCLC) staging, the curative treatment group included ea

...

Cited by: 23

Author: Jong Young Choi

Publish Year: 2011

[\[PDF\] Changing TACTICS in intermediate HCC: TACE plus ...](#)

<https://gut.bmj.com/content/gutjnl/early/2020/03/13/gutjnl-2020-320692.full.pdf>

Mar 13, 2020 · line treatment for **intermediate- stage** HCC (**Barcelona Clinic Liver Cancer** (BCLC) **stage B**), despite the fact that this **stage** comprises **patients** with a wide range of **liver** function, variable tumour number and size.^{1–4} In clinical practice, only 50%–60% of **patients** with BCLC **stage B** benefit from TACE, so TACE is usually

Author: Pompilia Radu, Jean-François Dufour **Publish Year:** 2020

iThenticate®

63093_Auto_Edited.docx

Quotes Excluded
Bibliography Excluded

12%
Similarity

13

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 63093

Manuscript Type: MINIREVIEWS

Rethinking the Barcelona Clinic liver cancer guidelines: Intermediate stage and Child-Pugh B patients are suitable for surgery?

Fabrizio Romano, Marco Chiarelli, Mattia Garancini, Mauro Scotti, Mauro Zago, Gerardo Cioffi, Matilde De Simone, Ugo Cioffi

Match Overview

1

Internet 77 words
crawled on 04-Sep-2020
synapse.koreamed.org

2%

2

Crossref 63 words
Giammauro Berardi, Zenichi Morise, Carlo Sposito, Kazuharu Igarashi et al. "Development of a nomogram to predic ...

2%

3

Internet 50 words
crawled on 22-Apr-2020
journals.lww.com

1%

4

Crossref 36 words
G.K. Gkantounis, A. Palkouras, M.-C. Stylianidi, H. Milonis, P. Tzimas, D. Roukos, G. Pentheroudakis, E. Felekouras. *

1%

5

Crossref 33 words
Philipp Renner, Jürgen Schuhbaum, Alexander Kroemer, Florian Zeman et al. "Morbidity of hepatic resection for inte: ...

1%

6

Internet 27 words
crawled on 16-Sep-2020

1%

国内版 国际版

Rethinking the Barcelona clinic liver cancer guidelines: Intermediate



ALL IMAGES VIDEOS

5,000 Results

Any time ▾

Treatment algorithm for intermediate and advanced stage ...

<https://pubmed.ncbi.nlm.nih.gov/22212948>

The practice guideline for hepatocellular carcinoma (HCC) in Korea was revised in 2009. It was based on clinical evidence. The treatment algorithm was divided into curative and noncurative treatments. According to Barcelona Clinic Liver Cancer (BCLC) staging, the curative treatment group included ea ...

Cited by: 23

Author: Jong Young Choi

Publish Year: 2011

[PDF] Changing TACTICS in intermediate HCC: TACE plus ...

<https://gut.bmj.com/content/gutjnl/early/2020/03/13/gutjnl-2020-320692.full.pdf>

Mar 13, 2020 · line treatment for intermediate- stage HCC (Barcelona Clinic Liver Cancer (BCLC) stage B), despite the fact that this stage comprises patients with a wide range of liver function, variable tumour number and size.1–4 In clinical practice, only 50%–60% of patients with BCLC stage B benefit from TACE, so TACE is usually

Author: Pompilia Radu, Jean-François Dufour Publish Year: 2020

PEOPLE ALSO ASK

What are the stages of liver cancer in Barcelona? ▾

What is HCC in Child-Pugh Class B liver function? ▾

What are the guidelines for the diagnosis of hepatocellular carcinoma? ▾

Which is the best prognosis system for liver cancer? ▾

Feedback

[PDF] Changing TACTICS in intermediate HCC: TACE plus ...

<https://gut.bmj.com/content/gutjnl/69/8/1374.full.pdf>

intermediate-stage HCC (Barcelona Clinic Liver Cancer (BCLC) stage B), despite the fact that this stage comprises patients with a wide range of liver function, variable tumour number and size. 1–4 In clinical prac-tice, only 50%–60% of patients with BCLC stage B benefit from TACE, so TACE is usually repeated to achieve maximum tumour recession.