

[Clinicopathological features and outcome for ...](#)

<https://pubmed.ncbi.nlm.nih.gov/30062861>

GEJ-NENs were mostly poorly differentiated carcinomas, and all of Chinese patients were NEC/MANEC. The outcome of MANEC was preferable to NECs. Both lymph nodes metastasis and distant disease were independent predictors of prognosis. Surgical resection can improve survival, but postoperative adjuvan ...

Cited by: 5 Author: Panpan Zhang, Wei Wang, Ming Lu, Cha...
Publish Year: 2018

[Clinicopathological significance of gastric poorly ...](#)

<https://pubmed.ncbi.nlm.nih.gov/27108877>

Clinicopathological characteristics, prognosis, and ...

<https://www.sciencedirect.com/science/article/pii/S1078143920302775>

Jan 01, 2021 · There were 232 patients (92.8%) with pathological stage T2 to T4 diseases and 56 patients (22.4%) initially diagnosed with stage IV disease. The most common sites of metastasis were lung (39.6%), distant lymph nodes (39.2%), bone (19.6%), liver (18.0%), and adrenal gland (7.2%), respectively, and 10.4% of patients appeared local recurrence.

Author: Xu Li, Siming Li, Zhihong Chi, Chuanli... Publish Year: 2021

Analysis of clinicopathological characteristics associated ...

<https://www.spandidos-publications.com/10.3892/ol.2016.5120>

Clinical characteristics. Clinicopathological parameters of 232 patients with OSCC, including 175 fatal cases and 57 non-fatal cases, were analyzed. The 232 patients with OSCC included 169 males and 63 females, with an age range of 24–83 years. The 57 non-fatal cases included 37 males and 20 females, of which 26 patients were aged >60 years.

Cited by: 1 Author: Xin Jin, Dongxuan Liu, Xin Zhao, Yu Zhou, ... Publish Year: 2016

[Comparison of clinicopathological characteristics and ...

<https://www.ncbi.nlm.nih.gov/pubmed/21186408>

Search Tools

Turn off Hover Translation (关闭翻译)

激活 Windows
转到“设置”以激活 Windows。

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 63158

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Analysis of the clinicopathological characteristics and prognosis of 232 patients with poorly differentiated gastric neuroendocrine neoplasms

Poorly differentiated gastric neuroendocrine neoplasms

Deng Han, Yuang-Liang Li, Zhi-Wei Zhou, Fei Yin, Jie Chen, Fang Liu, Yanfen Shi, Wei Wang, Yu Zhang, Xianjun Yu, Jian-Ming Xu, Runxiang Yang, Chao Tian, Jie Luo, Huang-Ying Tan

Match Overview

Rank	Source	Words	Match %
1	Crossref "Abstracts", Journal of Neuroendocrinology, 2021	49 words	1%
2	Internet crawled on 17-Jul-2020 www.wjgnet.com	32 words	1%
3	Internet crawled on 06-Jan-2021 pesquisa.bvsalud.org	28 words	1%
4	Crossref "Abstracts", Neuroendocrinology, 2019	27 words	1%
5	Internet crawled on 01-Dec-2020 www.researchsquare.com	15 words	<1%
6	Crossref Rui Mao, Kan Li, Jian-Qiang Cai, Sheng Luo, Megan Turner, Dan Blazer, Hong Zhao, Adjuvant Chemotherapy We	15 words	<1%

国内版 国际版

Analysis of the clinicopathological characteristics and prognosis of 2



ALL IMAGES VIDEOS

57,200 Results Any time ▾

Multicenter Analysis of the Clinicopathological ...

<https://www.enets.org/multicenter-analysis-of-the...>

#2683 Clinicopathological Features and Prognostic Analysis of 227 Cases with Well-Differentiated Gastric Neuroendocrine Tumors Introduction: Gastric neuroendocrine neoplasms (g-NENs) are a group of rare tumors with a strong heterogeneity, subdivided as well-differentiated neuroendocrine tumors (NETs, type 1-3 g-NETs) and poorly-differentiated ...

Clinicopathological features and outcome for ...

<https://pubmed.ncbi.nlm.nih.gov/30062861>

GEJ-NENs were mostly poorly differentiated carcinomas, and all of Chinese patients were NEC/MANEC. The outcome of MANEC was preferable to NECs. Both lymph nodes metastasis and distant disease were independent predictors of prognosis. Surgical resection can improve survival, but postoperative adjuvan ...

Cited by: 5 Author: Panpan Zhang, Wei Wang, Ming Lu, Cha...

Publish Year: 2018

Clinicopathological characteristics and prognosis of ...

<https://www.researchgate.net/publication/5258230...>

To clarify the oncological outcome of rectal well-differentiated neuroendocrine tumors (W/D NETs), we examined the clinicopathological characteristics and prognosis of patients with this neoplasm.

Clinicopathological features and prognosis of patients ...

<https://onlinelibrary.wiley.com/doi/full/10.1002/cam4.1683>

Oct 11, 2018 · Neuroendocrine tumors (NETs) are a type of tumor derived from the diffuse neuroendocrine system, and they account for 1%-2% of malignant tumors. 1 Gastric neuroendocrine tumors (G-NETs) are rare, with prevalences of 3.2 and 1.7 per 100 000 people in European countries and the United States, respectively. 2

Cited by: 4 Author: Fang-Xi Tian, Yu-Qing Cai, Lv-Ping Zhuan...

Publish Year: 2018

Clinicopathological features and prognosis factors for ...

<https://journals.lww.com/md-journal/Fulltext/2019/...>

study was analysis of the data from Surveillance Epidemiology and End Results database. A total of 4608 patients with "one primary only" histologically pNETs were confirmed and 653 were older than 75